

Two group intervals

Brian Caffo, Jeff Leek, Roger Peng

May 19, 2016

Independent group t confidence intervals

- ▶ Suppose that we want to compare the mean blood pressure between two groups in a randomized trial; those who received the treatment to those who received a placebo
- ▶ We cannot use the paired t test because the groups are independent and may have different sample sizes
- ▶ We now present methods for comparing independent groups

Notation

- ▶ Let X_1, \dots, X_{n_x} be iid $N(\mu_x, \sigma^2)$
- ▶ Let Y_1, \dots, Y_{n_y} be iid $N(\mu_y, \sigma^2)$
- ▶ Let \bar{X} , \bar{Y} , S_x , S_y be the means and standard deviations
- ▶ Using the fact that linear combinations of normals are again normal, we know that $\bar{Y} - \bar{X}$ is also normal with mean $\mu_y - \mu_x$ and variance $\sigma^2(\frac{1}{n_x} + \frac{1}{n_y})$
- ▶ The pooled variance estimator

$$S_p^2 = \{(n_x - 1)S_x^2 + (n_y - 1)S_y^2\} / (n_x + n_y - 2)$$

is a good estimator of σ^2

Note

- ▶ The pooled estimator is a mixture of the group variances, placing greater weight on whichever has a larger sample size
- ▶ If the sample sizes are the same the pooled variance estimate is the average of the group variances
- ▶ The pooled estimator is unbiased

$$\begin{aligned} E[S_p^2] &= \frac{(n_x - 1)E[S_x^2] + (n_y - 1)E[S_y^2]}{n_x + n_y - 2} \\ &= \frac{(n_x - 1)\sigma^2 + (n_y - 1)\sigma^2}{n_x + n_y - 2} \end{aligned}$$

- ▶ The pooled variance estimate is independent of $\bar{Y} - \bar{X}$ since S_x is independent of \bar{X} and S_y is independent of \bar{Y} and the groups are independent

Result

- ▶ The sum of two independent Chi-squared random variables is Chi-squared with degrees of freedom equal to the sum of the degrees of freedom of the summands
- ▶ Therefore

$$(n_x + n_y - 2)S_p^2/\sigma^2 = (n_x - 1)S_x^2/\sigma^2 + (n_y - 1)S_y^2/\sigma^2$$

$$= \chi_{n_x-1}^2 + \chi_{n_y-1}^2$$

$$= \chi_{n_x+n_y-2}^2$$

Putting this all together

- ▶ The statistic

$$\frac{\frac{\bar{Y} - \bar{X} - (\mu_y - \mu_x)}{\sigma \left(\frac{1}{n_x} + \frac{1}{n_y} \right)^{1/2}}}{\sqrt{\frac{(n_x + n_y - 2) S_p^2}{(n_x + n_y - 2) \sigma^2}}} = \frac{\bar{Y} - \bar{X} - (\mu_y - \mu_x)}{S_p \left(\frac{1}{n_x} + \frac{1}{n_y} \right)^{1/2}}$$

is a standard normal divided by the square root of an independent Chi-squared divided by its degrees of freedom

- ▶ Therefore this statistic follows Gosset's t distribution with $n_x + n_y - 2$ degrees of freedom
- ▶ Notice the form is (estimator - true value) / SE

Confidence interval

- ▶ Therefore a $(1 - \alpha) \times 100\%$ confidence interval for $\mu_y - \mu_x$ is

$$\bar{Y} - \bar{X} \pm t_{n_x+n_y-2, 1-\alpha/2} S_p \left(\frac{1}{n_x} + \frac{1}{n_y} \right)^{1/2}$$

- ▶ Remember this interval is assuming a constant variance across the two groups
- ▶ If there is some doubt, assume a different variance per group, which we will discuss later

Example

Based on Rosner, Fundamentals of Biostatistics


- ▶ Comparing SBP for 8 oral contraceptive users versus 21 controls
- ▶ $\bar{X}_{OC} = 132.86$ mmHg with $s_{OC} = 15.34$ mmHg
- ▶ $\bar{X}_C = 127.44$ mmHg with $s_C = 18.23$ mmHg
- ▶ Pooled variance estimate

```
sp <- sqrt((7 * 15.34^2 + 20 * 18.23^2) / (8 + 21 - 2))  
132.86 - 127.44 + c(-1, 1) * qt(.975, 27) * sp * (1 / 8 + 1 / 21)
```

```
## [1] -9.521097 20.361097
```

```
data(sleep)  
x1 <- sleep$extra[sleep$group == 1]  
x2 <- sleep$extra[sleep$group == 2]  
n1 <- length(x1)
```


Ignoring pairing



Unequal variances

- ▶ Under unequal variances

$$\bar{Y} - \bar{X} \sim N\left(\mu_y - \mu_x, \frac{s_x^2}{n_x} + \frac{\sigma_y^2}{n_y}\right)$$

- ▶ The statistic

$$\frac{\bar{Y} - \bar{X} - (\mu_y - \mu_x)}{\left(\frac{s_x^2}{n_x} + \frac{\sigma_y^2}{n_y}\right)^{1/2}}$$

approximately follows Gosset's t distribution with degrees of freedom equal to

$$\frac{\left(s_x^2/n_x + s_y^2/n_y\right)^2}{\left(\frac{s_x^2}{n_x}\right)^2/(n_x - 1) + \left(\frac{s_y^2}{n_y}\right)^2/(n_y - 1)}$$

Example

- ▶ Comparing SBP for 8 oral contraceptive users versus 21 controls
- ▶ $\bar{X}_{OC} = 132.86$ mmHg with $s_{OC} = 15.34$ mmHg
- ▶ $\bar{X}_C = 127.44$ mmHg with $s_C = 18.23$ mmHg
- ▶ $df = 15.04$, $t_{15.04, .975} = 2.13$
- ▶ Interval

$$132.86 - 127.44 \pm 2.13 \left(\frac{15.34^2}{8} + \frac{18.23^2}{21} \right)^{1/2} = [-8.91, 19.75]$$

- ▶ In R, `t.test(..., var.equal = FALSE)`

Comparing other kinds of data

- ▶ For binomial data, there's lots of ways to compare two groups
- ▶ Relative risk, risk difference, odds ratio.
- ▶ Chi-squared tests, normal approximations, exact tests.
- ▶ For count data, there's also Chi-squared tests and exact tests.
- ▶ We'll leave the discussions for comparing groups of data for binary and count data until covering glms in the regression class.
- ▶ In addition, Mathematical Biostatistics Boot Camp 2 covers many special cases relevant to biostatistics.