

Internship Report in R

Internship Report

```
rm(list=ls())
```

Importing Packages

Data Preprocessing Steps

```
##### Read the data in #####
data <- read.csv(file='insurance.csv')

##### Print the first rows #####
print(head(data, 5))
```

```
##   age    sex    bmi children smoker   region   charges
## 1  19 female 27.900         0    yes southwest 16884.924
## 2  18  male 33.770         1    no  southeast  1725.552
## 3  28  male 33.000         3    no  southeast  4449.462
## 4  33  male 22.705         0    no northwest 21984.471
## 5  32  male 28.880         0    no northwest  3866.855
```

```
##### Print the columns' names #####
print(colnames(data))
```

```
## [1] "age"      "sex"      "bmi"      "children" "smoker"   "region"   "charges"
```

```
##### Print number of rows #####
print(nrow(data))
```

```
## [1] 1338
```

```
##### Converting to Numeric Variables #####
sex <- ifelse(data["sex"] == "female", 0, 1)
smoker <- ifelse(data["smoker"] == "yes", 1, 0)
region <- as.numeric(data$region)
```

```
##### Replacing columns in the Data #####
data["sex"] <- sex
data["smoker"] <- smoker
data["region"] <- region
```

Linear Models - using the purrr package to get individual models

```
##### Linear Regression #####
vars = c('age', 'sex', 'bmi', 'children', 'smoker', 'region')
#Using the purrr package to run all the models corresponding to the predictors
models <- vars %>% paste ('charges ~', .) %>% map(as.formula) %>% map(lm, data = data)
```

Summaries of the Models

Age

```
# age summary
summary(models[[1]])

##
## Call:
## .f(formula = .x[[i]], data = ..1)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -8059  -6671  -5939   5440  47829
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)   3165.9      937.1    3.378 0.000751 ***
## age           257.7       22.5   11.453 < 2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 11560 on 1336 degrees of freedom
## Multiple R-squared:  0.08941,    Adjusted R-squared:  0.08872
## F-statistic: 131.2 on 1 and 1336 DF,  p-value: < 2.2e-16
```

Sex

```
# sex summary
summary(models[[2]])

##
## Call:
## .f(formula = .x[[i]], data = ..1)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -12835  -8435  -3980   3476  51201
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)  12569.6      470.1   26.740 <2e-16 ***
## sex          1387.2      661.3    2.098  0.0361 *
##
```

```
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 12090 on 1336 degrees of freedom
## Multiple R-squared:  0.003282,    Adjusted R-squared:  0.002536
## F-statistic:    4.4 on 1 and 1336 DF,  p-value: 0.03613
```

BMI

```
# bmi summary
summary(models[[3]])
```

```
##
## Call:
## .f(formula = .x[[i]], data = ..1)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -20956  -8118  -3757   4722  49442
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)  1192.94    1664.80   0.717   0.474
## bmi           393.87     53.25   7.397 2.46e-13 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 11870 on 1336 degrees of freedom
## Multiple R-squared:  0.03934,    Adjusted R-squared:  0.03862
## F-statistic: 54.71 on 1 and 1336 DF,  p-value: 2.459e-13
```

Children

```
# children summary
summary(models[[4]])
```

```
##
## Call:
## .f(formula = .x[[i]], data = ..1)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -11585  -8759  -4071   3468  51248
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)  12522.5     446.5  28.049  <2e-16 ***
## children       683.1      274.2   2.491  0.0129 *
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 12090 on 1336 degrees of freedom
## Multiple R-squared:  0.004624,    Adjusted R-squared:  0.003879
## F-statistic: 6.206 on 1 and 1336 DF,  p-value: 0.01285
```

Smoker

```
# smoker summary
summary(models[[5]])
```

```
##
## Call:
## .f(formula = .x[[i]], data = ..1)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -19221  -5042   -919    3705   31720
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)   8434.3      229.0    36.83  <2e-16 ***
## smoker       23616.0      506.1    46.66  <2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 7470 on 1336 degrees of freedom
## Multiple R-squared:  0.6198, Adjusted R-squared:  0.6195
## F-statistic: 2178 on 1 and 1336 DF, p-value: < 2.2e-16
```

Region

```
# region summary
summary(models[[6]])
```

```
##
## Call:
## .f(formula = .x[[i]], data = ..1)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -12116  -8517  -3930   3347   50533
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept) 13441.60     823.85  16.316  <2e-16 ***
## region       -68.04     299.86  -0.227    0.821
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 12110 on 1336 degrees of freedom
## Multiple R-squared:  3.854e-05, Adjusted R-squared: -0.0007099
## F-statistic: 0.05149 on 1 and 1336 DF, p-value: 0.8205
```

Linear Model with All Predictors

```
##### Model with all the predictors #####
allpreds <- lm(charges ~ ., data = data)
```

Summary of the Model

```
##### Summary #####
summary(allpreds)

##
## Call:
## lm(formula = charges ~ ., data = data)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -11343  -2807  -1017   1408  29752
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept) -11461.81     983.00  -11.660  < 2e-16 ***
## age          257.29       11.89   21.647  < 2e-16 ***
## sex         -131.11       332.81   -0.394  0.693681
## bmi          332.57       27.72   11.997  < 2e-16 ***
## children     479.37       137.64    3.483  0.000513 ***
## smoker      23820.43       411.84   57.839  < 2e-16 ***
## region      -353.64       151.93   -2.328  0.020077 *
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 6060 on 1331 degrees of freedom
## Multiple R-squared:  0.7507, Adjusted R-squared:  0.7496
## F-statistic: 668.1 on 6 and 1331 DF,  p-value: < 2.2e-16
```

Linear Model with the Most Relevant Predictors

```
most_rel <- lm(charges ~ age + bmi + children + smoker, data = data)
```

Summary of the Model

```
summary(most_rel)
```

```
##
## Call:
## lm(formula = charges ~ age + bmi + children + smoker, data = data)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -11897.9  -2920.8   -986.6   1392.2  29509.6
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept) -12102.77     941.98  -12.848  < 2e-16 ***
## age           257.85       11.90   21.675  < 2e-16 ***
## bmi           321.85       27.38   11.756  < 2e-16 ***
## children      473.50      137.79    3.436 0.000608 ***
## smoker       23811.40     411.22   57.904  < 2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 6068 on 1333 degrees of freedom
## Multiple R-squared:  0.7497, Adjusted R-squared:  0.7489
## F-statistic: 998.1 on 4 and 1333 DF,  p-value: < 2.2e-16
```

For Comparison with Python Models (Links)

GitHub Pages for the Internship | GitHub Repository
 Heroku App - using Dash and Plotly