

Arbeit zur Erlangung des akademischen Grades  
Bachelor of Science

**Search for single top-quark production  
in the s-channel with the ATLAS  
experiment**

Alfredo Manente  
geboren in Dortmund

2025

Lehrstuhl für Experimentelle Physik IV  
Fakultät Physik  
Technische Universität Dortmund

Erstgutachter: PD Dr. Andrea Helen Knue  
Zweitgutachter: Dr. Salvatore La Cagnina  
Abgabedatum: 15. Dezember 2025

## Kurzfassung

In dieser Arbeit werden Studien zur Messung des Wirkungsquerschnitts für die Erzeugung einzelner Top-Quarks im s-Kanal bei Proton-Proton Kollisionen mit einer Schwerpunktsenergie von  $\sqrt{s} = 13 \text{ TeV}$  durchgeführt. Die Daten stammen aus Kollisionsexperimenten des LHC und wurden vom ATLAS Detektor in den Jahren 2015 bis 2018 aufgenommen. Zur Extrahierung des Wirkungsquerschnitts wird ein Profile Likelihood Fit verwendet, wobei als Diskriminante die Ergebnisse eines *Deep Neural Networks*(DNN) fungieren. Die Signal Signatur besteht aus einem geladenen Lepton, entweder Elektron oder Myon, zwei b-tagged Jets und fehlendem transversalen Impuls. Eine Vielzahl an von kinematischen Variablen wird hinsichtlich ihrer Trennkraft zwischen Signal und Untergrund untersucht.

Was mache ich denn eigentlich genau in meiner Arbeit? Was könnte ich noch dazu schreiben?

- Run 2 Daten
- integrierte Luminosität
- Wie funktioniert die DNN? Es nutzt high-level kinematic variables. Was ist das überhaupt?
- Mache deutlicher dass ich simulierte Daten verwende
- Erkenntnisse aus dem Resultat vielleicht hier noch mit rein schreiben?
- Welche Uncertainties behindern uns am meisten? Background modelling, signal modelling, detector modelling
- kleiner Wirkungsquerschnitt des s-kanals und großer hintergrund bei ähnlichen signaturen
- s kanal bis jetzt nicht präzise gemessen, andere single top channel schon präzise genug gemessen
- Machen wir einen Asimov Fit? Was ist das genau?

## Abstract

The abstract is a short summary of the thesis in English, together with the German summary it has to fit on this page.

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Properties of the Top Quark within the Standard Model</b>	<b>2</b>
2.1	Overview of the Standard Model . . . . .	2
2.2	The Top Quark . . . . .	4
<b>3</b>	<b>detector</b>	<b>7</b>
<b>4</b>	<b>object definition and event selection</b>	<b>9</b>
4.1	Monte Carlo Samples . . . . .	9
4.2	object definition . . . . .	10
4.3	event selection . . . . .	10
<b>5</b>	<b>main study and results</b>	<b>11</b>
5.1	Analysis strategy . . . . .	11
5.2	DNN Model variables . . . . .	11
5.3	Separation power . . . . .	11
5.4	profile likelihood fit . . . . .	11
5.5	systematic uncertainties . . . . .	11
<b>6</b>	<b>Summary and outlook</b>	<b>12</b>
<b>A</b>	<b>Ein Anhangskapitel</b>	<b>13</b>
	<b>Bibliography</b>	<b>14</b>

# 1 Introduction

The Standard Model (SM) of particle physics is the best descriptor for the behaviour of elementary particles and the interactions between them. Despite its major successes, like the discovery of the Higgs boson (citation) or the top quark(citation), many phenomena remain unexplained. Physics beyond the Standard Model (BSM) is needed to make breakthroughs in dark matter, neutrino oscillations or the matter-antimatter disbalance.

As the heaviest fermion with a mass of about 172.5 GeV, the top quark is the strongest coupler with the Higgs field and may act as a potential probe for electroweak symmetry breaking. Therefore in it lies potential in uncovering BSM physics. The challenge lies in its short lifetime of  $5 \times 10^{-25}$  s, which makes hadronisation impossible and almost exclusively leads to a  $W$ -boson and a bottom-quark as decay products.

At the ATLAS experiment of the Large Hadron Collider (LHC) top-quarks predominantly come in pairs of  $t\bar{t}$  via the strong interaction. Although, top quarks can be produced singly via the electroweak interaction, with the single-top production in the s- and t-channel, as well as in  $tW$  associated production. The s-channel has the smallest cross section of 10.32 pb of all mentioned processes and is dominated heavily by background processes. Unlike the other single modes, the s-channel has not yet been measured precisely with a significance over  $5\sigma$ , with the latest ATLAS measurement achieving an observed (expected) significance of 3.3(3.9). Even with its small cross section, the major hindrance in the measurement is the systematic uncertainties. To get a better measurement, this thesis will use a more modern approach, with a DNN output as the discriminant for a profile likelihood fit. Simulated data from proton-proton collisions in the LHC at  $\sqrt{s} = 13$  TeV corresponding to the full Run 2 dataset with a luminosity of  $140 \text{ fb}^{-1}$  from the ATLAS detector is used.


The structure of this work constitutes the following content. In chapter 2 an overview of the SM, the s-channel production mode as well as the dominant backgrounds and the previous measurement is given. chapter 3 presents the ATLAS detector. chapter 4 lists the samples, the object definitions and event selections used. chapter 5 describes the main study presented in this thesis. Finally chapter 6 gives a short summary of the results and an outlook for further advances in the measurement of the single top s-channel.

## 2 Properties of the Top Quark within the Standard Model

### 2.1 Overview of the Standard Model

The SM of particle physics is a gauge invariant theory, which describes the fundamental particles and the interactions between them. It was developed throughout the 20th century (citation) by combining the discoveries of quantum mechanics and special relativity into a quantum field theory. Three of the four fundamental forces, electromagnetism, the strong interaction and the weak interaction are characterized by it, only excluding gravity. Whereas electromagnetism and gravity act on infinite ranges, the strong and weak force act on subatomic scales.

The SM divides the fundamental particles into two main categories. First fermions, which constitute all known matter and carry half integer spin. Second bosons, which act out the interactions between fermions and carry a whole integer spin. Further, bosons can be divided into vector bosons, with spin 1 and a scalar boson, the Higgs boson ( $H$ ) with spin 0. The fermions are comprised of quarks and leptons, where both interact with the electromagnetic (EM) force but only quarks experience the strong force. Neutrinos, which take part in the lepton category, only interact via the weak force. The fundamental particles are further split into three generations in relation to their masses and time of discovery. The first generation is comprised of the electron ( $e$ ) and the corresponding electron neutrino ( $\nu_e$ ) for the lepton part, and the up quark ( $u$ ) as well as the down quark ( $d$ ). The second generation leptons include the muon ( $\mu$ ) and the muon neutrino ( $\nu$ ), while the charm quark ( $c$ ) and the strange quark ( $s$ ) constitute the second generation quarks. The third generation includes the heaviest fermions, with the tau lepton ( $\tau$ ) and its corresponding tau neutrino ( $\nu_{\text{tau}}$ ) on the lepton side and the top quark ( $t$ ) and bottom quark ( $b$ ) on the quark side. The electric charge defines the interaction strength with the EM force. All charged leptons possess an electric charge of -1 elementary charge ( $e$ ). Whereas up-type quarks ( $u, c, t$ ) carry a charge of  $+2/3e$  and down-type quarks ( $d, s, b$ ) carry a charge of  $-1/3e$ . For every fermion there exists an antiparticle counterpart with opposite electric charge and spin. Especially the first generation of fermions constitute to the overwhelming part of matter, where protons and neutrons are built with the  $u$ - and  $d$ -quarks and electrons as part of atoms.



## 2.2 The Top Quark


First postulated by Makoto Kobayashi and Toshihide Maskawa in 1973 and later discovered at Fermilab with the CDF and DØ experiments of the Tevatron collider in 1995 in  $p\bar{p}$  collisions, the top quark is the last discovered quark in the SM. As up-type quark in the third generation, it carries the heaviest mass of all quarks at  $m_t = (172.53 \pm 0.29) \text{ GeV}$  (citation), it possesses a large decay width of  $\Gamma_t = 1.42^{+0.19}_{-0.15} \text{ GeV}$  (citation). A large decay width leads to a short mean life time of  $\tau = \hbar/\Gamma_t \approx 5 \times 10^{-25} \text{ s}$ . Unlike other quarks, which hadronise due to colour confinement, the top quark decays before hadronisation is possible, due to its short lifetime. The top's properties, like spin information or kinetic properties, are then to be decoded by studying the decay particles, which makes its analysis especially interesting. The top quark decays almost exclusively into a W-boson and a  $b$ -quark.

### 2.2.1 Single top-quark production in the s-channel

The s-channel is one of the electroweak single top-production modes. In the single top-quark s-channel, a top-quark and a bottom antiquark are created from a virtual (off-shell) W-boson, after the annihilation of a quark and antiquark. The s-channel shows a  $tWb$  vertex, where the CKM matrix element  $V_{tb} \approx 1$  is involved. Therefore the s-channel is a strong probe for the electroweak properties of the top quark. In the LHC, the annihilating particles are predominantly valence up quarks ( $u$ ) and antidown ( $\bar{d}$ ) from the pool of sea quarks in the colliding protons. The parton density function (PDF) is a representation of the probability distribution with which a parton carries a fraction of the momentum of a hadron.

- s-channel is our production mode of interest
- $t\bar{t}$  via gluon gluon fusion is predominant mode
- s-channel has the smallest cross section
- Anti quarks from sea quarks in the pdf
- Explain PDFs to explain sea and partons
- How does the theoretical cross section get calculated? Via factorisation theorem
- s-channel feynman diagram
- what does the signature look like?





**Figure 2.2:** Feynman Diagram of the single top-quark s-channel production mode and further decay of the top-quark.

### 2.2.2 Top quark decay

- decay via weak interaction
- decay into W boson and bottom quark because of CKM  $V_{tb}$  approx 1
- W boson can decay hadronically 2/3 or leptonically 1/3. Hadronic channel has bigger phase space because of more color charge possibilities
- dileptonic, semileptonic or hadronic challenge
- hadronic decay can be detected through jets
- leptonic decay leads to undetected neutrino. This leads to  $E_t^{\text{miss}}$  (how is it calculated?)
- What do the different channel signatures look like?
- We use semileptonic channel because of multijet background


### 2.2.3 Background Processes

Hier kann ich im Prinzip alles aus der Präsi rein hauen

- What is background (Processes with similar signature)
- What are the most relevant backgrounds
- Feynman diagrams of background
- erklären wann background als signature gesehen werden kann
- Also im Prinzip präsi. Bei Noah steht nicht viel

### 2.2.4 Previous measurement

In Figure 2.3 previous measurements of the three different single top-quark production modes at different center of mass energies  $\sqrt{s}$  are shown, taken at the ATLAS and CMS experiments at the LHC with data from  $pp$ -collisions.



**Figure 2.3:** Chart of all measurements taken of the three different single top production modes at the ATLAS and CMS experiments [prev\_measure]


The most recent measurement of the s-channel was done by ATLAS with a center of mass energy of  $\sqrt{s} = 13$  TeV, with data collected between the years 2015 and 2018, corresponding to the Run 2 dataset at an integrated luminosity of  $140 \text{ fb}^{-1}$ . The cross section of the measured s-channel resulted in  $\sigma_{\text{s-chan}} = 8.2 \pm 0.6(\text{stat.})_{-2.8}^{+3.4}(\text{syst.})\text{pb}$ , in comparison to the theoretical estimate of  $\sigma_{\text{s-chan}}^{\text{theo.}} = 10.32_{-0.36}^{+0.40} \text{ pb}$ . This is equivalent with an observed (expected) signal significance over the background only hypothesis of  $3.3(3.9)\sigma$ . The analysis strategy consisted of a preselection and then a split into one signal region and three validation regions, for better modelling of the background processes. To extract the signal strength from the events, a profile likelihood fit was done, with a matrix element method (MEM) discriminant. The limiting factors of the previous measurement were noticeably the systematic uncertainties.

### 3 detector

- 4  $\pi$  detector
- central barrel and two endcaps
- Detector measurements
- magnet system
- inner detector
- electromagnetic calorimeter
- hadronic calorimeter
- Muon spectrometer
- LHCs Coordinate System
- pseudorapidity and distance between particles, definition of transverse momentum

#### Extra Punkte

- Infos zum LHC hinzufügen?
- Inner detector: Pixel Detector, Semiconductor detector, Transition Radiation Tracker
- Trigger system?



**Figure 3.1:** Schematic overview of the ATLAS Detector and all its subsystems [1]

## 4 objet definition and event selection

### 4.1 Monte Carlo Samples

Das hier ist im Prinzip was ich verstehen muss und dazu kann ich mehr in Niklas Master These lesen

- analysis by studying simulations of signal and background processes and seperating them
- Processes are simulated with different monte carlo simulators, which each simulate any process better or worse
- what part is simulated with data? Multijet?
- hard interaction with pertubative theory
- strong coupling non-pertubative
- parton showers and hadronisation described by phenomological models
- PYTHIA: Lund String model, HERWIG: heavy cluster model,
- Matrix element calculation by POWHEG BOX, SHERPA or aMC@NLO
- EVTGEN for heavy flavour decays
- Welche Programme wurden für welche Samples von welchen Prozessen benutzt?
- Wo kann ich diese nachlesen? Die stehen halt im Container Namen ;( + mehr Details bei AMI nachlesen
- Wieso wurden diese Programme und Generatoren für die verschiedenen Prozesse benutzt?
- Welche Prozesse habe ich jetzt genutzt?
  - single top s-channel ist unser signal
  - *t $\bar{t}$ background* dominant background
  - single top *t* channel

- single top  $\bar{t}$
- $\bar{t} + W$ ? Was ist das für ein Prozess?
- $t + W$  ?
- $W$  lep nu, mit veto. Wieso genau haben wir diese benutzt?
- $z$  ee ?
- $z$  mu mu?
- $z$  tau tau?
- was gibt es dort für wichtige Parameter?  $p_T^{\text{hard}}$ ?  $h_{\text{damp}}$ ?
- Zuerst Signal simulation und dann background von der Reihenfolge her
- ATLAS Detector Response simulation via GEANT4 (fullsim) or ATLfast(fast-sim)

## 4.2 object definition

- Nach welchen teilchen schauen wir in der Signatur? leptons (e,  $\mu$ ), neutrinos through  $E_T^{\text{miss}}$ , jets and b-tagged jets
- Ansprechen dass Tau nicht gemessen wird wegen short lifetime, aber decay particles can be possible
- Likelihood criteria? isolation criteria?
- $p_T$  und  $|\eta|$  definitionen für tight and loose particles
- transverse and longitudinal impact parameter specification:  $|z_0 \sin \theta| < 0.5 \text{ mm}$  und  $|d_0/\sigma(d_0)| < 5$
- anti- $k_t$  jet algo?

## 4.3 event selection

- state preselection (TopCP)
- table with final selection (FF)
-

## **5 main study and results**

### **5.1 Analysis strategy**

- 

### **5.2 DNN Model variables**

- 

### **5.3 Separation power**

- 

### **5.4 profile likelihood fit**

- 

### **5.5 systematic uncertainties**

-

## 6 Summary and outlook

??



## A Ein Anhangskapitel

Hier könnte ein Anhang stehen, falls Sie z. B. Code, Konstruktionszeichnungen oder Ähnliches mit in die Arbeit bringen wollen. Im Normalfall stehen jedoch alle Ihre Resultate im Hauptteil der Bachelorarbeit und ein Anhang ist überflüssig.

## Bibliography

- [1] Riccardo Maria Bianchi and ATLAS Collaboration. “ATLAS experiment schematic or layout illustration.” General Photo. 2022. URL: <https://cds.cern.ch/record/2837191>.
- [2] Nicola Serra. *Standard Model*. 2025. URL: <https://www.physik.uzh.ch/groups/serra/StandardModel.html>.

# Eidesstattliche Versicherung

## (Affidavit)

Name, Vorname  
(surname, first name)

Matrikelnummer  
(student ID number)

☐ Bachelorarbeit  
(Bachelor's thesis)

☐ Masterarbeit  
(Master's thesis)

Titel  
(Title)

Ich versichere hiermit an Eides statt, dass ich die vorliegende Abschlussarbeit mit dem oben genannten Titel selbstständig und ohne unzulässige fremde Hilfe erbracht habe. Ich habe keine anderen als die angegebenen Quellen und Hilfsmittel benutzt sowie wörtliche und sinngemäße Zitate kenntlich gemacht. Die Arbeit hat in gleicher oder ähnlicher Form noch keiner Prüfungsbehörde vorgelegen.

I declare in lieu of oath that I have completed the present thesis with the above-mentioned title independently and without any unauthorized assistance. I have not used any other sources or aids than the ones listed and have documented quotations and paraphrases as such. The thesis in its current or similar version has not been submitted to an auditing institution before.

Ort, Datum  
(place, date)

Unterschrift  
(signature)

### Belehrung:

Wer vorsätzlich gegen eine die Täuschung über Prüfungsleistungen betreffende Regelung einer Hochschulprüfungsordnung verstößt, handelt ordnungswidrig. Die Ordnungswidrigkeit kann mit einer Geldbuße von bis zu 50.000,00 € geahndet werden. Zuständige Verwaltungsbehörde für die Verfolgung und Ahndung von Ordnungswidrigkeiten ist der Kanzler/die Kanzlerin der Technischen Universität Dortmund. Im Falle eines mehrfachen oder sonstigen schwerwiegenden Täuschungsversuches kann der Prüfling zudem exmatrikuliert werden. (§ 63 Abs. 5 Hochschulgesetz - HG - ).

Die Abgabe einer falschen Versicherung an Eides statt wird mit Freiheitsstrafe bis zu 3 Jahren oder mit Geldstrafe bestraft.

Die Technische Universität Dortmund wird ggf. elektronische Vergleichswerkzeuge (wie z.B. die Software „turnitin“) zur Überprüfung von Ordnungswidrigkeiten in Prüfungsverfahren nutzen.

Die oben stehende Belehrung habe ich zur Kenntnis genommen:

### Official notification:

Any person who intentionally breaches any regulation of university examination regulations relating to deception in examination performance is acting improperly. This offense can be punished with a fine of up to EUR 50,000.00. The competent administrative authority for the pursuit and prosecution of offenses of this type is the Chancellor of TU Dortmund University. In the case of multiple or other serious attempts at deception, the examinee can also be unenrolled, Section 63 (5) North Rhine-Westphalia Higher Education Act (*Hochschulgesetz, HG*).

The submission of a false affidavit will be punished with a prison sentence of up to three years or a fine.

As may be necessary, TU Dortmund University will make use of electronic plagiarism-prevention tools (e.g. the "turnitin" service) in order to monitor violations during the examination procedures.

I have taken note of the above official notification:\*

Ort, Datum  
(place, date)

Unterschrift  
(signature)

**\*Please be aware that solely the German version of the affidavit ("Eidesstattliche Versicherung") for the Bachelor's/ Master's thesis is the official and legally binding version.**