

---

## **Topic 5. Linear Model Selection and Regularization**

# Review: Linear Regression

$$Y = \beta_0 + \beta_1 X_1 + \beta_2 X_2 + \dots + \beta_p X_p + \varepsilon$$

$\beta_0$  ---- intercept (i.e., the average value of  $Y$  if all inputs are zero)

$\beta_j$  ---- slope for the  $j$ th predictor (the average increase in  $Y$  when  $X_j$  is increased by 1 and all other predictors are held constant)


# Review: Estimating Coefficients

- Given a training data set

$$\{(\mathbf{x}_1, y_1), (\mathbf{x}_2, y_2), \dots, (\mathbf{x}_n, y_n)\}$$
$$\mathbf{x}_i = (x_1, x_2, \dots, x_p)$$

- Use Least Squares (LS) method to find coefficient estimates

$$\text{minimize } RSS = \sum_{i=1}^n (y_i - \hat{y}_i)^2$$
$$= \sum_{i=1}^n (y_i - \hat{\beta}_0 - \hat{\beta}_1 x_1 - \dots - \hat{\beta}_p x_p)^2$$



# Improving the Least Squares Estimates

- We want to improve the linear regression model, by replacing the ordinary least square (OLS) fitting with some alternative fitting procedure.
- Two reasons for improving the OLS model
  - Prediction Accuracy
  - Model Interpretability

# 1. Prediction Accuracy

- The least squares estimates have relatively low bias and low variability especially when
  - the relationship between  $Y$  and  $X$  is linear and
  - # observations is way bigger than # predictors ( $n \gg p$ )
- When  $n \approx p$ , OLS fit may have **high variance**, and result in overfitting and poor estimates on unseen observations.
- When  $n < p$ , OLS fit does not work, the variance of these estimates is infinite so this method cannot be used at all.

## 2. Model Interpretability

- When we have a large number of  $X$  variables in the model there will generally be many that have little or no effect on  $Y$ .
- Leaving these variables (terms) in the model makes it harder to see the “big picture”, i.e., the effect of the “important variables”.
- The model would be easier to interpret by removing the unimportant variables (i.e., setting the coefficients of those variables to zero).

# Solution

## ➤ Subset Selection

- Identifying a subset of the  $p$  predictors that we believe to be related to the response, and then fitting the model using this subset.
- E.g., best subset selection and stepwise selection

## ➤ Shrinkage

- Involves shrinking the estimates of coefficients toward zero
- The shrinkage reduces the variance.
- Some of the coefficients may shrink to exactly zero, and hence shrinkage methods can also perform variable selection.
- E.g. Ridge regression and the Lasso

## ➤ Dimension Reduction

- Involves projecting the  $p$  predictors into an  $M$ -dimensional space, where  $M < p$ , and then fitting regression model of  $Y$  on the projections.
- E.g. Principle Components Regression

# Subset Selection



# Subset Selection

- From the  $p$  predictor variables, choose a subset of them such that the linear model with this subset as predictors has best **prediction accuracy**.
- Popular methods
  - Best subset selection
  - Stepwise selection

# 1. Best Subset Selection

- In this approach, we run a linear regression for each possible combination of the predictors.
- The set of possible models include: models that contain exactly one predictor, models that contain exactly two predictors, ..., and the model with all the  $p$  predictors.
- There are  $\binom{p}{k}$ ,  $k = 1, \dots, p$  models that contain exactly  $k$  predictors. For example,
  - There are  $\binom{p}{1} = p$  models that contain exactly **one** predictor.
  - There are  $\binom{p}{2} = p(p - 1)/2$  models that contain exactly **two** predictors.

# Procedure

- Step 1: find the best model among the models with the same number of predictors
  - (Model<sub>0</sub>) the model with no predictors
  - (Model<sub>1</sub>) the best model among the models with 1 predictor
  - (Model<sub>2</sub>) the best model among the models with 2 predictors
  - .....
  - (Model <sub>$p - 1$</sub> ) the best model among the models with  $p - 1$  predictors
  - (Model <sub>$p$</sub> ) the model with all the  $p$  predictors
- Step 2: find the overall best model among the models obtained in Step 1

# The Algorithm of Best Subset Selection

---

**Algorithm 6.1** *Best subset selection*

---

1. Let  $\mathcal{M}_0$  denote the *null model*, which contains no predictors. This model simply predicts the sample mean for each observation.
  2. For  $k = 1, 2, \dots, p$ :
    - (a) Fit all  $\binom{p}{k}$  models that contain exactly  $k$  predictors.
    - (b) Pick the best among these  $\binom{p}{k}$  models, and call it  $\mathcal{M}_k$ . Here *best* is defined as having the smallest RSS, or equivalently largest  $R^2$ .
  3. Select a single best model from among  $\mathcal{M}_0, \dots, \mathcal{M}_p$  using cross-validated prediction error,  $C_p$  (AIC), BIC, or adjusted  $R^2$ .
-

# Model Selection Criteria

- **Algorithm-2(b):** use  $RSS$  or  $R^2$  as criteria
- **Be cautious:** when the number of predictors in the model increases,  $RSS$  decreases monotonically, and  $R^2$  increases monotonically.
- Therefore, if we use them to select *the single best model*, we will always end up with a model involving all the predictors.
- The problem is that a low  $RSS$  or a high  $R^2$  indicates a model with low *training* error, not test error.  
(in other words, more predictors always fit data better in training.)
- **Algorithm-3:** will be discussed later.

# Example: Credit Data

## ➤ Example: Credit data set

### *Response*

**balance**: average credit card debt

### *Predictors*

**age**

**cards**: number of credit cards

**education**: years of education

**income**: in thousands of dollars

**limit**: credit limit

**rating**: credit rating

**gender**: male/female (1 dummy variable)



**student**: student status (1 dummy variable)

**status**: marital status (1 dummy variable)

**Ethnicity**: Caucasian/African American/Asian (2 dummy variables)

# Example: $RSS$ and $R^2$

- The **red line** tracks the best model for a given number of predictors, according to  $RSS$  and  $R^2$ .
- As expected, the  $RSS/R^2$  will always decline/increase as the number of variables increases.
- From the three-predictor model on, adding predictors lead to little improvement.



# Drawback of Best Subset Selection

- Best subset selection is a conceptually simple approach. But it is computationally intensive especially when we have a large number of predictors (large  $p$ ).
- The number of possible models  $= 2^p$ . If  $p = 10$ , that means we will consider 1024 possible models. If  $p = 20$ , there are over one million possibilities!
- This method becomes computationally infeasible for values of  $p$  greater than around 40, even with extremely fast modern computers.



## 2. Stepwise Selection

- Best subset selection searches the enormous space of possible models, whereas stepwise selection explores a far more **restricted** set of models, which are more computationally efficient.
- Two methods:
  - Forward stepwise selection: Begins with the model containing no predictor, and then adds one predictor at a time that improves the model the most until no further improvement is possible
  - Backward stepwise selection: Begins with the model containing all predictors, and then deleting one predictor at a time that improves the model the most until no further improvement is possible

# Forward Stepwise Selection

---

**Algorithm 6.2** *Forward stepwise selection*

---

1. Let  $\mathcal{M}_0$  denote the *null* model, which contains no predictors.
  2. For  $k = 0, \dots, p - 1$ :
    - (a) Consider all  $p - k$  models that augment the predictors in  $\mathcal{M}_k$  with one additional predictor.
    - (b) Choose the *best* among these  $p - k$  models, and call it  $\mathcal{M}_{k+1}$ . Here *best* is defined as having smallest RSS or highest  $R^2$ .
  3. Select a single best model from among  $\mathcal{M}_0, \dots, \mathcal{M}_p$  using cross-validated prediction error,  $C_p$  (AIC), BIC, or adjusted  $R^2$ .
-

# A Simple Example of Forward Stepwise Selection

➤ Assume predictors  $X_1, X_2, X_3$ ,  $p = 3$

➤ Algorithm-1

The null model  $\rightarrow M_0 = Y \sim \text{intercept}$

➤ Algorithm-2

1. Consider adding one predictor ( $X_1$  or  $X_2$  or  $X_3$ )

3 models with one predictor:  $Y \sim X_1$ ,  $Y \sim X_2$ ,  $Y \sim X_3$

2. Choose the best among them  $\rightarrow M_1 = Y \sim X_2$

3. Then consider adding one more predictor ( $X_1$  or  $X_3$ )

2 models with two predictors:  $Y \sim X_2 + X_1$ ,  $Y \sim X_2 + X_3$

4. Choose the best among them  $\rightarrow M_2 = Y \sim X_2 + X_1$

5. Consider adding one more predictor ( $X_3$ )  $\rightarrow M_3 = Y \sim X_2 + X_1 + X_3$

➤ Algorithm-3

Choose the best model among  $M_0, M_1, M_2, M_3$

# Computational Advantage

- Best subset selection involves  $2^p$  models. Forward stepwise selection involves  $1 + p(p + 1)/2$  models.
- When  $p = 20$ , best subset selection requires fitting 1,048,576 models.
- Forward stepwise selection requires fitting only 211 models.

# Example: Credit Data

- However, since forward stepwise selection only searches part of the space of possible models, there is no guarantee that it will find the best possible model as best subset selection does.
- Results on Credit data set: the first four selected models from best subset selection and forward stepwise selection

# Variables	Best subset	Forward stepwise
One	rating	rating
Two	rating, income	rating, income
Three	rating, income, student	rating, income, student
Four	cards, income student, limit	rating, income, student, limit

# Backward Stepwise Selection

---

**Algorithm 6.3** *Backward stepwise selection*

---

1. Let  $\mathcal{M}_p$  denote the *full* model, which contains all  $p$  predictors.
  2. For  $k = p, p - 1, \dots, 1$ :
    - (a) Consider all  $k$  models that contain all but one of the predictors in  $\mathcal{M}_k$ , for a total of  $k - 1$  predictors.
    - (b) Choose the *best* among these  $k$  models, and call it  $\mathcal{M}_{k-1}$ . Here *best* is defined as having smallest RSS or highest  $R^2$ .
  3. Select a single best model from among  $\mathcal{M}_0, \dots, \mathcal{M}_p$  using cross-validated prediction error,  $C_p$  (AIC), BIC, or adjusted  $R^2$ .
-

# A Simple Example of Backward Stepwise Selection

➤ Assume predictors  $X_1, X_2, X_3$ ,  $p = 3$

➤ Algorithm-1

The full model  $\rightarrow M_3 = Y \sim X_1 + X_2 + X_3$

➤ Algorithm-2

1. Consider removing one predictor

3 models with two predictors:  $Y \sim X_2 + X_3$ ,  $Y \sim X_1 + X_3$ ,  $Y \sim X_1 + X_2$

2. Choose the best among them  $\rightarrow M_2 = Y \sim X_2 + X_3$

3. Then consider removing one more predictor ( $X_2$  or  $X_3$ )

2 models with one predictor:  $Y \sim X_2$ ,  $Y \sim X_3$

4. Choose the best among them  $\rightarrow M_1 = Y \sim X_2$

5. Consider removing one more predictor ( $X_2$ )  $\rightarrow M_0 = Y \sim \text{intercept}$

➤ Algorithm-3

Choose the best model among  $M_0, M_1, M_2, M_3$

# Properties about Backward Stepwise Selection

- Like forward stepwise selection, backward selection searches through only  $1 + p(p + 1)/2$  models, and so can be applied in settings where  $p$  is too large to apply best subset selection.
- Also, there is no guarantee to yield the best possible model.
- **Backward vs. Forward:** backward selection requires  $n > p$ , so that the full model can be fit. Forward selection can be used even when  $n < p$ . So forward selection is the only viable subset method when the number of predictors is very large.
- Hybrid approach: variables are added to the model sequentially, in analogy to forward selection. After adding each new variable, remove a variable that no longer provides an improvement in the model fit.




# Selecting the Single Best Model

- Subset selection results in a set of “good” models, each of which contains a subset of the  $p$  predictors. How do we determine which model is the “best” among them in terms of **prediction accuracy**?
- One simple approach is to take the model with the smallest RSS or the largest  $R^2$ .
- As we discussed, unfortunately, the model that includes all the variables will always have the largest  $R^2$  (and smallest RSS). This means that RSS and  $R^2$  are not suitable for selecting the best model among a collection of models with different numbers of predictors.

# Approaches

- **Keep in mind:** we want to find the model that has the best prediction accuracy (or lowest test error).
- Two common approaches
  1. Indirectly estimate the test error by making an adjustment to the training error to account for the bias due to overfitting



2. Directly estimate the test error, using a validation set approach or cross validation

# Approach 1

- These methods add penalty to RSS for the number of predictors (i.e. complexity) in the model (linear regression).

- **AIC** (Akaike information criterion)

$$\text{AIC} = \frac{1}{n\hat{\sigma}^2} (\text{RSS} + 2d\hat{\sigma}^2)$$

- **BIC** (Bayesian information criterion)

$$\text{BIC} = \frac{1}{n} (\text{RSS} + \log(n)d\hat{\sigma}^2)$$

- **$C_p$**  (equivalent to AIC for linear regression)

$$C_p = \frac{1}{n} (\text{RSS} + 2d\hat{\sigma}^2)$$

- **Adjusted  $R^2$**




$$\text{Adjusted } R^2 = 1 - \frac{\text{RSS}/(n - d - 1)}{\text{TSS}/(n - 1)}$$

# Approach 1

- For AIC, BIC and  $C_p$ , a smaller value indicates lower test error, and thus a better model.
- For adjusted  $R^2$ , a larger value indicates a better model.
- AIC vs. BIC: BIC places a heavier penalty on models with many predictors, hence results in the selection of smaller models.
- Adjusted  $R^2$  is not as well motivated in statistical theory as other three.
- All of these measures are simple to use and compute.

# Results of Credit Data: $C_p$ , BIC and Adjusted $R^2$

- $C_p$  (AIC) selects the 6-variable model containing **income**, **limit**, **rating**, **cards**, **age** and **student**.
- BIC selects the 4-variable model containing only **income**, **limit**, **cards** and **student**.
- Adjusted  $R^2$  selects the 7-variable model containing those selected by  $C_p$  (AIC) and **gender**.





# Approach 2

- Compute the validation set error or the cross-validation error for each model, and then select the model that results in smallest test error.
- Advantage over Approach 1: it provides a direct estimate of the test error, and makes fewer assumptions about the true underlying model (Approach 1 mainly applies to linear regression).
- In the past, performing cross validation was computationally prohibitive for problems with large  $p$  and/or  $n$ , so those simple measures, AIC, BIC,  $C_p$  and adjusted  $R^2$ , were often used. Nowadays, with fast computers, cross validation becomes more attractive.

# Results of Credit Data: Cross Validation

- Validation set and cross validation methods both select the 6-predictor model.
- The two methods suggest that the 4-, 5- and 6-predictor models are roughly equivalent in terms of test error.



# **Shrinkage Methods**



# Shrinkage Methods

- Subset selection methods fit a linear model that contains a subset of the predictors through least squares.
- Shrinkage methods fit a model containing all  $p$  predictors using a technique that shrinks the coefficient estimates towards zero.
- Two popular shrinkage methods: Ridge regression and LASSO.

# 1. Ridge Regression

- Ordinary Least Squares (OLS) estimates  $\beta$ s by minimizing

$$\text{RSS} = \sum_{i=1}^n \left( y_i - \beta_0 - \sum_{j=1}^p \beta_j x_{ij} \right)^2$$

- Ridge Regression uses a slightly different equation

$$\sum_{i=1}^n \left( y_i - \beta_0 - \sum_{j=1}^p \beta_j x_{ij} \right)^2 + \lambda \sum_{j=1}^p \beta_j^2 = \text{RSS} + \lambda \sum_{j=1}^p \beta_j^2$$

$\lambda \geq 0$  is a *tuning parameter*

# Ridge Regression Adds a Penalty on $\beta_s$ !

- The effect of this equation is to add a penalty of the form


$$\lambda \sum_{j=1}^p \beta_j^2$$

where the tuning parameter  $\lambda$  is a positive value.

- This has the effect of “*shrinking*” large  $\beta_s$  towards zero.
- Notice that when  $\lambda = 0$ , we get the OLS!
- The shrinkage penalty is applied to coefficients  $\beta_1, \dots, \beta_p$ , not to the intercept  $\beta_0$ . The data matrix  $\mathbf{X}$  should be centered (have mean zero) before ridge regression is performed.


# Credit Data: Ridge Regression

- As  $\lambda$  increases, the coefficients shrink towards zero.




# Credit Data: Ridge Regression

- Another perspective of the change in estimated coefficients



# What Makes the Method Work?

- The OLS estimates to minimize RSS



# Another Formulation for Ridge Regression

- The two formulations are equivalent


$$\underset{\beta}{\text{minimize}} \sum_{i=1}^n \left( y_i - \beta_0 - \sum_{j=1}^p \beta_j x_{ij} \right)^2 + \lambda \sum_{j=1}^p \beta_j^2 = \text{RSS} + \lambda \sum_{j=1}^p \beta_j^2$$



$$\underset{\beta}{\text{minimize}} \left\{ \sum_{i=1}^n \left( y_i - \beta_0 - \sum_{j=1}^p \beta_j x_{ij} \right)^2 \right\} \quad \text{subject to} \quad \sum_{j=1}^p \beta_j^2 \leq s$$

# Ridge Regression Estimates

- Ellipses that are centered around  $\hat{\beta}$  represent regions of constant RSS (i.e., all the points on a given ellipse share a common value of the RSS).
- The solid blue area is the constraint region of ridge regression, which is  $\beta_1^2 + \beta_2^2 \leq s$  in the two-dimension case.
- The ridge regression coefficient estimates are given by **the first point at which an ellipse contacts the constraint region.**





# Why Shrinking towards Zero Is A Good Thing?

## ➤ OLS estimates



- Generally have low bias but can have high variance
- When  $n$  and  $p$  are of similar size or  $n < p$ , the OLS estimates will be extremely variable

## ➤ The penalty term in the ridge regression estimates

- Increases bias
- But can substantially reduce variance

# Simulation Study

- Black: Bias
- Green: Variance
- Purple: Test MSE
- Increase in  $\lambda$  increases bias but decreases variance



# Computational Advantages

- If  $p$  is large, the best subset selection approach requires searching through enormous numbers of possible models.
- With Ridge Regression, for any given  $\lambda$ , we only need to fit one model and the computation turns out to be very simple.
- Ridge Regression can even be used when  $n < p$ , a situation where OLS fails completely!

## 2. LASSO

- Ridge Regression isn't perfect.
- One significant problem: the penalty term will never force any of the coefficients to be **exactly zero**. That means the final model will include all variables, which makes it hard to interpret.
- A more modern alternative is the LASSO.
- The LASSO works in a similar way to Ridge Regression, except that it uses a different penalty term.

# LASSO's Penalty Term

➤ Ridge Regression minimizes

$$\sum_{i=1}^n \left( y_i - \beta_0 - \sum_{j=1}^p \beta_j x_{ij} \right)^2 + \lambda \sum_{j=1}^p \beta_j^2 = \text{RSS} + \lambda \sum_{j=1}^p \beta_j^2.$$

➤ The LASSO minimizes

$$\sum_{i=1}^n \left( y_i - \beta_0 - \sum_{j=1}^p \beta_j x_{ij} \right)^2 + \lambda \sum_{j=1}^p |\beta_j| = \text{RSS} + \lambda \sum_{j=1}^p |\beta_j|.$$

# What's the Big Deal?

- This seems like a very similar idea but there is a big difference.
- Using this penalty, it could be proven mathematically that some coefficients end up being set to **exactly zero**.
- With LASSO, we can produce a model that **has high predictive power and it is simple to interpret**.

# Another Formulation for LASSO



- The two formulations are equivalent

$$\underset{\beta}{\text{minimize}} \sum_{i=1}^n \left( y_i - \beta_0 - \sum_{j=1}^p \beta_j x_{ij} \right)^2 + \lambda \sum_{j=1}^p |\beta_j| = \text{RSS} + \lambda \sum_{j=1}^p |\beta_j|.$$





$$\underset{\beta}{\text{minimize}} \left\{ \sum_{i=1}^n \left( y_i - \beta_0 - \sum_{j=1}^p \beta_j x_{ij} \right)^2 \right\} \quad \text{subject to} \quad \sum_{j=1}^p |\beta_j| \leq s$$

# Reason for the Magic of LASSO







# Credit Data: LASSO





# Ridge Regression vs. LASSO

- Simulation study 1: All 45 predictors are related with response
- Left: Squared bias (black), variance (green), test MSE (purple) of Lasso
- Right: Comparison of lasso (solid) and ridge regression (dotted)



# Ridge Regression vs. LASSO

- Simulation study 1: Response is a function of only 2 predictors
- Left: Squared bias (black), variance (green), test MSE (purple) of Lasso
- Right: Comparison of lasso (solid) and ridge regression (dotted)





# Ridge Regression vs. LASSO

- Neither ridge regression nor the lasso will universally dominate the other in terms of prediction accuracy.
- Ridge regression performs better when the response depends on many predictors, all with coefficients of roughly equal size.
- LASSO performs better when a relatively small number of predictors have substantial coefficients, and the remaining predictors have very small or zero coefficients.

# A Simple Special Case

- Consider a simple case with  $n = p$  and  $\mathbf{X} = \mathbf{I}_p$ , then  $\hat{\beta}_j^{ols} = y_j$ ,
- Ridge regression multiplies  $\hat{\beta}_j^{ridge}$  by a constant,  $\hat{\beta}_j^{ridge} = y_j/(1 + \lambda)$ .
- Lasso truncates  $\hat{\beta}_j^{ridge}$  towards zero by a constant,  $\hat{\beta}_j^{lasso} = \text{sign}(y_j)(|y_j| - \lambda/2)_+$ .



# Selecting the Tuning Parameter



- In subset selection methods, we need to determine which model is the best.
- Similarly, in shrinkage methods, we need to select the optimal value for the tuning parameter  $\lambda$ .
- Different values of the tuning parameter will lead to a model with different levels of prediction accuracy.
- **Question:** Which value of  $\lambda$  will produce the model with lowest test error?

# Approach — Cross Validation

- Select a grid of potential values for  $\lambda$
- For each value of  $\lambda$ , use cross validation to estimate the test error rate
- Select the value that gives the lowest error rate

# Example



- 10-fold cross validation (LOOCV) on Lasso





# Credit Data: Selecting Tuning Parameter

- Leave-one-out cross validation (LOOCV) on Ridge regression



# Other Extensions of LASSO

- Group lasso: if the  $p$  variables are partitioned into  $J$  groups, and then it is desirable to include or exclude the whole group

$$\min_{\beta} \frac{1}{2} \|\mathbf{y} - \mathbf{X}\beta\|^2 + \lambda \sum_{j=1}^p \|\vec{\beta}_j\|_2$$

where  $\vec{\beta}_j$  is a coefficient vector for the  $j$ th group.

- Elastic net:

$$\min_{\beta} \frac{1}{2} \|\mathbf{y} - \mathbf{X}\beta\|^2 + \lambda_1 \|\beta\|_1 + \lambda_2 \|\beta\|_2^2$$

- Fused lasso:

$$\min_{\beta} \frac{1}{2} \|\mathbf{y} - \mathbf{X}\beta\|^2 + \lambda \sum_{j=2}^p \|\beta_j - \beta_{j-1}\|_1$$

# Questions (1)

Read slides of Topic 5. Then answer the following questions.

- Why does the OLS fitting need to be improved?
- What are the methods to improve the OLS fitting?
- What are the similarities and differences of the best subset selection vs. stepwise selection algorithms?
- What are the suitable situations of best subset selection and stepwise selection?
- Can we select the single best model using RSS or  $R^2$ ?
- What are the two approaches to select the single best model?
- Why can AIC, BIC and  $C_p$  be used to select the best model for prediction?
- Why does a smaller value of AIC, BIC and  $C_p$  indicate a better model?

## Questions (2)

- What is the difference between subset selection methods and shrinkage methods?
- Why can shrinking coefficients towards zero improve prediction performance?
- In Ridge regression and LASSO, what is the effect of the tuning parameter  $\lambda$  ?
- What is the advantage of LASSO over Ridge regression?
- When should we use Ridge regression, and when use LASSO?