

This problem is NPC, because we can reduce hamiltonian path on grids to it.

1. dfs. A naive upper bound is $O^*(3^k)$, where k is the number of nonempty cells. We should be able to get more refined bounds, and some related papers are asymptotic number of hamiltonian paths on planar graphs [1], and on grid graphs [2]. An upper bound is the number of self-avoiding walks on the square graph \mathbb{Z}^2 , which is estimated to be $O^*(2.639^k)$ [3, 4].

other related papers: [5].

2. We can use bitmask DP, or planar separator theorem + meet in the middle.

References

- [1] Sudip Biswas, Stephane Durocher, Debajyoti Mondal, and Rahnuma Islam Nishat. Hamiltonian paths and cycles in planar graphs. In *International Conference on Combinatorial Optimization and Applications*, pages 83–94. Springer, 2012.
- [2] Olga Bodroza-Pantic, Bojana Pantic, Ilija Pantic, and Marija Bodroza-Solarov. Enumeration of hamiltonian cycles in some grid graphs. *MATCH Commun. Math. Comput. Chem*, 70(1):181–204, 2013.
- [3] Geoffrey R Grimmett and Zhongyang Li. Counting self-avoiding walks. *arXiv preprint arXiv:1304.7216*, 2013.
- [4] Geoffrey R Grimmett and Zhongyang Li. Self-avoiding walks and connective constants. In *Sojourns in Probability Theory and Statistical Physics-III*, pages 215–241. Springer, 2019.
- [5] Anthony J Guttmann. Self-avoiding walks and polygons—an overview. *arXiv preprint arXiv:1212.3448*, 2012.