

This problem is NPC, because we can reduce hamiltonian path on grids to it.

1. dfs. A naive upper bound is  $O^*(3^k)$ , where  $k$  is the number of nonempty cells. We should be able to get more refined bounds, and some related papers are asymptotic number of hamiltonian paths on planar graphs [1], and on grid graphs [2]. An upper bound is the number of self-avoiding walks on the square graph  $\mathbb{Z}^2$ , which is estimated to be  $O^*(2.639^k)$  [4, 5].

other related papers: [6].

my article: <https://leetcode-cn.com/problems/path-with-maximum-gold/solution/fu-za-du-fen-xi-wei-shi-yao-dfsbu-hui-ch-0v9f/>

2. We can use bitmask DP, or planar separator theorem + meet in the middle.

3. This the (node weighted) longest path problem in planar graphs.  $2^{O(\sqrt{k} \log k)}$  [3].

## References

- [1] Sudip Biswas, Stephane Durocher, Debajyoti Mondal, and Rahnuma Islam Nishat. Hamiltonian paths and cycles in planar graphs. In *International Conference on Combinatorial Optimization and Applications*, pages 83–94. Springer, 2012.
- [2] Olga Bodroza-Pantic, Bojana Pantic, Ilija Pantic, and Marija Bodroza-Solarov. Enumeration of hamiltonian cycles in some grid graphs. *MATCH Commun. Math. Comput. Chem*, 70(1):181–204, 2013.
- [3] Marek Cygan, Fedor V Fomin, Łukasz Kowalik, Daniel Lokshtanov, Dániel Marx, Marcin Pilipczuk, Michał Pilipczuk, and Saket Saurabh. *Parameterized algorithms*, volume 5. Springer, 2015.
- [4] Geoffrey R Grimmett and Zhongyang Li. Counting self-avoiding walks. *arXiv preprint arXiv:1304.7216*, 2013.
- [5] Geoffrey R Grimmett and Zhongyang Li. Self-avoiding walks and connective constants. In *Sojourns in Probability Theory and Statistical Physics-III*, pages 215–241. Springer, 2019.
- [6] Anthony J Guttmann. Self-avoiding walks and polygons—an overview. *arXiv preprint arXiv:1212.3448*, 2012.