

Lecture 6. Elementary Matrices

Elementary Matrices

Elementary Matrices are the matrix representations of three *elementary row operations* as discussed in Lecture 5.

Practice 6.1 Compute E_1A , E_2A , and E_3A , and describe how these products can be obtained by elementary row operations:

$$E_1 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 5 \end{bmatrix}, E_2 = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}, E_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -4 & 0 & 1 \end{bmatrix}, A = \begin{bmatrix} 2 & -1 \\ 1 & 4 \\ 0 & -2 \end{bmatrix}.$$

Elementary Matrices

Elementary Matrices are the matrix representations of three *elementary row operations* as discussed in Lecture 5.

Practice 6.1 Compute E_1A , E_2A , and E_3A , and describe how these products can be obtained by elementary row operations:

$$E_1 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 5 \end{bmatrix}, E_2 = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}, E_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -4 & 0 & 1 \end{bmatrix}, A = \begin{bmatrix} 2 & -1 \\ 1 & 4 \\ 0 & -2 \end{bmatrix}.$$

Practice 6.1 illustrates the following general fact about elementary matrices.

- If an elementary row operation is performed on an $m \times n$ matrix A , the resulting matrix can be written as EA , where E is the $m \times m$ matrix performing the same row operation on A .

Inverse of Elementary Matrices

Theorem 6.2. Each elementary matrix E is invertible. The inverse of E is the elementary matrix of the same type that transforms E back into I .

Inverse of Elementary Matrices

Theorem 6.2. Each elementary matrix E is invertible. The inverse of E is the elementary matrix of the same type that transforms E back into I .

In fact, the inverse of an elementary matrix is constructed by doing the *reverse row operation*.

- If E is obtained by switching rows i and j , then E^{-1} is also obtained by switching rows i and j .

Inverse of Elementary Matrices

Theorem 6.2. Each elementary matrix E is invertible. The inverse of E is the elementary matrix of the same type that transforms E back into I .

In fact, the inverse of an elementary matrix is constructed by doing the *reverse row operation*.

- If E is obtained by switching rows i and j , then E^{-1} is also obtained by switching rows i and j .
- If E is obtained by multiplying row i by the scalar k , then E^{-1} is obtained by multiplying row i by the scalar $\frac{1}{k}$.

Inverse of Elementary Matrices

Theorem 6.2. Each elementary matrix E is invertible. The inverse of E is the elementary matrix of the same type that transforms E back into I .

In fact, the inverse of an elementary matrix is constructed by doing the *reverse row operation*.

- If E is obtained by switching rows i and j , then E^{-1} is also obtained by switching rows i and j .
- If E is obtained by multiplying row i by the scalar k , then E^{-1} is obtained by multiplying row i by the scalar $\frac{1}{k}$.
- E is obtained by adding k times row i to row j , then E^{-1} is obtained by subtracting k times row i from row j , (that is, by adding $-k$ times row i to row j .)

Example 6.3. Find the inverse of $E_1 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -4 & 0 & 1 \end{bmatrix}$.

Example 6.3. Find the inverse of $E_1 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -4 & 0 & 1 \end{bmatrix}$.

- **Solution.** To transform E_1 into I , do $4R_1 + R_3 \rightarrow R_3$. The elementary matrix that does this is

$$E_1^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ +4 & 0 & 1 \end{bmatrix}$$

Practice 6.4. Find the inverse of each matrix.

$$E_1 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 2 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, \quad E_2 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad E_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -5 \end{bmatrix}.$$

Matrix Representation of The Inverse Procedure

- Recall that the procedure of finding A^{-1}

$$[A | I] \Leftrightarrow [I | A^{-1}]$$

Matrix Representation of The Inverse Procedure

- Recall that the procedure of finding A^{-1}

$$[A | I] \Leftrightarrow [I | A^{-1}]$$

- So, if A is a $n \times n$ nonsingular matrix, then A is row equivalent to I_n . Then, since each step of the row reduction of A corresponds to left-multiplication by an elementary matrix,

Matrix Representation of The Inverse Procedure

- Recall that the procedure of finding A^{-1}

$$[A \mid I] \Leftrightarrow [I \mid A^{-1}]$$

- So, if A is a $n \times n$ nonsingular matrix, then A is row equivalent to I_n . Then, since each step of the row reduction of A corresponds to left-multiplication by an elementary matrix, there exist elementary matrices E_1, E_2, \dots, E_p such that

$$\begin{aligned}[A \mid I_n] &\Leftrightarrow [E_1 A \mid E_1 I_n] \\&\Leftrightarrow [E_2(E_1 A) \mid E_2(E_1 I_n)] \\&\quad \vdots \\&\Leftrightarrow [E_p(E_{p-1}\dots E_1 A) \mid E_p(E_{p-1}\dots E_1)I_n] \\&\Leftrightarrow [\quad \quad I_n \quad \quad | \quad E_p(E_{p-1}\dots E_1 I_n)]\end{aligned}$$

- Therefore,

Matrix Representation of The Inverse Procedure

- Recall that the procedure of finding A^{-1}

$$[A \mid I] \Leftrightarrow [I \mid A^{-1}]$$

- So, if A is a $n \times n$ nonsingular matrix, then A is row equivalent to I_n . Then, since each step of the row reduction of A corresponds to left-multiplication by an elementary matrix, there exist elementary matrices E_1, E_2, \dots, E_p such that

$$\begin{aligned}[A \mid I_n] &\Leftrightarrow [E_1 A \mid E_1 I_n] \\&\Leftrightarrow [E_2(E_1 A) \mid E_2(E_1 I_n)] \\&\quad \vdots \\&\Leftrightarrow [E_p(E_{p-1}\dots E_1 A) \mid E_p(E_{p-1}\dots E_1)I_n] \\&\Leftrightarrow [\quad \quad I_n \quad \quad | \quad E_p(E_{p-1}\dots E_1 I_n)]\end{aligned}$$

- Therefore,

$$E_p E_{p-1} \dots E_1 A = I_n \text{ and } A^{-1} = E_p E_{p-1} \dots E_1$$

$$E_p E_{p-1} \dots E_1 A = I_n \text{ and } A^{-1} = E_p E_{p-1} \dots E_1 \quad (1)$$

- Since the product $E_p E_{p-1} \dots E_1$ of invertible matrices is invertible, the equation (1) leads to

$$(E_p E_{p-1} \dots E_1)^{-1} (E_p E_{p-1} \dots E_1) A = (E_p E_{p-1} \dots E_1)^{-1} I_n \Leftrightarrow A = (E_p E_{p-1} \dots E_1)^{-1}$$

$$E_p E_{p-1} \dots E_1 A = I_n \text{ and } A^{-1} = E_p E_{p-1} \dots E_1 \quad (1)$$

- Since the product $E_p E_{p-1} \dots E_1$ of invertible matrices is invertible, the equation (1) leads to

$$(E_p E_{p-1} \dots E_1)^{-1} (E_p E_{p-1} \dots E_1) A = (E_p E_{p-1} \dots E_1)^{-1} I_n \Leftrightarrow A = (E_p E_{p-1} \dots E_1)^{-1}$$

Example 6.5. Find the elementary matrices such that $A^{-1} = E_p E_{p-1} \dots E_1$.

$$A = \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix}$$

Practice 6.6. Find the elementary matrices such that $A^{-1} = E_p E_{p-1} \dots E_1$.

$$A = \begin{bmatrix} 0 & 1 & 2 \\ 1 & 0 & 3 \\ 4 & -3 & 8 \end{bmatrix}$$