



B Genomic context of TE intervals and peak midpoint proxies
Canonical TE reference
all TE interval midpoints · n=3,651,522
Native peaks
all peak midpoints · n=68,126
Native peaks
TE-positive peak midpoints · n=5,734
Published peaks (mapped mm10)
all peak midpoints · n=339,735
Published peaks (mapped mm10)
TE-positive peak midpoints · n=54,191
0%
20%
40%
60%
80%
100%
Percentage of intervals; one mutually exclusive context per midpoint
Promoter <=1 kb
Promoter 1-2 kb
Promoter 2-3 kb
5' UTR
3' UTR
First exon
Other exon
First intron
Other intron
Downstream <=300 bp
Distal intergenic




Full BED0 half−open regions; each peak ID counted once. External coordinate benchmark.

Official archived release; identity with paper authors local files is unproved.


C Gene-body and TE membership of peak midpoint proxies






Each peak ID counted once. Circle areas are schematic, not proportional to counts.

D Canonical TE families at peak midpoint proxies




Numbers beside bars are unique peak IDs. Families may overlap; percentages need not sum to 100%.


E TE-family association with peak midpoint proxies




6 zero-hit families omitted from log axes; all 43 retained in tables.



6 zero-hit families omitted from log axes; all 43 retained in tables.



2 zero-hit families omitted from log axes; all 43 retained in tables.



2 zero-hit families omitted from log axes; all 43 retained in tables.

DNA LINE LTR RC Retroposon SINE

Dashed threshold: FDR 0.05; dotted: exploratory FDR 0.10. Interval-defined RepeatMasker loci; one-sided hypergeometric test.
Chromosome/length/GC sensitivity is descriptive; mappability is unmatched. Coordinate association does not establish TE activation.