



9,511 published ATAC Microglia cells / 20 animals — job 8769240






Circles: female; triangles: male. Four mice per condition. Descriptive region annotation; no condition test.
Categories may overlap. These are not TE fragment counts or RNA TE expression.



TE class representation in each mouse: all 20 animals, equal-mouse means
DNA
LINE
SINE
LTR
RC
Retroposon
Accessible peak overlap (%)
WT3
WT6
WT21
FAD3
APOE4
TREM2
Female circles / male triangles; four mice per condition. Different per-mouse detected peak sets; descriptive, no genotype/age test.




Exact paired label-swap tests; BH includes all tested TE levels. Local genomic correlation may inflate significance.
No animal genotype/age effect or autonomous TE activity is established.



Exact paired label-swap tests; BH includes all tested TE levels. Local genomic correlation may inflate significance.
No animal genotype/age effect or autonomous TE activity is established.



Exact paired label-swap tests; BH includes all tested TE levels. Local genomic correlation may inflate significance.
No animal genotype/age effect or autonomous TE activity is established.



Actual accepted job: 8769240; 9,511 published ATAC Microglia cells; all 20 animals.
Primary callable Microglia-accessible regions: 62,956; complete-covariate query regions: 62,950.
Matched query/control pairs: 53,607; query retained: 85.16% of complete-covariate query.
Test status: matching_failed_no_P_values

Primary peaks: positive in >=max(3, ceil(1% of mouse Microglia cells)) cells per mouse,
with at least two qualifying mice. Any blacklist-overlapping peak is excluded entirely.
Matching: chromosome/gene context + GC, length, mappability, TSS distance and
non-Microglia detection prevalence. The result is conditional on this matched subset.

Published MAPQ >30 / primary / properly paired processing can underrepresent repetitive TEs.
GenMap K76E2 is a single-read proxy, not correction of paired-fragment multimapping selection.
Local genomic correlation may inflate exploratory paired-test significance.
These are accessible-region annotations, not TE-fragment counts or RNA TE expression.
No animal condition differential test or autonomous TE activity is established.
The 22,251 selected RNA cells are a different assay/selection from these 9,511 ATAC cells.