Target-only ATAC peak-bound TE loci: normalization and aggregation sensitivity
Blue: WT_3mo; red: target; four mice per group; circles: female, triangles: male

WT_6mo
9,361 loci

WT_21mo
9,725 loci

5xFAD_3mo
7,240 loci

APOE4_TREM2_3mo
6,438 loci

Total-TE normalization
Z-score of the set total

TMM normalization
Z-score of the set total

Gene-RNA-depth normalization
Z-score of the set total

Previous locus-wise scaling
Mean of individual locus z-scores

WT_3mo

WT_6mo

WT_21mo

5xFAD_3mo

APOE4_TREM2_3mo

WT_3mo-only ATAC peak-bound TE loci: normalization and aggregation sensitivity
Blue: WT_3mo; red: target; four mice per group; circles: female, triangles: male

WT_6mo
8,188 loci

WT_21mo
8,767 loci

5xFAD_3mo
10,136 loci

APOE4_TREM2_3mo
10,615 loci


Total-TE normalization
Z-score of the set total

TMM normalization
Z-score of the set total

Gene-RNA-depth normalization
Z-score of the set total

Previous locus-wise scaling
Mean of individual locus z-scores

WT_3mo
WT_6mo
WT_21mo
5xFAD_3mo
APOE4_TREM2_3mo


5xFAD target-only loci: abundance and locus-average summaries answer different questions
Descriptive per-locus mean TE-CPM direction
Higher in 5xFAD
Lower in 5xFAD
TE loci
0
200
400
600
800
1000
0
1–4
5–19
20+
Assigned RNA counts per locus across 8 mice
Contributions to the original set total
Change in summed TE-CPM, 5xFAD − WT_3mo
4000
3000
2000
1000
0
−1000
−2000
−3000
Positive locus contributions
Negative locus contributions
Net change