
Microglia RNA TE overview


22,251 provisional RNA Microglia-like cells | 20 mice | 5 conditions | fractional expected TE allocations

A Actual RNA cohort: 20 mice, five conditions




Labels are unique within each condition; full animal IDs are retained in cohort_compact_labelmap.tsv.

B Saved gene RNA UMAP: original provisional states




Exact selected-cell join; coordinates retain the original 52,016-cell gene fit.

C Gene RNA layout




TE composition layout




Separate embeddings: distances and positions are not aligned. No joint RNA–ATAC or label-transfer inference.


D Actual cell QC distributions; descriptive cells, biological replicates are mice




E Provisional TE composition groups




Original gene states on TE layout



Condition on TE layout




TE allocation on TE layout




TE-only composition embedding; depth and biological covariates are diagnostic overlays, not validated states.

E: TE groups are provisional and depth-associated.
TE composition PC1 correlates with log1p TE allocation (Pearson r = 0.920949); this is not validated biological state separation.

F Mouse-level TE family composition



G Equal-mouse TE family composition



Dot area = fraction of four mice with signal. Denominator: each mouse's total exported TE allocation; no DA test.

Separate gene and TE RNA layouts: no joint RNA-ATAC embedding or coordinate alignment. Gene coordinates retain the original 52,016-cell fit; exact selected IDs are shown. Family composition divides by each mouse's total exported TE allocation; mice are biological replicates. Expected allocations are not observed molecule counts or whole-transcriptome expression. Individual panel PDFs retain readable full-size axes and tables remain the source of exact values.