STAT 1210 - Quiz 4

Part IV: Inference for Population Means - Version 1

Fall 2026

Name:

_____

Student number:

_____

Instructions

- You have 40 minutes to complete the quiz.
- All solutions must be handwritten on this quiz and the supplied answer pages.
- Show the calculations needed to support each numerical answer.
- Explain conclusions in the context of the question.
- Report numerical answers to at least three decimal places when appropriate.
- A t-distribution table excerpt is provided on the final page.
- R coding is not required for this quiz.

Empty Page

Question 1: Shell Length of Cultured PEI Mussels (8 marks)

A mussel grower in Prince Edward Island historically harvested blue mussels with a population mean shell length of 60 mm. This season, the grower selects a random sample of 25 harvest-ready mussels from a lease containing several thousand mussels. The sample has mean shell length $\bar{x} = 57.6$ mm and sample standard deviation $s = 6.0$ mm. A dotplot of the sample is approximately symmetric and shows no strong outliers.

The grower wants to determine whether the population mean shell length this season differs from the historical value. Use $\alpha = 0.05$.

1. Define the population parameter μ. State the null and alternative hypotheses.
2. Calculate the standard error, the test statistic and the degrees of freedom. You may use

$$t = \frac{\bar{x} - \mu_0}{s/\sqrt{n}}$$

3. Use the t-distribution table to give a range for the two-sided P-value.
4. Construct a 95% confidence interval using $t^* = 2.064$.

Answer Page for Question 1

Question 2: Potato Yield under Two Disease-Control Programs (12 marks)

An agricultural research station in Prince Edward Island compares two disease-control programs for a potato crop. The researchers randomly assign 34 similar field plots to either an enhanced program or the current standard program. At harvest, they measure marketable potato yield in tonnes per hectare. The two groups are independent, and the yield distributions are approximately symmetric with no strong outliers.

Disease-control program	Number of plots n	Mean yield $\bar{x}$	Standard deviation s
Enhanced	18	35.8 tonnes/ha	4.2 tonnes/ha
Standard	16	32.4 tonnes/ha	3.6 tonnes/ha

The research question specified before the experiment is whether the enhanced program increases the population mean marketable yield. Define the difference as enhanced minus standard and use $\alpha = 0.05$.

1. Define μ_E and μ_S. State the null and alternative hypotheses.
2. Calculate the point estimate $\bar{x}_E - \bar{x}_S$ and its standard error. You may use

$$\text{SE} = \sqrt{\frac{s_E^2}{n_E} + \frac{s_S^2}{n_S}}$$

3. Calculate the Welch degrees of freedom and the test statistic. Use the t-distribution table to give a range for the one-sided P-value. State the test decision and conclusion in context. You may use

$$df = \frac{\left(\frac{s_E^2}{n_E} + \frac{s_S^2}{n_S}\right)^2}{\frac{1}{n_E - 1}\left(\frac{s_E^2}{n_E}\right)^2 + \frac{1}{n_S - 1}\left(\frac{s_S^2}{n_S}\right)^2}$$

$$t = \frac{(\bar{x}_E - \bar{x}_S) - 0}{\sqrt{s_E^2/n_E + s_S^2/n_S}}$$

4. Construct a two-sided 95% confidence interval for $\mu_E - \mu_S$ using $t^* = 2.037$.
5. Explain what random assignment contributes to the study. Can the results automatically be generalized to every potato field in Prince Edward Island?

Answer Page for Question 2

Answer Page for Question 2

Answer Page for Question 2

Student's t Critical-Value Table

Each entry is a positive critical value t^* with the indicated probability to its right. For a two-sided P-value, double the corresponding one-sided tail probability. When the calculated degrees of freedom are not listed, use the next lower available value.

	Probability to the right of t^*										
df	.250	.200	.150	.100	.050	.025	.010	.005	.0025	.001	.0005
1	1.000	1.376	1.963	3.078	6.314	12.706	31.821	63.657	127.321	318.309	636.619
2	0.816	1.061	1.386	1.886	2.920	4.303	6.965	9.925	14.089	22.327	31.599
3	0.765	0.978	1.250	1.638	2.353	3.182	4.541	5.841	7.453	10.215	12.924
4	0.741	0.941	1.190	1.533	2.132	2.776	3.747	4.604	5.598	7.173	8.610
5	0.727	0.920	1.156	1.476	2.015	2.571	3.365	4.032	4.773	5.893	6.869
6	0.718	0.906	1.134	1.440	1.943	2.447	3.143	3.707	4.317	5.208	5.959
7	0.711	0.896	1.119	1.415	1.895	2.365	2.998	3.499	4.029	4.785	5.408
8	0.706	0.889	1.108	1.397	1.860	2.306	2.896	3.355	3.833	4.501	5.041
9	0.703	0.883	1.100	1.383	1.833	2.262	2.821	3.250	3.690	4.297	4.781
10	0.700	0.879	1.093	1.372	1.812	2.228	2.764	3.169	3.581	4.144	4.587
11	0.697	0.876	1.088	1.363	1.796	2.201	2.718	3.106	3.497	4.025	4.437
12	0.695	0.873	1.083	1.356	1.782	2.179	2.681	3.055	3.428	3.930	4.318
13	0.694	0.870	1.079	1.350	1.771	2.160	2.650	3.012	3.372	3.852	4.221
14	0.692	0.868	1.076	1.345	1.761	2.145	2.624	2.977	3.326	3.787	4.140
15	0.691	0.866	1.074	1.341	1.753	2.131	2.602	2.947	3.286	3.733	4.073
16	0.690	0.865	1.071	1.337	1.746	2.120	2.583	2.921	3.252	3.686	4.015
17	0.689	0.863	1.069	1.333	1.740	2.110	2.567	2.898	3.222	3.646	3.965
18	0.688	0.862	1.067	1.330	1.734	2.101	2.552	2.878	3.197	3.610	3.922
19	0.688	0.861	1.066	1.328	1.729	2.093	2.539	2.861	3.174	3.579	3.883
20	0.687	0.860	1.064	1.325	1.725	2.086	2.528	2.845	3.153	3.552	3.850
21	0.686	0.859	1.063	1.323	1.721	2.080	2.518	2.831	3.135	3.527	3.819
22	0.686	0.858	1.061	1.321	1.717	2.074	2.508	2.819	3.119	3.505	3.792
23	0.685	0.858	1.060	1.319	1.714	2.069	2.500	2.807	3.104	3.485	3.768
24	0.685	0.857	1.059	1.318	1.711	2.064	2.492	2.797	3.091	3.467	3.745
25	0.684	0.856	1.058	1.316	1.708	2.060	2.485	2.787	3.078	3.450	3.725
26	0.684	0.856	1.058	1.315	1.706	2.056	2.479	2.779	3.067	3.435	3.707
27	0.684	0.855	1.057	1.314	1.703	2.052	2.473	2.771	3.057	3.421	3.690
28	0.683	0.855	1.056	1.313	1.701	2.048	2.467	2.763	3.047	3.408	3.674
29	0.683	0.854	1.055	1.311	1.699	2.045	2.462	2.756	3.038	3.396	3.659
30	0.683	0.854	1.055	1.310	1.697	2.042	2.457	2.750	3.030	3.385	3.646
40	0.681	0.851	1.050	1.303	1.684	2.021	2.423	2.704	2.971	3.307	3.551
60	0.679	0.848	1.045	1.296	1.671	2.000	2.390	2.660	2.915	3.232	3.460
80	0.678	0.846	1.043	1.292	1.664	1.990	2.374	2.639	2.887	3.195	3.416
100	0.677	0.845	1.042	1.290	1.660	1.984	2.364	2.626	2.871	3.174	3.390
1000	0.675	0.842	1.037	1.282	1.646	1.962	2.330	2.581	2.813	3.098	3.300
∞	0.674	0.842	1.036	1.282	1.645	1.960	2.326	2.576	2.807	3.090	3.291

For negative test statistics, use symmetry and compare the absolute value $|t|$ with the positive entries in the table.