

A. Cottage Village

time limit per test: 2 seconds

memory limit per test: 64 megabytes

input: standard input

output: standard output

A new cottage village called «Flatville» is being built in Flatland. By now they have already built in «Flatville» n square houses with the centres on the Ox -axis. The houses' sides are parallel to the coordinate axes. It's known that no two houses overlap, but they can touch each other.

The architect bureau, where Peter works, was commissioned to build a new house in «Flatville». The customer wants his future house to be on the Ox -axis, to be square in shape, have a side t , and touch at least one of the already built houses. For sure, its sides should be parallel to the coordinate axes, its centre should be on the Ox -axis and it shouldn't overlap any of the houses in the village.

Peter was given a list of all the houses in «Flatville». Would you help him find the amount of possible positions of the new house?

Input

The first line of the input data contains numbers n and t ($1 \leq n, t \leq 1000$). Then there follow n lines, each of them contains two space-separated integer numbers: $x_i a_i$, where x_i — x -coordinate of the centre of the i -th house, and a_i — length of its side ($-1000 \leq x_i \leq 1000, 1 \leq a_i \leq 1000$).

Output

Output the amount of possible positions of the new house.

Examples

input
2 2 0 4 6 2
output
4

input
2 2 0 4 5 2
output
3

input
2 3 0 4 5 2
output
2

Note

It is possible for the x -coordinate of the new house to have non-integer value.