Verification outcome · Two-arm binary · Odds Ratio · Comparison: Intensive individualized cognitive behavioural therapy plus structured clinical follow-up / Standard clinical care · Follow-up: first

| Study or subgroup | Intensive individualized cognitive behavioural therapy plus structured clinical follow-up<br>Events / Total | Standard clinical care<br>Events / Total | Odds ratio, IV, random (95% CI) | Weight (%) | Odds ratio, IV, random (95% CI) |
|---|---|---|---|---|---|
| Fixture 1 | 4 / 20 | 6 / 22 | | 12.4 | 0.666667 (0.157558 to 2.820825) |
| Fixture 2 | 5 / 21 | 7 / 23 | | 14.3 | 0.714286 (0.186917 to 2.729572) |
| Fixture 3 | 6 / 22 | 8 / 24 | | 16.1 | 0.750000 (0.211647 to 2.657722) |
| Fixture 4 | 7 / 23 | 9 / 25 | | 17.7 | 0.777778 (0.232730 to 2.599313) |
| Fixture 5 | 8 / 24 | 10 / 26 | | 19.1 | 0.800000 (0.250907 to 2.550748) |
| Fixture 6 | 9 / 25 | 11 / 27 | | 20.5 | 0.818182 (0.266738 to 2.509662) |
| **Random-effects model (DerSimonian–Laird)** | **39 / 135** | **51 / 147** | | **100.0** | **0.761368 (0.458516 to 1.264253)** |




Heterogeneity: $\tau^2 = 0.000000$; $Q = 0.065861$, $df = 5$ ($P = 0.999942$); $I^2 = 0.0\%$

Test for overall effect: $Z = -1.053719$ ($P = 0.292012$)

95% prediction interval: [0.458516, 1.264253]