Verification outcome · Two-arm binary · Odds Ratio · Comparison: Intensive individualized cognitive behavioural therapy plus structured clinical follow-up / Standard clinical care · Follow-up: first

| Study | Intensive individualized cognitive behavioural therapy plus structured clinical follow-up | | Standard clinical care | | Weight | Odds ratio IV, Random, 95% CI |
|---|---|---|---|---|---|---|
| | Events | Total | Events | Total | | |
| Fixture 1 | 4 | 20 | 6 | 22 | 12.4% | 0.666667 [0.157558, 2.820825] |
| Fixture 2 | 5 | 21 | 7 | 23 | 14.3% | 0.714286 [0.186917, 2.729572] |
| Fixture 3 | 6 | 22 | 8 | 24 | 16.1% | 0.750000 [0.211647, 2.657722] |
| Fixture 4 | 7 | 23 | 9 | 25 | 17.7% | 0.777778 [0.232730, 2.599313] |
| Fixture 5 | 8 | 24 | 10 | 26 | 19.1% | 0.800000 [0.250907, 2.550748] |
| Fixture 6 | 9 | 25 | 11 | 27 | 20.5% | 0.818182 [0.266738, 2.509662] |
| **Random-effects model (DerSimonian–Laird)** | | **135** | | **147** | **100.0%** | **0.761368 [0.458516, 1.264253]** |
| Total events: | 39 | | 51 | | | |

Odds ratio IV, Random, 95% CI




Heterogeneity: $\tau^2 = 0.000000$; $Q = 0.065861$, $df = 5$ ($P = 0.999942$); $I^2 = 0.0\%$

Test for overall effect: $Z = -1.053719$ ($P = 0.292012$)

95% prediction interval: [0.458516, 1.264253]