

Definition 7.1

- ▶ A k -coloring of a graph G is a labeling $f : V(G) \rightarrow \{1, 2, \dots, k\}$.

Definition 7.1

- ▶ A k -coloring of a graph G is a labeling $f : V(G) \rightarrow \{1, 2, \dots, k\}$.
- ▶ The labels are **colors**.

Definition 7.1

- ▶ A k -coloring of a graph G is a labeling $f : V(G) \rightarrow \{1, 2, \dots, k\}$.
- ▶ The labels are **colors**.
- ▶ The vertices with color i are a **color class**.

Definition 7.1

- ▶ A k -coloring of a graph G is a labeling $f : V(G) \rightarrow \{1, 2, \dots, k\}$.
- ▶ The labels are **colors**.
- ▶ The vertices with color i are a **color class**.
- ▶ A k -coloring f is **proper** if $f(x) \neq f(y)$ whenever x and y are adjacent.

Chromatic Number

Definition 7.1

- ▶ A k -coloring of a graph G is a labeling $f : V(G) \rightarrow \{1, 2, \dots, k\}$.
- ▶ The labels are **colors**.
- ▶ The vertices with color i are a **color class**.
- ▶ A k -coloring f is **proper** if $f(x) \neq f(y)$ whenever x and y are adjacent.
- ▶ The **chromatic number** $\chi(G)$ is the minimum k such that G is k -colorable.

Chromatic Number

Definition 7.1

- ▶ A k -coloring of a graph G is a labeling $f : V(G) \rightarrow \{1, 2, \dots, k\}$.
- ▶ The labels are **colors**.
- ▶ The vertices with color i are a **color class**.
- ▶ A k -coloring f is **proper** if $f(x) \neq f(y)$ whenever x and y are adjacent.
- ▶ The **chromatic number** $\chi(G)$ is the minimum k such that G is k -colorable.
- ▶ If $\chi(G) = k$, then G is **k -chromatic**.

Definition 7.1

- ▶ A k -coloring of a graph G is a labeling $f : V(G) \rightarrow \{1, 2, \dots, k\}$.
- ▶ The labels are **colors**.
- ▶ The vertices with color i are a **color class**.
- ▶ A k -coloring f is **proper** if $f(x) \neq f(y)$ whenever x and y are adjacent.
- ▶ The **chromatic number** $\chi(G)$ is the minimum k such that G is k -colorable.
- ▶ If $\chi(G) = k$, then G is **k -chromatic**.
- ▶ If $\chi(G) = k$, but $\chi(H) < k$ for every proper subgraph H of G , then G is **k -color-critical** or **k -critical**.

Definition 7.1

- ▶ A k -coloring of a graph G is a labeling $f : V(G) \rightarrow \{1, 2, \dots, k\}$.
- ▶ The labels are **colors**.
- ▶ The vertices with color i are a **color class**.
- ▶ A k -coloring f is **proper** if $f(x) \neq f(y)$ whenever x and y are adjacent.
- ▶ The **chromatic number** $\chi(G)$ is the minimum k such that G is k -colorable.
- ▶ If $\chi(G) = k$, then G is **k -chromatic**.
- ▶ If $\chi(G) = k$, but $\chi(H) < k$ for every proper subgraph H of G , then G is **k -color-critical** or **k -critical**.
- ▶ Let $\omega(G)$ denote the **clique number** of G , that is, the order of a largest complete subgraph of G .

Definition 7.1

- ▶ A k -coloring of a graph G is a labeling $f : V(G) \rightarrow \{1, 2, \dots, k\}$.
- ▶ The labels are **colors**.
- ▶ The vertices with color i are a **color class**.
- ▶ A k -coloring f is **proper** if $f(x) \neq f(y)$ whenever x and y are adjacent.
- ▶ The **chromatic number** $\chi(G)$ is the minimum k such that G is k -colorable.
- ▶ If $\chi(G) = k$, then G is **k -chromatic**.
- ▶ If $\chi(G) = k$, but $\chi(H) < k$ for every proper subgraph H of G , then G is **k -color-critical** or **k -critical**.
- ▶ Let $\omega(G)$ denote the **clique number** of G , that is, the order of a largest complete subgraph of G .
- ▶ Let $\alpha(G)$ denote the **independence number** of G , that is, the largest number of vertices of G no two of which are adjacent.

Facts About Chromatic Number

- ▶ For vertex coloring, all graphs will be considered simple.

Facts About Chromatic Number

- ▶ For vertex coloring, all graphs will be considered simple.
- ▶ $\chi(G) \geq \omega(G)$, and $\chi(G) = \omega(G)$ when G is complete.

Facts About Chromatic Number

- ▶ For vertex coloring, all graphs will be considered simple.
- ▶ $\chi(G) \geq \omega(G)$, and $\chi(G) = \omega(G)$ when G is complete.
- ▶ $\chi(G)$ may exceed $\omega(G)$,

Facts About Chromatic Number

- ▶ For vertex coloring, all graphs will be considered simple.
- ▶ $\chi(G) \geq \omega(G)$, and $\chi(G) = \omega(G)$ when G is complete.
- ▶ $\chi(G)$ may exceed $\omega(G)$, for example, consider $C_{2r+1} \vee K_s$, that is, the graph formed from the disjoint union of C_{2r+1} and K_s by joining each vertex of C_{2r+1} to each vertex of K_s .

Facts About Chromatic Number

- ▶ For vertex coloring, all graphs will be considered simple.
- ▶ $\chi(G) \geq \omega(G)$, and $\chi(G) = \omega(G)$ when G is complete.
- ▶ $\chi(G)$ may exceed $\omega(G)$, for example, consider $C_{2r+1} \vee K_s$, that is, the graph formed from the disjoint union of C_{2r+1} and K_s by joining each vertex of C_{2r+1} to each vertex of K_s .
- ▶ There are triangle-free graphs of arbitrarily high chromatic number (Mycielski)

Facts About Chromatic Number

- ▶ For vertex coloring, all graphs will be considered simple.
- ▶ $\chi(G) \geq \omega(G)$, and $\chi(G) = \omega(G)$ when G is complete.
- ▶ $\chi(G)$ may exceed $\omega(G)$, for example, consider $C_{2r+1} \vee K_s$, that is, the graph formed from the disjoint union of C_{2r+1} and K_s by joining each vertex of C_{2r+1} to each vertex of K_s .
- ▶ There are triangle-free graphs of arbitrarily high chromatic number (Mycielski)
 - ▶ Let $M_2 = K_2$.

Facts About Chromatic Number

- ▶ For vertex coloring, all graphs will be considered simple.
- ▶ $\chi(G) \geq \omega(G)$, and $\chi(G) = \omega(G)$ when G is complete.
- ▶ $\chi(G)$ may exceed $\omega(G)$, for example, consider $C_{2r+1} \vee K_s$, that is, the graph formed from the disjoint union of C_{2r+1} and K_s by joining each vertex of C_{2r+1} to each vertex of K_s .
- ▶ There are triangle-free graphs of arbitrarily high chromatic number (Mycielski)
 - ▶ Let $M_2 = K_2$.
 - ▶ Construct M_{n+1} from M_n by:

Facts About Chromatic Number

- ▶ For vertex coloring, all graphs will be considered simple.
- ▶ $\chi(G) \geq \omega(G)$, and $\chi(G) = \omega(G)$ when G is complete.
- ▶ $\chi(G)$ may exceed $\omega(G)$, for example, consider $C_{2r+1} \vee K_s$, that is, the graph formed from the disjoint union of C_{2r+1} and K_s by joining each vertex of C_{2r+1} to each vertex of K_s .
- ▶ There are triangle-free graphs of arbitrarily high chromatic number (Mycielski)
 - ▶ Let $M_2 = K_2$.
 - ▶ Construct M_{n+1} from M_n by:
 - ▶ Start with a copy M of M_n ;

Facts About Chromatic Number

- ▶ For vertex coloring, all graphs will be considered simple.
- ▶ $\chi(G) \geq \omega(G)$, and $\chi(G) = \omega(G)$ when G is complete.
- ▶ $\chi(G)$ may exceed $\omega(G)$, for example, consider $C_{2r+1} \vee K_s$, that is, the graph formed from the disjoint union of C_{2r+1} and K_s by joining each vertex of C_{2r+1} to each vertex of K_s .
- ▶ There are triangle-free graphs of arbitrarily high chromatic number (Mycielski)
 - ▶ Let $M_2 = K_2$.
 - ▶ Construct M_{n+1} from M_n by:
 - ▶ Start with a copy M of M_n ;
 - ▶ for each vertex v in M add a new vertex u and connect it to the neighbors of v in M ;

Facts About Chromatic Number

- ▶ For vertex coloring, all graphs will be considered simple.
- ▶ $\chi(G) \geq \omega(G)$, and $\chi(G) = \omega(G)$ when G is complete.
- ▶ $\chi(G)$ may exceed $\omega(G)$, for example, consider $C_{2r+1} \vee K_s$, that is, the graph formed from the disjoint union of C_{2r+1} and K_s by joining each vertex of C_{2r+1} to each vertex of K_s .
- ▶ There are triangle-free graphs of arbitrarily high chromatic number (Mycielski)
 - ▶ Let $M_2 = K_2$.
 - ▶ Construct M_{n+1} from M_n by:
 - ▶ Start with a copy M of M_n ;
 - ▶ for each vertex v in M add a new vertex u and connect it to the neighbors of v in M ;
 - ▶ add a new vertex w and connect it to all vertices not in M .

Facts About Chromatic Number

- ▶ For vertex coloring, all graphs will be considered simple.
- ▶ $\chi(G) \geq \omega(G)$, and $\chi(G) = \omega(G)$ when G is complete.
- ▶ $\chi(G)$ may exceed $\omega(G)$, for example, consider $C_{2r+1} \vee K_s$, that is, the graph formed from the disjoint union of C_{2r+1} and K_s by joining each vertex of C_{2r+1} to each vertex of K_s .
- ▶ There are triangle-free graphs of arbitrarily high chromatic number (Mycielski)
 - ▶ Let $M_2 = K_2$.
 - ▶ Construct M_{n+1} from M_n by:
 - ▶ Start with a copy M of M_n ;
 - ▶ for each vertex v in M add a new vertex u and connect it to the neighbors of v in M ;
 - ▶ add a new vertex w and connect it to all vertices not in M .
- ▶ $\chi(G) \geq |G|/\alpha(G)$

Facts About Chromatic Number

- ▶ For vertex coloring, all graphs will be considered simple.
- ▶ $\chi(G) \geq \omega(G)$, and $\chi(G) = \omega(G)$ when G is complete.
- ▶ $\chi(G)$ may exceed $\omega(G)$, for example, consider $C_{2r+1} \vee K_s$, that is, the graph formed from the disjoint union of C_{2r+1} and K_s by joining each vertex of C_{2r+1} to each vertex of K_s .
- ▶ There are triangle-free graphs of arbitrarily high chromatic number (Mycielski)
 - ▶ Let $M_2 = K_2$.
 - ▶ Construct M_{n+1} from M_n by:
 - ▶ Start with a copy M of M_n ;
 - ▶ for each vertex v in M add a new vertex u and connect it to the neighbors of v in M ;
 - ▶ add a new vertex w and connect it to all vertices not in M .
- ▶ $\chi(G) \geq |G|/\alpha(G)$
- ▶ $\chi(G) \leq \Delta(G) + 1$

Proof.

Color greedily:



Facts About Chromatic Number

- ▶ For vertex coloring, all graphs will be considered simple.
- ▶ $\chi(G) \geq \omega(G)$, and $\chi(G) = \omega(G)$ when G is complete.
- ▶ $\chi(G)$ may exceed $\omega(G)$, for example, consider $C_{2r+1} \vee K_s$, that is, the graph formed from the disjoint union of C_{2r+1} and K_s by joining each vertex of C_{2r+1} to each vertex of K_s .
- ▶ There are triangle-free graphs of arbitrarily high chromatic number (Mycielski)
 - ▶ Let $M_2 = K_2$.
 - ▶ Construct M_{n+1} from M_n by:
 - ▶ Start with a copy M of M_n ;
 - ▶ for each vertex v in M add a new vertex u and connect it to the neighbors of v in M ;
 - ▶ add a new vertex w and connect it to all vertices not in M .
- ▶ $\chi(G) \geq |G|/\alpha(G)$
- ▶ $\chi(G) \leq \Delta(G) + 1$

Proof.

Color greedily: Order the vertices arbitrarily as v_1, v_2, \dots, v_n .



Facts About Chromatic Number

- ▶ For vertex coloring, all graphs will be considered simple.
- ▶ $\chi(G) \geq \omega(G)$, and $\chi(G) = \omega(G)$ when G is complete.
- ▶ $\chi(G)$ may exceed $\omega(G)$, for example, consider $C_{2r+1} \vee K_s$, that is, the graph formed from the disjoint union of C_{2r+1} and K_s by joining each vertex of C_{2r+1} to each vertex of K_s .
- ▶ There are triangle-free graphs of arbitrarily high chromatic number (Mycielski)
 - ▶ Let $M_2 = K_2$.
 - ▶ Construct M_{n+1} from M_n by:
 - ▶ Start with a copy M of M_n ;
 - ▶ for each vertex v in M add a new vertex u and connect it to the neighbors of v in M ;
 - ▶ add a new vertex w and connect it to all vertices not in M .
- ▶ $\chi(G) \geq |G|/\alpha(G)$
- ▶ $\chi(G) \leq \Delta(G) + 1$

Proof.

Color greedily: Order the vertices arbitrarily as v_1, v_2, \dots, v_n . Starting with $k = 1$, color each vertex v_k with the smallest color not used among the vertices v_1, v_2, \dots, v_{k-1} that are neighbors of v_k . □

Brooks' Theorem

Theorem 7.2 (Brooks 1941)

If G is a connected simple graph other than a clique and an odd cycle, then $\chi(G) \leq \Delta(G)$.

5-Color Theorem

Theorem 7.3 (Heawood 1890)

Every loopless planar graph has a proper 5-coloring.

5-Color Theorem

Theorem 7.3 (Heawood 1890)

Every loopless planar graph has a proper 5-coloring.

Proof.

Suppose G is a plane graph that is a minimal counter-example.



5-Color Theorem

Theorem 7.3 (Heawood 1890)

Every loopless planar graph has a proper 5-coloring.

Proof.

Suppose G is a plane graph that is a minimal counter-example. Then G is simple, and so $\|G\| \leq 3|G| - 6$ by Corollary 6.15.



5-Color Theorem

Theorem 7.3 (Heawood 1890)

Every loopless planar graph has a proper 5-coloring.

Proof.

Suppose G is a plane graph that is a minimal counter-example. Then G is simple, and so $|E(G)| \leq 3|V(G)| - 6$ by Corollary 6.15. It follows that G has a vertex v of degree 5 or less.



5-Color Theorem

Theorem 7.3 (Heawood 1890)

Every loopless planar graph has a proper 5-coloring.

Proof.

Suppose G is a plane graph that is a minimal counter-example. Then G is simple, and so $|G| \leq 3|G| - 6$ by Corollary 6.15. It follows that G has a vertex v of degree 5 or less. Then $G - v$ has a 5-coloring f by the minimality of G .



5-Color Theorem

Theorem 7.3 (Heawood 1890)

Every loopless planar graph has a proper 5-coloring.

Proof.

Suppose G is a plane graph that is a minimal counter-example. Then G is simple, and so $\|G\| \leq 3|G| - 6$ by Corollary 6.15. It follows that G has a vertex v of degree 5 or less. Then $G - v$ has a 5-coloring f by the minimality of G . Since G is not 5-colorable, each color appears at one of the neighbors of v (and so $d(v) = 5$).



5-Color Theorem

Theorem 7.3 (Heawood 1890)

Every loopless planar graph has a proper 5-coloring.

Proof.

Suppose G is a plane graph that is a minimal counter-example. Then G is simple, and so $|E(G)| \leq 3|V(G)| - 6$ by Corollary 6.15. It follows that G has a vertex v of degree 5 or less. Then $G - v$ has a 5-coloring f by the minimality of G . Since G is not 5-colorable, each color appears at one of the neighbors of v (and so $d(v) = 5$). We may assume that the colors on the neighbors of v appear as 1, 2, 3, 4, and 5 as they are inspected clockwise.



5-Color Theorem

Theorem 7.3 (Heawood 1890)

Every loopless planar graph has a proper 5-coloring.

Proof.

Suppose G is a plane graph that is a minimal counter-example. Then G is simple, and so $\|G\| \leq 3|G| - 6$ by Corollary 6.15. It follows that G has a vertex v of degree 5 or less. Then $G - v$ has a 5-coloring f by the minimality of G . Since G is not 5-colorable, each color appears at one of the neighbors of v (and so $d(v) = 5$). We may assume that the colors on the neighbors of v appear as 1, 2, 3, 4, and 5 as they are inspected clockwise. Let $G_{i,j}$ denote the subgraph of $G - v$ induced by the colors i and j .



5-Color Theorem

Theorem 7.3 (Heawood 1890)

Every loopless planar graph has a proper 5-coloring.

Proof.

Suppose G is a plane graph that is a minimal counter-example. Then G is simple, and so $|E(G)| \leq 3|V(G)| - 6$ by Corollary 6.15. It follows that G has a vertex v of degree 5 or less. Then $G - v$ has a 5-coloring f by the minimality of G . Since G is not 5-colorable, each color appears at one of the neighbors of v (and so $d(v) = 5$). We may assume that the colors on the neighbors of v appear as 1, 2, 3, 4, and 5 as they are inspected clockwise. Let $G_{i,j}$ denote the subgraph of $G - v$ induced by the colors i and j . Note that we can exchange the two colors on any component of $G_{i,j}$ to obtain another proper coloring of $G - v$.



5-Color Theorem

Theorem 7.3 (Heawood 1890)

Every loopless planar graph has a proper 5-coloring.

Proof.

Suppose G is a plane graph that is a minimal counter-example. Then G is simple, and so $|E(G)| \leq 3|V(G)| - 6$ by Corollary 6.15. It follows that G has a vertex v of degree 5 or less. Then $G - v$ has a 5-coloring f by the minimality of G . Since G is not 5-colorable, each color appears at one of the neighbors of v (and so $d(v) = 5$). We may assume that the colors on the neighbors of v appear as 1, 2, 3, 4, and 5 as they are inspected clockwise. Let $G_{i,j}$ denote the subgraph of $G - v$ induced by the colors i and j . Note that we can exchange the two colors on any component of $G_{i,j}$ to obtain another proper coloring of $G - v$. If some two neighbors of v were in different components of $G_{i,j}$, then switching colors on one such component would result in two neighbors of G being colored the same,



5-Color Theorem

Theorem 7.3 (Heawood 1890)

Every loopless planar graph has a proper 5-coloring.

Proof.

Suppose G is a plane graph that is a minimal counter-example. Then G is simple, and so $|E(G)| \leq 3|V(G)| - 6$ by Corollary 6.15. It follows that G has a vertex v of degree 5 or less. Then $G - v$ has a 5-coloring f by the minimality of G . Since G is not 5-colorable, each color appears at one of the neighbors of v (and so $d(v) = 5$). We may assume that the colors on the neighbors of v appear as 1, 2, 3, 4, and 5 as they are inspected clockwise. Let $G_{i,j}$ denote the subgraph of $G - v$ induced by the colors i and j . Note that we can exchange the two colors on any component of $G_{i,j}$ to obtain another proper coloring of $G - v$. If some two neighbors of v were in different components of $G_{i,j}$, then switching colors on one such component would result in two neighbors of G being colored the same, Thus allowing to extend the coloring to v .



5-Color Theorem

Theorem 7.3 (Heawood 1890)

Every loopless planar graph has a proper 5-coloring.

Proof.

Suppose G is a plane graph that is a minimal counter-example. Then G is simple, and so $|E(G)| \leq 3|V(G)| - 6$ by Corollary 6.15. It follows that G has a vertex v of degree 5 or less. Then $G - v$ has a 5-coloring f by the minimality of G . Since G is not 5-colorable, each color appears at one of the neighbors of v (and so $d(v) = 5$). We may assume that the colors on the neighbors of v appear as 1, 2, 3, 4, and 5 as they are inspected clockwise. Let $G_{i,j}$ denote the subgraph of $G - v$ induced by the colors i and j . Note that we can exchange the two colors on any component of $G_{i,j}$ to obtain another proper coloring of $G - v$. If some two neighbors of v were in different components of $G_{i,j}$, then switching colors on one such component would result in two neighbors of G being colored the same. Thus allowing to extend the coloring to v . Thus we may assume that every two neighbors of v are in the same component of $G_{i,j}$. \square

Proof, Continued

Let $P_{i,j}$ be a path in $G_{i,j}$ joining the neighbors of v colored i and j .

Proof, Continued

Let $P_{i,j}$ be a path in $G_{i,j}$ joining the neighbors of v colored i and j . Consider the cycle C of G induced by $P_{1,3}$ together with v ,

Proof, Continued

Let $P_{i,j}$ be a path in $G_{i,j}$ joining the neighbors of v colored i and j . Consider the cycle C of G induced by $P_{1,3}$ together with v , which separates the neighbor of v colored 2 from the one colored 4.

Proof, Continued

Let $P_{i,j}$ be a path in $G_{i,j}$ joining the neighbors of v colored i and j . Consider the cycle C of G induced by $P_{1,3}$ together with v , which separates the neighbor of v colored 2 from the one colored 4. Hence $P_{2,4}$ must cross C , which is impossible.

Proof, Continued

Let $P_{i,j}$ be a path in $G_{i,j}$ joining the neighbors of v colored i and j . Consider the cycle C of G induced by $P_{1,3}$ together with v , which separates the neighbor of v colored 2 from the one colored 4. Hence $P_{2,4}$ must cross C , which is impossible.

Theorem 7.4 (4-Color Theorem, Appel and Haken 1977)

Every loopless planar graph has a proper 4-coloring.

Chromatic Polynomial

Let $P_G(x)$ denote the number of ways to properly color a (labeled) graph G with x colors.

Chromatic Polynomial

Let $P_G(x)$ denote the number of ways to properly color a (labeled) graph G with x colors. If G has loops, then $P_G(x) = 0$.

Chromatic Polynomial

Let $P_G(x)$ denote the number of ways to properly color a (labeled) graph G with x colors. If G has loops, then $P_G(x) = 0$. If G is edgeless of order n , then $P_G(x) = x^n$.

Chromatic Polynomial

Let $P_G(x)$ denote the number of ways to properly color a (labeled) graph G with x colors. If G has loops, then $P_G(x) = 0$. If G is edgeless of order n , then $P_G(x) = x^n$. If e is a non-loop non-multiple edge of G incident with u and v , then the proper colorings of $G \setminus e$ with x colors can be partitioned into two sets: A , in which u and v receive the same color, and B , in which they do not.

Chromatic Polynomial

Let $P_G(x)$ denote the number of ways to properly color a (labeled) graph G with x colors. If G has loops, then $P_G(x) = 0$. If G is edgeless of order n , then $P_G(x) = x^n$. If e is a non-loop non-multiple edge of G incident with u and v , then the proper colorings of $G \setminus e$ with x colors can be partitioned into two sets: A , in which u and v receive the same color, and B , in which they do not. Then A corresponds to proper colorings of G/e with x colors, and B corresponds to proper colorings of G .

Chromatic Polynomial

Let $P_G(x)$ denote the number of ways to properly color a (labeled) graph G with x colors. If G has loops, then $P_G(x) = 0$. If G is edgeless of order n , then $P_G(x) = x^n$. If e is a non-loop non-multiple edge of G incident with u and v , then the proper colorings of $G \setminus e$ with x colors can be partitioned into two sets: A , in which u and v receive the same color, and B , in which they do not. Then A corresponds to proper colorings of G/e with x colors, and B corresponds to proper colorings of G . Hence

$$P_G(x) = P_{G \setminus e}(x) - P_{G/e}(x).$$

Chromatic Polynomial

Let $P_G(x)$ denote the number of ways to properly color a (labeled) graph G with x colors. If G has loops, then $P_G(x) = 0$. If G is edgeless of order n , then $P_G(x) = x^n$. If e is a non-loop non-multiple edge of G incident with u and v , then the proper colorings of $G \setminus e$ with x colors can be partitioned into two sets: A , in which u and v receive the same color, and B , in which they do not. Then A corresponds to proper colorings of G/e with x colors, and B corresponds to proper colorings of G . Hence

$$P_G(x) = P_{G \setminus e}(x) - P_{G/e}(x).$$

$P_G(x)$ is called the **chromatic polynomial** of G .

Chromatic Polynomial

Let $P_G(x)$ denote the number of ways to properly color a (labeled) graph G with x colors. If G has loops, then $P_G(x) = 0$. If G is edgeless of order n , then $P_G(x) = x^n$. If e is a non-loop non-multiple edge of G incident with u and v , then the proper colorings of $G \setminus e$ with x colors can be partitioned into two sets: A , in which u and v receive the same color, and B , in which they do not. Then A corresponds to proper colorings of G/e with x colors, and B corresponds to proper colorings of G . Hence

$$P_G(x) = P_{G \setminus e}(x) - P_{G/e}(x).$$

$P_G(x)$ is called the **chromatic polynomial** of G .

Theorem 7.6 (Four-Color Theorem, restated)

If G is a planar loopless graph, then $P_G(4) > 0$.

Perfect Graphs

Definition 7.7

A graph G is **perfect** if $\chi(H) = \omega(H)$ for every induced subgraph H of G .

Perfect Graphs

Definition 7.7

A graph G is **perfect** if $\chi(H) = \omega(H)$ for every induced subgraph H of G .

Theorem 7.8 (Perfect Graph Theorem, Lovász 1972)

A graph is perfect if and only if its complement is perfect.

Perfect Graphs

Definition 7.7

A graph G is **perfect** if $\chi(H) = \omega(H)$ for every induced subgraph H of G .

Theorem 7.8 (Perfect Graph Theorem, Lovász 1972)

A graph is perfect if and only if its complement is perfect.

Theorem 7.9 (Strong Graph Theorem (formerly Berge's Strong Graph Conjecture), Chudnovsky, Robertson, Seymour, Thomas 2002)

A graph is perfect if and only if it has no induced subgraph that is an odd cycle of length at least five or its complement.

Definition 7.10

- ▶ A k -edge-coloring of a graph G is a labeling $f : E(G) \rightarrow \{1, 2, \dots, k\}$.

Edge Colorings

Definition 7.10

- ▶ A k -edge-coloring of a graph G is a labeling $f : E(G) \rightarrow \{1, 2, \dots, k\}$.
- ▶ The labels are colors and the edge-set with one color is a color class.

Edge Colorings

Definition 7.10

- ▶ A k -edge-coloring of a graph G is a labeling $f : E(G) \rightarrow \{1, 2, \dots, k\}$.
- ▶ The labels are colors and the edge-set with one color is a color class.
- ▶ A k -edge-coloring is proper if adjacent edges have different colors,

Edge Colorings

Definition 7.10

- ▶ A **k -edge-coloring** of a graph G is a labeling $f : E(G) \rightarrow \{1, 2, \dots, k\}$.
- ▶ The labels are **colors** and the edge-set with one color is a **color class**.
- ▶ A k -edge-coloring is **proper** if adjacent edges have different colors, or equivalently, if every color class is a matching.

Definition 7.10

- ▶ A k -edge-coloring of a graph G is a labeling $f : E(G) \rightarrow \{1, 2, \dots, k\}$.
- ▶ The labels are colors and the edge-set with one color is a color class.
- ▶ A k -edge-coloring is proper if adjacent edges have different colors, or equivalently, if every color class is a matching.
- ▶ A graph is k -edge-colorable if it has a proper k -edge-coloring.

Definition 7.10

- ▶ A **k -edge-coloring** of a graph G is a labeling $f : E(G) \rightarrow \{1, 2, \dots, k\}$.
- ▶ The labels are **colors** and the edge-set with one color is a **color class**.
- ▶ A k -edge-coloring is **proper** if adjacent edges have different colors, or equivalently, if every color class is a matching.
- ▶ A graph is **k -edge-colorable** if it has a proper k -edge-coloring.
- ▶ The **chromatic index** or **edge chromatic number** $\chi'(G)$ of a loopless graph G is the least k such that G is k -edge-colorable.

Edge Colorings

Definition 7.10

- ▶ A **k -edge-coloring** of a graph G is a labeling $f : E(G) \rightarrow \{1, 2, \dots, k\}$.
- ▶ The labels are **colors** and the edge-set with one color is a **color class**.
- ▶ A k -edge-coloring is **proper** if adjacent edges have different colors, or equivalently, if every color class is a matching.
- ▶ A graph is **k -edge-colorable** if it has a proper k -edge-coloring.
- ▶ The **chromatic index** or **edge chromatic number** $\chi'(G)$ of a loopless graph G is the least k such that G is k -edge-colorable.

Note 7.11

$$\Delta(G) \leq \chi'(G).$$

Edge-Coloring of Bipartite Graphs

Theorem 7.12 (König 1916)

If G is bipartite, then $\chi'(G) = \Delta(G)$.

Vizing's Theorem

Theorem 7.13 (Vizing 1964–65, Gupta 1966)

If G is simple, then $\chi'(G) \leq \Delta(G) + 1$.

List Colorings

Suppose G is a graph with the vertex set V , and $\mathcal{L} = (L_v)_{v \in V}$ associates with each vertex v a list L_v of colors available to color v .

List Colorings

Suppose G is a graph with the vertex set V , and $\mathcal{L} = (L_v)_{v \in V}$ associates with each vertex v a list L_v of colors available to color v . We say that G **admits an \mathcal{L} -coloring** if there is a proper coloring of G such that, for every vertex v , the color of v is in the list L_v .

List Colorings

Suppose G is a graph with the vertex set V , and $\mathcal{L} = (L_v)_{v \in V}$ associates with each vertex v a list L_v of colors available to color v . We say that G **admits an \mathcal{L} -coloring** if there is a proper coloring of G such that, for every vertex v , the color of v is in the list L_v . The graph G is **k -list-colorable** or **k -choosable** if G admits an \mathcal{L} -coloring for every $\mathcal{L} = (L_v)_{v \in V}$ with $|L_v| = k$ for every vertex v .

List Colorings

Suppose G is a graph with the vertex set V , and $\mathcal{L} = (L_v)_{v \in V}$ associates with each vertex v a list L_v of colors available to color v . We say that G **admits an \mathcal{L} -coloring** if there is a proper coloring of G such that, for every vertex v , the color of v is in the list L_v . The graph G is **k -list-colorable** or **k -choosable** if G admits an \mathcal{L} -coloring for every $\mathcal{L} = (L_v)_{v \in V}$ with $|L_v| = k$ for every vertex v . The smallest k such that G is k -choosable is called the **list-chromatic number of G** and is denoted by **$\text{ch}(G)$** .

List Colorings

Suppose G is a graph with the vertex set V , and $\mathcal{L} = (L_v)_{v \in V}$ associates with each vertex v a list L_v of colors available to color v . We say that G **admits an \mathcal{L} -coloring** if there is a proper coloring of G such that, for every vertex v , the color of v is in the list L_v . The graph G is **k -list-colorable** or **k -choosable** if G admits an \mathcal{L} -coloring for every $\mathcal{L} = (L_v)_{v \in V}$ with $|L_v| = k$ for every vertex v . The smallest k such that G is k -choosable is called the **list-chromatic number of G** and is denoted by **$\text{ch}(G)$** . **List colorings** of edges are defined analogously, as is the **list-chromatic index $\text{ch}'(G)$** .

List Colorings

Suppose G is a graph with the vertex set V , and $\mathcal{L} = (L_v)_{v \in V}$ associates with each vertex v a list L_v of colors available to color v . We say that G **admits an \mathcal{L} -coloring** if there is a proper coloring of G such that, for every vertex v , the color of v is in the list L_v . The graph G is **k -list-colorable** or **k -choosable** if G admits an \mathcal{L} -coloring for every $\mathcal{L} = (L_v)_{v \in V}$ with $|L_v| = k$ for every vertex v . The smallest k such that G is k -choosable is called the **list-chromatic number of G** and is denoted by **$\text{ch}(G)$** .

List colorings of edges are defined analogously, as is the **list-chromatic index $\text{ch}'(G)$** . Note that if $\mathcal{L} = (L_v)_{v \in V}$ is such that all L_v 's are identical and of cardinality k , then G admitting an \mathcal{L} -coloring is equivalent to G being k -colorable.

List Colorings

Suppose G is a graph with the vertex set V , and $\mathcal{L} = (L_v)_{v \in V}$ associates with each vertex v a list L_v of colors available to color v . We say that G **admits an \mathcal{L} -coloring** if there is a proper coloring of G such that, for every vertex v , the color of v is in the list L_v . The graph G is **k -list-colorable** or **k -choosable** if G admits an \mathcal{L} -coloring for every $\mathcal{L} = (L_v)_{v \in V}$ with $|L_v| = k$ for every vertex v . The smallest k such that G is k -choosable is called the **list-chromatic number of G** and is denoted by **$\text{ch}(G)$** .

List colorings of edges are defined analogously, as is the **list-chromatic index $\text{ch}'(G)$** . Note that if $\mathcal{L} = (L_v)_{v \in V}$ is such that all L_v 's are identical and of cardinality k , then G admitting an \mathcal{L} -coloring is equivalent to G being k -colorable. An analogous statement holds for edge-colorings.

List Colorings

Suppose G is a graph with the vertex set V , and $\mathcal{L} = (L_v)_{v \in V}$ associates with each vertex v a list L_v of colors available to color v . We say that G **admits an \mathcal{L} -coloring** if there is a proper coloring of G such that, for every vertex v , the color of v is in the list L_v . The graph G is **k -list-colorable** or **k -choosable** if G admits an \mathcal{L} -coloring for every $\mathcal{L} = (L_v)_{v \in V}$ with $|L_v| = k$ for every vertex v . The smallest k such that G is k -choosable is called the **list-chromatic number of G** and is denoted by **$\text{ch}(G)$** .

List colorings of edges are defined analogously, as is the **list-chromatic index $\text{ch}'(G)$** . Note that if $\mathcal{L} = (L_v)_{v \in V}$ is such that all L_v 's are identical and of cardinality k , then G admitting an \mathcal{L} -coloring is equivalent to G being k -colorable. An analogous statement holds for edge-colorings. Thus

$$\text{ch}(G) \geq \chi(G) \quad \text{and} \quad \text{ch}'(G) \geq \chi'(G).$$

List Colorings

Suppose G is a graph with the vertex set V , and $\mathcal{L} = (L_v)_{v \in V}$ associates with each vertex v a list L_v of colors available to color v . We say that G **admits an \mathcal{L} -coloring** if there is a proper coloring of G such that, for every vertex v , the color of v is in the list L_v . The graph G is **k -list-colorable** or **k -choosable** if G admits an \mathcal{L} -coloring for every $\mathcal{L} = (L_v)_{v \in V}$ with $|L_v| = k$ for every vertex v . The smallest k such that G is k -choosable is called the **list-chromatic number of G** and is denoted by **$\text{ch}(G)$** .

List colorings of edges are defined analogously, as is the **list-chromatic index $\text{ch}'(G)$** . Note that if $\mathcal{L} = (L_v)_{v \in V}$ is such that all L_v 's are identical and of cardinality k , then G admitting an \mathcal{L} -coloring is equivalent to G being k -colorable. An analogous statement holds for edge-colorings. Thus

$$\text{ch}(G) \geq \chi(G) \quad \text{and} \quad \text{ch}'(G) \geq \chi'(G).$$

But there are graphs for which $\text{ch}(G) \neq \chi(G)$.

List Colorings

Suppose G is a graph with the vertex set V , and $\mathcal{L} = (L_v)_{v \in V}$ associates with each vertex v a list L_v of colors available to color v . We say that G **admits an \mathcal{L} -coloring** if there is a proper coloring of G such that, for every vertex v , the color of v is in the list L_v . The graph G is **k -list-colorable** or **k -choosable** if G admits an \mathcal{L} -coloring for every $\mathcal{L} = (L_v)_{v \in V}$ with $|L_v| = k$ for every vertex v . The smallest k such that G is k -choosable is called the **list-chromatic number of G** and is denoted by **$\text{ch}(G)$** .

List colorings of edges are defined analogously, as is the **list-chromatic index $\text{ch}'(G)$** . Note that if $\mathcal{L} = (L_v)_{v \in V}$ is such that all L_v 's are identical and of cardinality k , then G admitting an \mathcal{L} -coloring is equivalent to G being k -colorable. An analogous statement holds for edge-colorings. Thus

$$\text{ch}(G) \geq \chi(G) \quad \text{and} \quad \text{ch}'(G) \geq \chi'(G).$$

But there are graphs for which $\text{ch}(G) \neq \chi(G)$. Consider $K_{3,3}$ where each side of the bipartition has lists $\{1, 2\}$, $\{1, 3\}$, and $\{2, 3\}$.

List Colorings

Suppose G is a graph with the vertex set V , and $\mathcal{L} = (L_v)_{v \in V}$ associates with each vertex v a list L_v of colors available to color v . We say that G **admits an \mathcal{L} -coloring** if there is a proper coloring of G such that, for every vertex v , the color of v is in the list L_v . The graph G is **k -list-colorable** or **k -choosable** if G admits an \mathcal{L} -coloring for every $\mathcal{L} = (L_v)_{v \in V}$ with $|L_v| = k$ for every vertex v . The smallest k such that G is k -choosable is called the **list-chromatic number of G** and is denoted by **$\text{ch}(G)$** .

List colorings of edges are defined analogously, as is the **list-chromatic index $\text{ch}'(G)$** . Note that if $\mathcal{L} = (L_v)_{v \in V}$ is such that all L_v 's are identical and of cardinality k , then G admitting an \mathcal{L} -coloring is equivalent to G being k -colorable. An analogous statement holds for edge-colorings. Thus

$$\text{ch}(G) \geq \chi(G) \quad \text{and} \quad \text{ch}'(G) \geq \chi'(G).$$

But there are graphs for which $\text{ch}(G) \neq \chi(G)$. Consider $K_{3,3}$ where each side of the bipartition has lists $\{1, 2\}$, $\{1, 3\}$, and $\{2, 3\}$. The list-chromatic number of this graph is 3, while the chromatic number is 2.

Every Planar Graph Is 5-Choosable

Theorem 7.14 (Thomassen 1994)

Every planar graph is 5-choosable.