

# Polygonal Paths

## Note 6.1

*A graph may be viewed as a **topological space** when*

# Polygonal Paths

## Note 6.1

*A graph may be viewed as a **topological space** when*

- ▶ *the vertices are points;*

# Polygonal Paths

## Note 6.1

A graph may be viewed as a *topological space* when

- ▶ the vertices are points;
- ▶ the edges are homeomorphic images of the unit interval; and

# Polygonal Paths

## Note 6.1

A graph may be viewed as a *topological space* when

- ▶ the vertices are points;
- ▶ the edges are homeomorphic images of the unit interval; and
- ▶ the incidences are reflected by the vertices (points) being in the closure of the edges.

# Polygonal Paths

## Note 6.1

A graph may be viewed as a *topological space* when

- ▶ the vertices are points;
- ▶ the edges are homeomorphic images of the unit interval; and
- ▶ the incidences are reflected by the vertices (points) being in the closure of the edges.

## Definition 6.2

A **polygonal curve** in the plane is the union of finitely many line segments such that each segment starts at the end of the previous one and no point lies in more than one segment, except the end of one segment and the beginning of the next one coincide.

A **simple open** polygonal curve is one homeomorphic to a closed interval.

A **simple closed** polygonal curve is one homeomorphic to a unit circle.

## Definition 6.3

- ▶ A **drawing** of a graph  $G$  is a function that maps each vertex  $v \in V(G)$  to a point  $f(v)$  in the plane, and each  $uv$ -edge to a simple polygonal  $f(u)f(v)$ -curve in the plane.

## Definition 6.3

- ▶ A **drawing** of a graph  $G$  is a function that maps each vertex  $v \in V(G)$  to a point  $f(v)$  in the plane, and each  $uv$ -edge to a simple polygonal  $f(u)f(v)$ -curve in the plane.
- ▶ A point  $f(e) \cap f(e')$  other than the a common endpoint is a **crossing**.

## Definition 6.3

- ▶ A **drawing** of a graph  $G$  is a function that maps each vertex  $v \in V(G)$  to a point  $f(v)$  in the plane, and each  $uv$ -edge to a simple polygonal  $f(u)f(v)$ -curve in the plane.
- ▶ A point  $f(e) \cap f(e')$  other than the a common endpoint is a **crossing**.
- ▶ A graph is **planar** if it has a drawing without crossings.



## Definition 6.3

- ▶ A **drawing** of a graph  $G$  is a function that maps each vertex  $v \in V(G)$  to a point  $f(v)$  in the plane, and each  $uv$ -edge to a simple polygonal  $f(u)f(v)$ -curve in the plane.
- ▶ A point  $f(e) \cap f(e')$  other than the a common endpoint is a **crossing**.
- ▶ A graph is **planar** if it has a drawing without crossings. Such a drawing is a **planar embedding** of  $G$ .

## Definition 6.3

- ▶ A **drawing** of a graph  $G$  is a function that maps each vertex  $v \in V(G)$  to a point  $f(v)$  in the plane, and each  $uv$ -edge to a simple polygonal  $f(u)f(v)$ -curve in the plane.
- ▶ A point  $f(e) \cap f(e')$  other than the a common endpoint is a **crossing**.
- ▶ A graph is **planar** if it has a drawing without crossings. Such a drawing is a **planar embedding** of  $G$ .
- ▶ A **plane graph** is a particular drawing of a a graph in the plane with no crossings.

## Definition 6.3

- ▶ A **drawing** of a graph  $G$  is a function that maps each vertex  $v \in V(G)$  to a point  $f(v)$  in the plane, and each  $uv$ -edge to a simple polygonal  $f(u)f(v)$ -curve in the plane.
- ▶ A point  $f(e) \cap f(e')$  other than the a common endpoint is a **crossing**.
- ▶ A graph is **planar** if it has a drawing without crossings. Such a drawing is a **planar embedding** of  $G$ .
- ▶ A **plane graph** is a particular drawing of a a graph in the plane with no crossings.

## Note 6.4

*A plane embedding corresponds to an embedding of the graph in the sphere through a **stereographic projection**.*

## Theorem 6.5 (Jordan Curve Theorem)

*If  $C$  is a simple closed polygonal curve in the plane, then the complement of  $C$  in the plane consists of two connected components each with  $C$  as the boundary.*

## Theorem 6.5 (Jordan Curve Theorem)

*If  $C$  is a simple closed polygonal curve in the plane, then the complement of  $C$  in the plane consists of two connected components each with  $C$  as the boundary.*

## Definition 6.6

- ▶ The connected components of the complement of a plane graph are the **faces** of the embedding.

## Theorem 6.5 (Jordan Curve Theorem)

*If  $C$  is a simple closed polygonal curve in the plane, then the complement of  $C$  in the plane consists of two connected components each with  $C$  as the boundary.*

## Definition 6.6

- ▶ The connected components of the complement of a plane graph are the **faces** of the embedding.
- ▶ The **length** of a face is the number of edges in the boundary of the face, with cut-edges counted twice.

## Theorem 6.5 (Jordan Curve Theorem)

*If  $C$  is a simple closed polygonal curve in the plane, then the complement of  $C$  in the plane consists of two connected components each with  $C$  as the boundary.*

## Definition 6.6

- ▶ The connected components of the complement of a plane graph are the **faces** of the embedding.
- ▶ The **length** of a face is the number of edges in the boundary of the face, with cut-edges counted twice.
- ▶ The **dual graph**  $G^*$  of a non-empty plane graph  $G$  is the graph such that
  - ▶ the vertices of  $G^*$  are the faces of  $G$ ;

## Theorem 6.5 (Jordan Curve Theorem)

*If  $C$  is a simple closed polygonal curve in the plane, then the complement of  $C$  in the plane consists of two connected components each with  $C$  as the boundary.*

## Definition 6.6

- ▶ The connected components of the complement of a plane graph are the **faces** of the embedding.
- ▶ The **length** of a face is the number of edges in the boundary of the face, with cut-edges counted twice.
- ▶ The **dual graph**  $G^*$  of a non-empty plane graph  $G$  is the graph such that
  - ▶ the vertices of  $G^*$  are the faces of  $G$ ;
  - ▶ the edges of  $G^*$  are the edges of  $G$ ;



## Theorem 6.5 (Jordan Curve Theorem)


*If  $C$  is a simple closed polygonal curve in the plane, then the complement of  $C$  in the plane consists of two connected components each with  $C$  as the boundary.*

## Definition 6.6

- ▶ The connected components of the complement of a plane graph are the **faces** of the embedding.
- ▶ The **length** of a face is the number of edges in the boundary of the face, with cut-edges counted twice.
- ▶ The **dual graph**  $G^*$  of a non-empty plane graph  $G$  is the graph such that
  - ▶ the vertices of  $G^*$  are the faces of  $G$ ;
  - ▶ the edges of  $G^*$  are the edges of  $G$ ;
  - ▶ a vertex and an edge of  $G^*$  are incident if and only if the edge is the boundary of the corresponding face of  $G$ .


## Example of a Dual Graph

Example 6.7




## Example of a Dual Graph

Example 6.7



## Example of a Dual Graph

Example 6.7



# Properties of Dual Graphs

## Note 6.8

*Different graphs may have the same dual.*

# Properties of Dual Graphs

## Note 6.8

*Different graphs may have the same dual.  $(G^*)^* = G$  if and only if  $G$  is connected.*

# Properties of Dual Graphs

## Note 6.8

*Different graphs may have the same dual.  $(G^*)^* = G$  if and only if  $G$  is connected.*

## Theorem 6.9

*Edges in a plane graph form a cycle if and only if the edges in the dual graph form a bond.*

## Properties of Dual Graphs, Continued

### Theorem 6.10

*If a plane graph  $G$  is connected, then the Tutte connectivity of  $G$  is the same as the Tutte connectivity of  $G^*$ .*



## Properties of Dual Graphs, Continued

### Theorem 6.10

*If a plane graph  $G$  is connected, then the Tutte connectivity of  $G$  is the same as the Tutte connectivity of  $G^*$ .*

### Theorem 6.11

*The following are equivalent for a plane graph  $G$ :*

## Properties of Dual Graphs, Continued

### Theorem 6.10

*If a plane graph  $G$  is connected, then the Tutte connectivity of  $G$  is the same as the Tutte connectivity of  $G^*$ .*

### Theorem 6.11

*The following are equivalent for a plane graph  $G$ :*

(A)  $G$  is bipartite;

## Properties of Dual Graphs, Continued

### Theorem 6.10

*If a plane graph  $G$  is connected, then the Tutte connectivity of  $G$  is the same as the Tutte connectivity of  $G^*$ .*

### Theorem 6.11

*The following are equivalent for a plane graph  $G$ :*

- (A)  *$G$  is bipartite;*
- (B) *every face of  $G$  has even length;*

## Properties of Dual Graphs, Continued

### Theorem 6.10

*If a plane graph  $G$  is connected, then the Tutte connectivity of  $G$  is the same as the Tutte connectivity of  $G^*$ .*

### Theorem 6.11

*The following are equivalent for a plane graph  $G$ :*

- (A)  $G$  is bipartite;
- (B) every face of  $G$  has even length;
- (C)  $G^*$  is Eulerian.

## Note 6.12

- ▶ *Deleting an edge or a vertex from a plane graph results in a plane graph.*

## Note 6.12

- ▶ *Deleting an edge or a vertex from a plane graph results in a plane graph.*
- ▶ *Contracting an edge in a plane graph can be visualized as sliding the two endvertices towards each other until they meet, pulling all incident edges along.*

## Note 6.12

- ▶ *Deleting an edge or a vertex from a plane graph results in a plane graph.*
- ▶ *Contracting an edge in a plane graph can be visualized as sliding the two endvertices towards each other until they meet, pulling all incident edges along.*
- ▶ *Thus the class of planar graphs is **minor-closed**, that is, all minors of planar graphs are also planar.*

# Euler's Formula

## Theorem 6.13 (Euler's Formula)

*If a connected non-empty plane graph has  $v$  vertices,  $e$  edges, and  $f$  faces, then  $v - e + f = 2$ .*



# Euler's Formula

## Theorem 6.13 (Euler's Formula)

*If a connected non-empty plane graph has  $v$  vertices,  $e$  edges, and  $f$  faces, then  $v - e + f = 2$ .*

## Proof.

We proceed by induction on  $v$ .



# Euler's Formula

## Theorem 6.13 (Euler's Formula)

*If a connected non-empty plane graph has  $v$  vertices,  $e$  edges, and  $f$  faces, then  $v - e + f = 2$ .*

## Proof.

We proceed by induction on  $v$ . If  $v = 1$ , then  $G$  has only loops, each a closed curve in the embedding.



# Euler's Formula

## Theorem 6.13 (Euler's Formula)

*If a connected non-empty plane graph has  $v$  vertices,  $e$  edges, and  $f$  faces, then  $v - e + f = 2$ .*

### Proof.

We proceed by induction on  $v$ . If  $v = 1$ , then  $G$  has only loops, each a closed curve in the embedding. If  $e = 0$ , then  $f = 1$ , and the formula holds.



# Euler's Formula

## Theorem 6.13 (Euler's Formula)

*If a connected non-empty plane graph has  $v$  vertices,  $e$  edges, and  $f$  faces, then  $v - e + f = 2$ .*

### Proof.

We proceed by induction on  $v$ . If  $v = 1$ , then  $G$  has only loops, each a closed curve in the embedding. If  $e = 0$ , then  $f = 1$ , and the formula holds. Each added loop adds one more edge and one more face, and so the formula holds when  $v = 1$ .



# Euler's Formula

## Theorem 6.13 (Euler's Formula)

*If a connected non-empty plane graph has  $v$  vertices,  $e$  edges, and  $f$  faces, then  $v - e + f = 2$ .*

### Proof.

We proceed by induction on  $v$ . If  $v = 1$ , then  $G$  has only loops, each a closed curve in the embedding. If  $e = 0$ , then  $f = 1$ , and the formula holds. Each added loop adds one more edge and one more face, and so the formula holds when  $v = 1$ .

Suppose  $v > 1$ .



# Euler's Formula

## Theorem 6.13 (Euler's Formula)

*If a connected non-empty plane graph has  $v$  vertices,  $e$  edges, and  $f$  faces, then  $v - e + f = 2$ .*

### Proof.

We proceed by induction on  $v$ . If  $v = 1$ , then  $G$  has only loops, each a closed curve in the embedding. If  $e = 0$ , then  $f = 1$ , and the formula holds. Each added loop adds one more edge and one more face, and so the formula holds when  $v = 1$ .

Suppose  $v > 1$ . Since  $G$  is connected, it has a non-loop edge.



# Euler's Formula

## Theorem 6.13 (Euler's Formula)

*If a connected non-empty plane graph has  $v$  vertices,  $e$  edges, and  $f$  faces, then  $v - e + f = 2$ .*

### Proof.

We proceed by induction on  $v$ . If  $v = 1$ , then  $G$  has only loops, each a closed curve in the embedding. If  $e = 0$ , then  $f = 1$ , and the formula holds. Each added loop adds one more edge and one more face, and so the formula holds when  $v = 1$ .

Suppose  $v > 1$ . Since  $G$  is connected, it has a non-loop edge. Contract such an edge to obtain a plane graph with  $v' = v - 1$  vertices,  $e' = e - 1$  edges, and  $f' = f$  faces.



# Euler's Formula

## Theorem 6.13 (Euler's Formula)

*If a connected non-empty plane graph has  $v$  vertices,  $e$  edges, and  $f$  faces, then  $v - e + f = 2$ .*

### Proof.

We proceed by induction on  $v$ . If  $v = 1$ , then  $G$  has only loops, each a closed curve in the embedding. If  $e = 0$ , then  $f = 1$ , and the formula holds. Each added loop adds one more edge and one more face, and so the formula holds when  $v = 1$ .

Suppose  $v > 1$ . Since  $G$  is connected, it has a non-loop edge. Contract such an edge to obtain a plane graph with  $v' = v - 1$  vertices,  $e' = e - 1$  edges, and  $f' = f$  faces. Applying the inductive hypothesis, we get  $v' - e' + f' = 2$ ,





# Euler's Formula

## Theorem 6.13 (Euler's Formula)

*If a connected non-empty plane graph has  $v$  vertices,  $e$  edges, and  $f$  faces, then  $v - e + f = 2$ .*

### Proof.

We proceed by induction on  $v$ . If  $v = 1$ , then  $G$  has only loops, each a closed curve in the embedding. If  $e = 0$ , then  $f = 1$ , and the formula holds. Each added loop adds one more edge and one more face, and so the formula holds when  $v = 1$ .

Suppose  $v > 1$ . Since  $G$  is connected, it has a non-loop edge. Contract such an edge to obtain a plane graph with  $v' = v - 1$  vertices,  $e' = e - 1$  edges, and  $f' = f$  faces. Applying the inductive hypothesis, we get  $v' - e' + f' = 2$ , and so  $(v - 1) - (e - 1) + (f) = v - e + f = 2$ , as desired.  $\square$

## Corollaries of Euler's Formula

### Note 6.14

- ▶ *Euler's Formula implies that all plane embeddings of connected graphs have the same number of faces.*

# Corollaries of Euler's Formula

## Note 6.14

- ▶ *Euler's Formula implies that all plane embeddings of connected graphs have the same number of faces.*
- ▶ *Contracting a non-loop edge of  $G$  has the effect of deleting the corresponding edge in  $G^*$ . Similarly, deleting a non-cut edge of  $G$  has the effect of contracting the corresponding edge in  $G^*$ .*

# Corollaries of Euler's Formula

## Note 6.14

- ▶ *Euler's Formula implies that all plane embeddings of connected graphs have the same number of faces.*
- ▶ *Contracting a non-loop edge of  $G$  has the effect of deleting the corresponding edge in  $G^*$ . Similarly, deleting a non-cut edge of  $G$  has the effect of contracting the corresponding edge in  $G^*$ .*
- ▶ *Euler's Formula (as stated) fails for disconnected graphs.*

# Corollaries of Euler's Formula

## Note 6.14

- ▶ *Euler's Formula implies that all plane embeddings of connected graphs have the same number of faces.*
- ▶ *Contracting a non-loop edge of  $G$  has the effect of deleting the corresponding edge in  $G^*$ . Similarly, deleting a non-cut edge of  $G$  has the effect of contracting the corresponding edge in  $G^*$ .*
- ▶ *Euler's Formula (as stated) fails for disconnected graphs.*

## Corollary 6.15

*If  $G$  is a planar graph whose order  $v$  is at least 3, whose size is  $e$ , and whose girth  $g$  is at least 3 but finite, then*

$$e \leq \frac{(v-2)g}{g-2}.$$

# Corollaries of Euler's Formula

## Note 6.14

- ▶ *Euler's Formula implies that all plane embeddings of connected graphs have the same number of faces.*
- ▶ *Contracting a non-loop edge of  $G$  has the effect of deleting the corresponding edge in  $G^*$ . Similarly, deleting a non-cut edge of  $G$  has the effect of contracting the corresponding edge in  $G^*$ .*
- ▶ *Euler's Formula (as stated) fails for disconnected graphs.*

## Corollary 6.15

*If  $G$  is a planar graph whose order  $v$  is at least 3, whose size is  $e$ , and whose girth  $g$  is at least 3 but finite, then*

$$e \leq \frac{(v-2)g}{g-2}.$$

*If  $G$  is simple, then  $e \leq 3v - 6$ .*

$$e \leq 3v - 6.$$

### Example 6.16

Is  $K_5$  planar?

No, since  $e = 10 > 3v - 6 = 9$ .

$$e \leq 3v - 6.$$

### Example 6.16

Is  $K_5$  planar?

No, since  $e = 10 > 3v - 6 = 9$ .

Is  $K_{3,3}$  planar?



$$e \leq 3v - 6.$$

### Example 6.16

Is  $K_5$  planar?

No, since  $e = 10 > 3v - 6 = 9$ .

Is  $K_{3,3}$  planar?

No, since

$$e = 9 > \frac{(v-2)g}{g-2} = 8.$$

# Statement of the Kuratowski Theorem

## Theorem 6.17 (Kuratowski 1930)

*A graph is planar if and only if it has neither  $K_5$  nor  $K_{3,3}$  as a topological minor.*

# Statement of the Kuratowski Theorem

## Theorem 6.17 (Kuratowski 1930)

*A graph is planar if and only if it has neither  $K_5$  nor  $K_{3,3}$  as a topological minor.*

## Theorem 6.18 (Wagner)

*A graph is planar if and only if it has neither  $K_5$  nor  $K_{3,3}$  as a minor.*

# Statement of the Kuratowski Theorem

## Theorem 6.17 (Kuratowski 1930)

*A graph is planar if and only if it has neither  $K_5$  nor  $K_{3,3}$  as a topological minor.*

## Theorem 6.18 (Wagner)

*A graph is planar if and only if it has neither  $K_5$  nor  $K_{3,3}$  as a minor.*

## Lemma 6.19

*If  $F$  is the edge-set of the boundary of a face of a plane graph  $G$ , then  $G$  has an plane embedding in which  $F$  is the boundary of the infinite face.*

# Minimally No-Planar Graphs

## Definition 6.20

A graph is **minimally non-planar** if it is non-planar, but every proper subgraph of it is planar.

# Minimally No-Planar Graphs

## Definition 6.20

A graph is **minimally non-planar** if it is non-planar, but every proper subgraph of it is planar.

## Lemma 6.21

*Every minimal non-planar graph is 2-connected.*

# Minimally No-Planar Graphs

## Definition 6.20

A graph is **minimally non-planar** if it is non-planar, but every proper subgraph of it is planar.

## Lemma 6.21

*Every minimal non-planar graph is 2-connected.*

## Proof.

If  $G$  is disconnected, we can embed one component of  $G$  inside one face of the rest of  $G$ .



# Minimally Non-Planar Graphs

## Definition 6.20

A graph is **minimally non-planar** if it is non-planar, but every proper subgraph of it is planar.

## Lemma 6.21

*Every minimal non-planar graph is 2-connected.*

## Proof.

If  $G$  is disconnected, we can embed one component of  $G$  inside one face of the rest of  $G$ . Similarly, if  $G$  has a cut-vertex  $v$ , let  $G_1, G_2, \dots, G_k$  be the subgraphs of  $G$  induced by  $v$  together with the components of  $G - v$ .





# Minimally Non-Planar Graphs

## Definition 6.20

A graph is **minimally non-planar** if it is non-planar, but every proper subgraph of it is planar.

## Lemma 6.21

*Every minimal non-planar graph is 2-connected.*

## Proof.

If  $G$  is disconnected, we can embed one component of  $G$  inside one face of the rest of  $G$ . Similarly, if  $G$  has a cut-vertex  $v$ , let  $G_1, G_2, \dots, G_k$  be the subgraphs of  $G$  induced by  $v$  together with the components of  $G - v$ . By the minimality of  $G$ , these subgraphs are planar.



# Minimally Non-Planar Graphs

## Definition 6.20

A graph is **minimally non-planar** if it is non-planar, but every proper subgraph of it is planar.

## Lemma 6.21

*Every minimal non-planar graph is 2-connected.*

## Proof.

If  $G$  is disconnected, we can embed one component of  $G$  inside one face of the rest of  $G$ . Similarly, if  $G$  has a cut-vertex  $v$ , let  $G_1, G_2, \dots, G_k$  be the subgraphs of  $G$  induced by  $v$  together with the components of  $G - v$ . By the minimality of  $G$ , these subgraphs are planar. It is easy to see that the plane embeddings of these subgraphs can be put together to form a plane embedding of  $G$ . □

## Lemma 6.22

*Suppose  $G = H_1 \oplus_2 H_2$  is non-planar.*

## Lemma 6.22

*Suppose  $G = H_1 \oplus_2 H_2$  is non-planar. Then at least one of  $H_1$  and  $H_2$  is non-planar.*

# Kuratowski Graphs and Topological Minors

## Lemma 6.22

*Suppose  $G = H_1 \oplus_2 H_2$  is non-planar. Then at least one of  $H_1$  and  $H_2$  is non-planar.*

## Proof.

Let  $e$  be the common edge of  $H_1$  and  $H_2$ .



## Lemma 6.22

*Suppose  $G = H_1 \oplus_2 H_2$  is non-planar. Then at least one of  $H_1$  and  $H_2$  is non-planar.*

## Proof.

Let  $e$  be the common edge of  $H_1$  and  $H_2$ . Suppose both  $H_1$  and  $H_2$  are planar.



## Lemma 6.22

*Suppose  $G = H_1 \oplus_2 H_2$  is non-planar. Then at least one of  $H_1$  and  $H_2$  is non-planar.*

## Proof.

Let  $e$  be the common edge of  $H_1$  and  $H_2$ . Suppose both  $H_1$  and  $H_2$  are planar. By Lemma 6.19, each of  $H_1$  and  $H_2$  can be embedded in the plane with  $e$  in the boundary of the infinite face.



## Lemma 6.22

*Suppose  $G = H_1 \oplus_2 H_2$  is non-planar. Then at least one of  $H_1$  and  $H_2$  is non-planar.*

## Proof.

Let  $e$  be the common edge of  $H_1$  and  $H_2$ . Suppose both  $H_1$  and  $H_2$  are planar. By Lemma 6.19, each of  $H_1$  and  $H_2$  can be embedded in the plane with  $e$  in the boundary of the infinite face. It is now easy to put together the embeddings of  $H_1$  and  $H_2$  into a plane embedding of  $G$ .  $\square$



# Kuratowski Graphs and Topological Minors

## Lemma 6.22

*Suppose  $G = H_1 \oplus_2 H_2$  is non-planar. Then at least one of  $H_1$  and  $H_2$  is non-planar.*

## Proof.

Let  $e$  be the common edge of  $H_1$  and  $H_2$ . Suppose both  $H_1$  and  $H_2$  are planar. By Lemma 6.19, each of  $H_1$  and  $H_2$  can be embedded in the plane with  $e$  in the boundary of the infinite face. It is now easy to put together the embeddings of  $H_1$  and  $H_2$  into a plane embedding of  $G$ .  $\square$

## Definition 6.23

- ▶ A **Kuratowski subgraph** is a subgraph isomorphic to a subdivision of  $K_5$  or of  $K_{3,3}$ .

# Kuratowski Graphs and Topological Minors

## Lemma 6.22

*Suppose  $G = H_1 \oplus_2 H_2$  is non-planar. Then at least one of  $H_1$  and  $H_2$  is non-planar.*

## Proof.

Let  $e$  be the common edge of  $H_1$  and  $H_2$ . Suppose both  $H_1$  and  $H_2$  are planar. By Lemma 6.19, each of  $H_1$  and  $H_2$  can be embedded in the plane with  $e$  in the boundary of the infinite face. It is now easy to put together the embeddings of  $H_1$  and  $H_2$  into a plane embedding of  $G$ .  $\square$

## Definition 6.23

- ▶ A **Kuratowski subgraph** is a subgraph isomorphic to a subdivision of  $K_5$  or of  $K_{3,3}$ .
- ▶ A vertex of a graph  $G$  is a **branch vertex** of a Kuratowski subgraph  $H$  of  $G$ , if its degree in  $H$  exceeds two.

## Lemma 6.24

*If  $G/e$  has a Kuratowski subgraph, then so does  $G$ .*

# Tutte's Version of Kuratowski's Theorem

## Definition 6.25

A plane embedding is **convex** if every face except the infinite one is a convex polygon.

# Tutte's Version of Kuratowski's Theorem

## Definition 6.25

A plane embedding is **convex** if every face except the infinite one is a convex polygon.

## Theorem 6.26 (Tutte 1960–63)

*If  $G$  is a simple 3-connected graph with neither  $K_5$  nor  $K_{3,3}$  as the topological minor, then  $G$  has a convex embedding in the plane with no three vertices on a line.*