

Proof of Hall's Theorem

Recall the **Hall's Condition**: $|N(S)| \geq |S|$ for every $S \subseteq X$.

Proof of Hall's Theorem

Recall the **Hall's Condition**: $|N(S)| \geq |S|$ for every $S \subseteq X$.
Necessity is clear.

Proof of Hall's Theorem

Recall the **Hall's Condition**: $|N(S)| \geq |S|$ for every $S \subseteq X$.

Necessity is clear.

To prove sufficiency, suppose the Hall's condition holds, let M be a maximum matching, and suppose $u \in X$ is unsaturated.

Proof of Hall's Theorem

Recall the **Hall's Condition**: $|N(S)| \geq |S|$ for every $S \subseteq X$.

Necessity is clear.

To prove sufficiency, suppose the Hall's condition holds, let M be a maximum matching, and suppose $u \in X$ is unsaturated. Let S and T be subsets of X and Y , respectively, that are reachable from u by M -alternating paths.

Proof of Hall's Theorem

Recall the **Hall's Condition**: $|N(S)| \geq |S|$ for every $S \subseteq X$.

Necessity is clear.

To prove sufficiency, suppose the Hall's condition holds, let M be a maximum matching, and suppose $u \in X$ is unsaturated. Let S and T be subsets of X and Y , respectively, that are reachable from u by M -alternating paths. These paths reach Y from u along edges not in M , and reach X along edges in M .

Proof of Hall's Theorem

Recall the **Hall's Condition**: $|N(S)| \geq |S|$ for every $S \subseteq X$.

Necessity is clear.

To prove sufficiency, suppose the Hall's condition holds, let M be a maximum matching, and suppose $u \in X$ is unsaturated. Let S and T be subsets of X and Y , respectively, that are reachable from u by M -alternating paths. These paths reach Y from u along edges not in M , and reach X along edges in M . Hence every vertex in $S - u$ is reached along an edge in M from a vertex in T .

Proof of Hall's Theorem

Recall the **Hall's Condition**: $|N(S)| \geq |S|$ for every $S \subseteq X$.

Necessity is clear.

To prove sufficiency, suppose the Hall's condition holds, let M be a maximum matching, and suppose $u \in X$ is unsaturated. Let S and T be subsets of X and Y , respectively, that are reachable from u by M -alternating paths. These paths reach Y from u along edges not in M , and reach X along edges in M . Hence every vertex in $S - u$ is reached along an edge in M from a vertex in T . Since there are no augmenting paths, every vertex in T is saturated.

Proof of Hall's Theorem

Recall the **Hall's Condition**: $|N(S)| \geq |S|$ for every $S \subseteq X$.

Necessity is clear.

To prove sufficiency, suppose the Hall's condition holds, let M be a maximum matching, and suppose $u \in X$ is unsaturated. Let S and T be subsets of X and Y , respectively, that are reachable from u by M -alternating paths. These paths reach Y from u along edges not in M , and reach X along edges in M . Hence every vertex in $S - u$ is reached along an edge in M from a vertex in T . Since there are no augmenting paths, every vertex in T is saturated. Hence the edges of M establish a bijection between T and $S - u$.

Proof of Hall's Theorem

Recall the **Hall's Condition**: $|N(S)| \geq |S|$ for every $S \subseteq X$.

Necessity is clear.

To prove sufficiency, suppose the Hall's condition holds, let M be a maximum matching, and suppose $u \in X$ is unsaturated. Let S and T be subsets of X and Y , respectively, that are reachable from u by M -alternating paths. These paths reach Y from u along edges not in M , and reach X along edges in M . Hence every vertex in $S - u$ is reached along an edge in M from a vertex in T . Since there are no augmenting paths, every vertex in T is saturated. Hence the edges of M establish a bijection between T and $S - u$. Note that an edge between S and $y \in Y - T$ would be an edge not in M , and thus create an M -augmenting path to y , which contradicts $y \notin T$.

Proof of Hall's Theorem

Recall the **Hall's Condition**: $|N(S)| \geq |S|$ for every $S \subseteq X$.

Necessity is clear.

To prove sufficiency, suppose the Hall's condition holds, let M be a maximum matching, and suppose $u \in X$ is unsaturated. Let S and T be subsets of X and Y , respectively, that are reachable from u by M -alternating paths. These paths reach Y from u along edges not in M , and reach X along edges in M . Hence every vertex in $S - u$ is reached along an edge in M from a vertex in T . Since there are no augmenting paths, every vertex in T is saturated. Hence the edges of M establish a bijection between T and $S - u$. Note that an edge between S and $y \in Y - T$ would be an edge not in M , and thus create an M -augmenting path to y , which contradicts $y \notin T$. Hence $T = N(S)$,

Proof of Hall's Theorem

Recall the **Hall's Condition**: $|N(S)| \geq |S|$ for every $S \subseteq X$.

Necessity is clear.

To prove sufficiency, suppose the Hall's condition holds, let M be a maximum matching, and suppose $u \in X$ is unsaturated. Let S and T be subsets of X and Y , respectively, that are reachable from u by M -alternating paths. These paths reach Y from u along edges not in M , and reach X along edges in M . Hence every vertex in $S - u$ is reached along an edge in M from a vertex in T . Since there are no augmenting paths, every vertex in T is saturated. Hence the edges of M establish a bijection between T and $S - u$. Note that an edge between S and $y \in Y - T$ would be an edge not in M , and thus create an M -augmenting path to y , which contradicts $y \notin T$. Hence $T = N(S)$, and $|N(S)| = |T| = |S| - 1 < |S|$; a contradiction.