

Definition 5.1

- ▶ A **separating set** or a **vertex cut** of a graph G is a set $S \subseteq V(G)$ such that $G - S$ has more than one component.

Definition 5.1

- ▶ A **separating set** or a **vertex cut** of a graph G is a set $S \subseteq V(G)$ such that $G - S$ has more than one component.
- ▶ **Vertex connectivity** or **connectivity** $\kappa(G)$ of a graph G is defined as follows:

Definition 5.1

- ▶ A **separating set** or a **vertex cut** of a graph G is a set $S \subseteq V(G)$ such that $G - S$ has more than one component.
- ▶ **Vertex connectivity** or **connectivity** $\kappa(G)$ of a graph G is defined as follows:
 - ▶ $\kappa(G) = 0$ if G is disconnected;

Definition 5.1

- ▶ A **separating set** or a **vertex cut** of a graph G is a set $S \subseteq V(G)$ such that $G - S$ has more than one component.
- ▶ **Vertex connectivity** or **connectivity** $\kappa(G)$ of a graph G is defined as follows:
 - ▶ $\kappa(G) = 0$ if G is disconnected;
 - ▶ $\kappa(G) = |G| - 1$ if G is connected, but has no pair of distinct non-adjacent vertices.

Definition 5.1

- ▶ A **separating set** or a **vertex cut** of a graph G is a set $S \subseteq V(G)$ such that $G - S$ has more than one component.
- ▶ **Vertex connectivity** or **connectivity** $\kappa(G)$ of a graph G is defined as follows:
 - ▶ $\kappa(G) = 0$ if G is disconnected;
 - ▶ $\kappa(G) = |G| - 1$ if G is connected, but has no pair of distinct non-adjacent vertices.
 - ▶ $\kappa(G) = j$ if G is connected, but has a pair of non-adjacent vertices, and j is the smallest integer such that G has a j -element vertex cut.

Definition 5.1

- ▶ A **separating set** or a **vertex cut** of a graph G is a set $S \subseteq V(G)$ such that $G - S$ has more than one component.
- ▶ **Vertex connectivity** or **connectivity** $\kappa(G)$ of a graph G is defined as follows:
 - ▶ $\kappa(G) = 0$ if G is disconnected;
 - ▶ $\kappa(G) = |G| - 1$ if G is connected, but has no pair of distinct non-adjacent vertices.
 - ▶ $\kappa(G) = j$ if G is connected, but has a pair of non-adjacent vertices, and j is the smallest integer such that G has a j -element vertex cut.
- ▶ If k is a positive integer, then G is **k -connected** or **k -vertex-connected** if $k \leq \kappa(G)$.

Vertex Connectivity

Definition 5.1

- ▶ A **separating set** or a **vertex cut** of a graph G is a set $S \subseteq V(G)$ such that $G - S$ has more than one component.
- ▶ **Vertex connectivity** or **connectivity** $\kappa(G)$ of a graph G is defined as follows:
 - ▶ $\kappa(G) = 0$ if G is disconnected;
 - ▶ $\kappa(G) = |G| - 1$ if G is connected, but has no pair of distinct non-adjacent vertices.
 - ▶ $\kappa(G) = j$ if G is connected, but has a pair of non-adjacent vertices, and j is the smallest integer such that G has a j -element vertex cut.
- ▶ If k is a positive integer, then G is **k -connected** or **k -vertex-connected** if $k \leq \kappa(G)$.

Note 5.2

- ▶ *Vertex connectivity is not affected by adding or deleting loops and parallel edges.*

Vertex Connectivity

Definition 5.1

- ▶ A **separating set** or a **vertex cut** of a graph G is a set $S \subseteq V(G)$ such that $G - S$ has more than one component.
- ▶ **Vertex connectivity** or **connectivity** $\kappa(G)$ of a graph G is defined as follows:
 - ▶ $\kappa(G) = 0$ if G is disconnected;
 - ▶ $\kappa(G) = |G| - 1$ if G is connected, but has no pair of distinct non-adjacent vertices.
 - ▶ $\kappa(G) = j$ if G is connected, but has a pair of non-adjacent vertices, and j is the smallest integer such that G has a j -element vertex cut.
- ▶ If k is a positive integer, then G is **k -connected** or **k -vertex-connected** if $k \leq \kappa(G)$.

Note 5.2

- ▶ *Vertex connectivity is not affected by adding or deleting loops and parallel edges.*
- ▶ *K_1 is connected although $\kappa(K_1) = 0$.*

Example 5.3

- ▶ $\kappa(K_n) = n - 1$ for $n \geq 2$;

Example 5.3

- ▶ $\kappa(K_n) = n - 1$ for $n \geq 2$;
- ▶ $\kappa(K_{m,n}) = \min(m, n)$;

Example 5.3

- ▶ $\kappa(K_n) = n - 1$ for $n \geq 2$;
- ▶ $\kappa(K_{m,n}) = \min(m, n)$;
- ▶ If T is a non-trivial tree, then $\kappa(T) = 1$.

Example 5.3

- ▶ $\kappa(K_n) = n - 1$ for $n \geq 2$;
- ▶ $\kappa(K_{m,n}) = \min(m, n)$;
- ▶ If T is a non-trivial tree, then $\kappa(T) = 1$.
- ▶ $\kappa(C_n) = 2$ for all $n \geq 3$.

Example 5.3

- ▶ $\kappa(K_n) = n - 1$ for $n \geq 2$;
- ▶ $\kappa(K_{m,n}) = \min(m, n)$;
- ▶ If T is a non-trivial tree, then $\kappa(T) = 1$.
- ▶ $\kappa(C_n) = 2$ for all $n \geq 3$.
- ▶ An **n -wheel** W_n is obtained from C_n by adding a new vertex and joining it to all vertices of C_n . If $n \geq 3$, then $\kappa(W_n) = 3$.

Edge Connectivity

Definition 5.4

- ▶ A **disconnecting set of edges** of a graph G with $|G| > 1$ is a set $F \subseteq E(G)$ such that $G \setminus F$ has more than one component.

Edge Connectivity

Definition 5.4

- ▶ A **disconnecting set of edges** of a graph G with $|G| > 1$ is a set $F \subseteq E(G)$ such that $G \setminus F$ has more than one component.
- ▶ A graph is **k -edge-connected** if every disconnecting set has at least k edges.

Edge Connectivity

Definition 5.4

- ▶ A **disconnecting set of edges** of a graph G with $|G| > 1$ is a set $F \subseteq E(G)$ such that $G \setminus F$ has more than one component.
- ▶ A graph is **k -edge-connected** if every disconnecting set has at least k edges.
- ▶ The **edge connectivity** of G , written $\kappa'(G)$ is the maximum k such that G is k -edge-connected.

Definition 5.4

- ▶ A **disconnecting set of edges** of a graph G with $|G| > 1$ is a set $F \subseteq E(G)$ such that $G \setminus F$ has more than one component.
- ▶ A graph is **k -edge-connected** if every disconnecting set has at least k edges.
- ▶ The **edge connectivity** of G , written $\kappa'(G)$ is the maximum k such that G is k -edge-connected.
- ▶ Given $S, T \subseteq V(G)$, we write $[S, T]$ for the set of edges with one endpoint in S and the other in T .

Edge Connectivity

Definition 5.4

- ▶ A **disconnecting set of edges** of a graph G with $|G| > 1$ is a set $F \subseteq E(G)$ such that $G \setminus F$ has more than one component.
- ▶ A graph is **k -edge-connected** if every disconnecting set has at least k edges.
- ▶ The **edge connectivity** of G , written $\kappa'(G)$ is the maximum k such that G is k -edge-connected.
- ▶ Given $S, T \subseteq V(G)$, we write $[S, T]$ for the set of edges with one endpoint in S and the other in T .
- ▶ An **edge cut** is a set of edges of the form $[S, \overline{S}]$ where S is a non-empty proper subset of $V(G)$.

Edge Connectivity

Definition 5.4


- ▶ A **disconnecting set of edges** of a graph G with $|G| > 1$ is a set $F \subseteq E(G)$ such that $G \setminus F$ has more than one component.
- ▶ A graph is **k -edge-connected** if every disconnecting set has at least k edges.
- ▶ The **edge connectivity** of G , written $\kappa'(G)$ is the maximum k such that G is k -edge-connected.
- ▶ Given $S, T \subseteq V(G)$, we write $[S, T]$ for the set of edges with one endpoint in S and the other in T .
- ▶ An **edge cut** is a set of edges of the form $[S, \overline{S}]$ where S is a non-empty proper subset of $V(G)$.
- ▶ A **bond** is a minimal non-empty edge cut.

Edge Connectivity

Definition 5.4

- ▶ A **disconnecting set of edges** of a graph G with $|G| > 1$ is a set $F \subseteq E(G)$ such that $G \setminus F$ has more than one component.
- ▶ A graph is **k -edge-connected** if every disconnecting set has at least k edges.
- ▶ The **edge connectivity** of G , written $\kappa'(G)$ is the maximum k such that G is k -edge-connected.
- ▶ Given $S, T \subseteq V(G)$, we write $[S, T]$ for the set of edges with one endpoint in S and the other in T .
- ▶ An **edge cut** is a set of edges of the form $[S, \overline{S}]$ where S is a non-empty proper subset of $V(G)$.
- ▶ A **bond** is a minimal non-empty edge cut.

Example 5.5




Edge Connectivity

Definition 5.4

- ▶ A **disconnecting set of edges** of a graph G with $|G| > 1$ is a set $F \subseteq E(G)$ such that $G \setminus F$ has more than one component.
- ▶ A graph is **k -edge-connected** if every disconnecting set has at least k edges.
- ▶ The **edge connectivity** of G , written $\kappa'(G)$ is the maximum k such that G is k -edge-connected.
- ▶ Given $S, T \subseteq V(G)$, we write $[S, T]$ for the set of edges with one endpoint in S and the other in T .
- ▶ An **edge cut** is a set of edges of the form $[S, \overline{S}]$ where S is a non-empty proper subset of $V(G)$.
- ▶ A **bond** is a minimal non-empty edge cut.

Example 5.5

disconnecting set, but
not an edge cut




Edge Connectivity

Definition 5.4

- ▶ A **disconnecting set of edges** of a graph G with $|G| > 1$ is a set $F \subseteq E(G)$ such that $G \setminus F$ has more than one component.
- ▶ A graph is **k -edge-connected** if every disconnecting set has at least k edges.
- ▶ The **edge connectivity** of G , written $\kappa'(G)$ is the maximum k such that G is k -edge-connected.
- ▶ Given $S, T \subseteq V(G)$, we write $[S, T]$ for the set of edges with one endpoint in S and the other in T .
- ▶ An **edge cut** is a set of edges of the form $[S, \overline{S}]$ where S is a non-empty proper subset of $V(G)$.
- ▶ A **bond** is a minimal non-empty edge cut.

Example 5.5

disconnecting set, but
not an edge cut




Edge Connectivity

Definition 5.4

- ▶ A **disconnecting set of edges** of a graph G with $|G| > 1$ is a set $F \subseteq E(G)$ such that $G \setminus F$ has more than one component.
- ▶ A graph is **k -edge-connected** if every disconnecting set has at least k edges.
- ▶ The **edge connectivity** of G , written $\kappa'(G)$ is the maximum k such that G is k -edge-connected.
- ▶ Given $S, T \subseteq V(G)$, we write $[S, T]$ for the set of edges with one endpoint in S and the other in T .
- ▶ An **edge cut** is a set of edges of the form $[S, \overline{S}]$ where S is a non-empty proper subset of $V(G)$.
- ▶ A **bond** is a minimal non-empty edge cut.

Example 5.5

edge cut, but not a bond




Edge Connectivity

Definition 5.4

- ▶ A **disconnecting set of edges** of a graph G with $|G| > 1$ is a set $F \subseteq E(G)$ such that $G \setminus F$ has more than one component.
- ▶ A graph is **k -edge-connected** if every disconnecting set has at least k edges.
- ▶ The **edge connectivity** of G , written $\kappa'(G)$ is the maximum k such that G is k -edge-connected.
- ▶ Given $S, T \subseteq V(G)$, we write $[S, T]$ for the set of edges with one endpoint in S and the other in T .
- ▶ An **edge cut** is a set of edges of the form $[S, \overline{S}]$ where S is a non-empty proper subset of $V(G)$.
- ▶ A **bond** is a minimal non-empty edge cut.

Example 5.5

edge cut, but not a bond




Edge Connectivity

Definition 5.4

- ▶ A **disconnecting set of edges** of a graph G with $|G| > 1$ is a set $F \subseteq E(G)$ such that $G \setminus F$ has more than one component.
- ▶ A graph is **k -edge-connected** if every disconnecting set has at least k edges.
- ▶ The **edge connectivity** of G , written $\kappa'(G)$ is the maximum k such that G is k -edge-connected.
- ▶ Given $S, T \subseteq V(G)$, we write $[S, T]$ for the set of edges with one endpoint in S and the other in T .
- ▶ An **edge cut** is a set of edges of the form $[S, \overline{S}]$ where S is a non-empty proper subset of $V(G)$.
- ▶ A **bond** is a minimal non-empty edge cut.

Example 5.5

bond




Edge Connectivity

Definition 5.4

- ▶ A **disconnecting set of edges** of a graph G with $|G| > 1$ is a set $F \subseteq E(G)$ such that $G \setminus F$ has more than one component.
- ▶ A graph is **k -edge-connected** if every disconnecting set has at least k edges.
- ▶ The **edge connectivity** of G , written $\kappa'(G)$ is the maximum k such that G is k -edge-connected.
- ▶ Given $S, T \subseteq V(G)$, we write $[S, T]$ for the set of edges with one endpoint in S and the other in T .
- ▶ An **edge cut** is a set of edges of the form $[S, \overline{S}]$ where S is a non-empty proper subset of $V(G)$.
- ▶ A **bond** is a minimal non-empty edge cut.

Example 5.5

bond



Whitney's Theorem

Note 5.6

The edge connectivity of a graph is unaffected by adding or deleting loops, but is affected by adding and deleting edges in parallel.

Whitney's Theorem

Note 5.6

The edge connectivity of a graph is unaffected by adding or deleting loops, but is affected by adding and deleting edges in parallel.

Theorem 5.7 (Whitney 1932)

If G is graph with $|G| > 1$, then $\kappa(G) \leq \kappa'(G) \leq \delta(G)$.

Whitney's Theorem

Note 5.6

The edge connectivity of a graph is unaffected by adding or deleting loops, but is affected by adding and deleting edges in parallel.

Theorem 5.7 (Whitney 1932)

If G is graph with $|G| > 1$, then $\kappa(G) \leq \kappa'(G) \leq \delta(G)$.

Proof.

The edges incident to a vertex form a disconnecting set, so $\kappa' \leq \delta$.



Whitney's Theorem

Note 5.6

The edge connectivity of a graph is unaffected by adding or deleting loops, but is affected by adding and deleting edges in parallel.

Theorem 5.7 (Whitney 1932)

If G is graph with $|G| > 1$, then $\kappa(G) \leq \kappa'(G) \leq \delta(G)$.

Proof.

The edges incident to a vertex form a disconnecting set, so $\kappa' \leq \delta$.
Clearly, $\kappa(G) \leq |G| - 1$.



Whitney's Theorem

Note 5.6

The edge connectivity of a graph is unaffected by adding or deleting loops, but is affected by adding and deleting edges in parallel.

Theorem 5.7 (Whitney 1932)

If G is graph with $|G| > 1$, then $\kappa(G) \leq \kappa'(G) \leq \delta(G)$.

Proof.

The edges incident to a vertex form a disconnecting set, so $\kappa' \leq \delta$.
Clearly, $\kappa(G) \leq |G| - 1$. Suppose $[S, \overline{S}]$ is a minimum edge cut of size $k' = \kappa'(G)$.



Whitney's Theorem

Note 5.6

The edge connectivity of a graph is unaffected by adding or deleting loops, but is affected by adding and deleting edges in parallel.

Theorem 5.7 (Whitney 1932)

If G is graph with $|G| > 1$, then $\kappa(G) \leq \kappa'(G) \leq \delta(G)$.

Proof.

The edges incident to a vertex form a disconnecting set, so $\kappa' \leq \delta$. Clearly, $\kappa(G) \leq |G| - 1$. Suppose $[S, \overline{S}]$ is a minimum edge cut of size $k' = \kappa'(G)$. If every vertex in S is adjacent to every vertex in \overline{S} , then $k' = |S||\overline{S}| \geq |G| - 1$, and the inequality follows.



Whitney's Theorem

Note 5.6

The edge connectivity of a graph is unaffected by adding or deleting loops, but is affected by adding and deleting edges in parallel.

Theorem 5.7 (Whitney 1932)

If G is graph with $|G| > 1$, then $\kappa(G) \leq \kappa'(G) \leq \delta(G)$.

Proof.

The edges incident to a vertex form a disconnecting set, so $\kappa' \leq \delta$. Clearly, $\kappa(G) \leq |G| - 1$. Suppose $[S, \overline{S}]$ is a minimum edge cut of size $k' = \kappa'(G)$. If every vertex in S is adjacent to every vertex in \overline{S} , then $k' = |S||\overline{S}| \geq |G| - 1$, and the inequality follows. Hence we may assume that there are vertices $s \in S$ and $\overline{s} \in \overline{S}$ that are non-adjacent.



Whitney's Theorem

Note 5.6

The edge connectivity of a graph is unaffected by adding or deleting loops, but is affected by adding and deleting edges in parallel.

Theorem 5.7 (Whitney 1932)

If G is graph with $|G| > 1$, then $\kappa(G) \leq \kappa'(G) \leq \delta(G)$.

Proof.

The edges incident to a vertex form a disconnecting set, so $\kappa' \leq \delta$. Clearly, $\kappa(G) \leq |G| - 1$. Suppose $[S, \overline{S}]$ is a minimum edge cut of size $k' = \kappa'(G)$. If every vertex in S is adjacent to every vertex in \overline{S} , then $k' = |S||\overline{S}| \geq |G| - 1$, and the inequality follows. Hence we may assume that there are vertices $s \in S$ and $\overline{s} \in \overline{S}$ that are non-adjacent. Let T be the vertex set consisting of all neighbors of s in \overline{S} and all vertices in $S - s$ that have neighbors in \overline{S} .



Whitney's Theorem

Note 5.6

The edge connectivity of a graph is unaffected by adding or deleting loops, but is affected by adding and deleting edges in parallel.

Theorem 5.7 (Whitney 1932)

If G is graph with $|G| > 1$, then $\kappa(G) \leq \kappa'(G) \leq \delta(G)$.

Proof.

The edges incident to a vertex form a disconnecting set, so $\kappa' \leq \delta$. Clearly, $\kappa(G) \leq |G| - 1$. Suppose $[S, \overline{S}]$ is a minimum edge cut of size $k' = \kappa'(G)$. If every vertex in S is adjacent to every vertex in \overline{S} , then $k' = |S||\overline{S}| \geq |G| - 1$, and the inequality follows. Hence we may assume that there are vertices $s \in S$ and $\overline{s} \in \overline{S}$ that are non-adjacent. Let T be the vertex set consisting of all neighbors of s in \overline{S} and all vertices in $S - s$ that have neighbors in \overline{S} . Then T is a vertex cut consisting of one endpoint of each edge in $[S, \overline{S}]$. □

Whitney's Theorem

Note 5.6

The edge connectivity of a graph is unaffected by adding or deleting loops, but is affected by adding and deleting edges in parallel.

Theorem 5.7 (Whitney 1932)


If G is graph with $|G| > 1$, then $\kappa(G) \leq \kappa'(G) \leq \delta(G)$.

Proof.

The edges incident to a vertex form a disconnecting set, so $\kappa' \leq \delta$. Clearly, $\kappa(G) \leq |G| - 1$. Suppose $[S, \overline{S}]$ is a minimum edge cut of size $k' = \kappa'(G)$. If every vertex in S is adjacent to every vertex in \overline{S} , then $k' = |S||\overline{S}| \geq |G| - 1$, and the inequality follows. Hence we may assume that there are vertices $s \in S$ and $\overline{s} \in \overline{S}$ that are non-adjacent. Let T be the vertex set consisting of all neighbors of s in \overline{S} and all vertices in $S - s$ that have neighbors in \overline{S} . Then T is a vertex cut consisting of one endpoint of each edge in $[S, \overline{S}]$. Hence $\kappa \leq \kappa'$. □


Connectivity Example

Example 5.8



Connectivity Example


Example 5.8



$$\kappa = 1 < \kappa' = 2 < \delta = 3$$

Connectivity Example

Example 5.8




$$\kappa = 1 < \kappa' = 2 < \delta = 3$$

Theorem 5.9

If G is a connected graph and S is a non-empty proper subset of $V(G)$, then $F = [S, \overline{S}]$ is a bond if and only if $G \setminus F$ has two components.

Connectivity Example

Example 5.8




$$\kappa = 1 < \kappa' = 2 < \delta = 3$$

Theorem 5.9

If G is a connected graph and S is a non-empty proper subset of $V(G)$, then $F = [S, \overline{S}]$ is a bond if and only if $G \setminus F$ has two components. Equivalently, if and only if the subgraphs of G induced by each of S and \overline{S} are connected.

Connectivity Example

Example 5.8



$$\kappa = 1 < \kappa' = 2 < \delta = 3$$

Theorem 5.9

If G is a connected graph and S is a non-empty proper subset of $V(G)$, then $F = [S, \overline{S}]$ is a bond if and only if $G \setminus F$ has two components. Equivalently, if and only if the subgraphs of G induced by each of S and \overline{S} are connected.


Proof.

If $G \setminus F$ has two components, then F is a bond, since $G \setminus F'$ is connected for every proper subset F' of F .



Connectivity Example

Example 5.8



$$\kappa = 1 < \kappa' = 2 < \delta = 3$$

Theorem 5.9

If G is a connected graph and S is a non-empty proper subset of $V(G)$, then $F = [S, \overline{S}]$ is a bond if and only if $G \setminus F$ has two components. Equivalently, if and only if the subgraphs of G induced by each of S and \overline{S} are connected.

Proof.


If $G \setminus F$ has two components, then F is a bond, since $G \setminus F'$ is connected for every proper subset F' of F .

If $G \setminus F$ has more than two components, then we may assume $S = A \cup B$ with no edges between A and B .



Connectivity Example

Example 5.8



$$\kappa = 1 < \kappa' = 2 < \delta = 3$$

Theorem 5.9

If G is a connected graph and S is a non-empty proper subset of $V(G)$, then $F = [S, \overline{S}]$ is a bond if and only if $G \setminus F$ has two components. Equivalently, if and only if the subgraphs of G induced by each of S and \overline{S} are connected.

Proof.

If $G \setminus F$ has two components, then F is a bond, since $G \setminus F'$ is connected for every proper subset F' of F .

If $G \setminus F$ has more than two components, then we may assume $S = A \cup B$ with no edges between A and B . Then $[A, \overline{A}]$ is an edge cut which is a proper subset of F ; a contradiction. \square

Definition 5.10

- ▶ A **k -separation** of a graph G is a pair of subgraphs $\{A, B\}$ of G such that each of A and B has size at least k , $A \neq G$, $B \neq G$, $A \cup B = G$, and $A \cap B$ is trivial of order at most k .

Definition 5.10

- ▶ A **k -separation** of a graph G is a pair of subgraphs $\{A, B\}$ of G such that each of A and B has size at least k , $A \neq G$, $B \neq G$, $A \cup B = G$, and $A \cap B$ is trivial of order at most k .
- ▶ If G has a k -separation for some k , then **Tutte connectivity** of G is $\min\{j : G \text{ has a } j \text{ separation}\}$, and ∞ if no k -separation exists.


Tutte Connectivity

Definition 5.10

- ▶ A **k -separation** of a graph G is a pair of subgraphs $\{A, B\}$ of G such that each of A and B has size at least k , $A \neq G$, $B \neq G$, $A \cup B = G$, and $A \cap B$ is trivial of order at most k .
- ▶ If G has a k -separation for some k , then **Tutte connectivity** of G is $\min\{j : G \text{ has a } j \text{ separation}\}$, and ∞ if no k -separation exists.

Example 5.11

1-separations:



Tutte Connectivity

Definition 5.10

- ▶ A **k -separation** of a graph G is a pair of subgraphs $\{A, B\}$ of G such that each of A and B has size at least k , $A \neq G$, $B \neq G$, $A \cup B = G$, and $A \cap B$ is trivial of order at most k .
- ▶ If G has a k -separation for some k , then **Tutte connectivity** of G is $\min\{j : G \text{ has a } j \text{ separation}\}$, and ∞ if no k -separation exists.

Example 5.11

1-separations:




Tutte Connectivity

Definition 5.10


- ▶ A **k -separation** of a graph G is a pair of subgraphs $\{A, B\}$ of G such that each of A and B has size at least k , $A \neq G$, $B \neq G$, $A \cup B = G$, and $A \cap B$ is trivial of order at most k .
- ▶ If G has a k -separation for some k , then **Tutte connectivity** of G is $\min\{j : G \text{ has a } j \text{ separation}\}$, and ∞ if no k -separation exists.

Example 5.11

1-separations:



2-separations:




Tutte Connectivity

Definition 5.10


- ▶ A **k -separation** of a graph G is a pair of subgraphs $\{A, B\}$ of G such that each of A and B has size at least k , $A \neq G$, $B \neq G$, $A \cup B = G$, and $A \cap B$ is trivial of order at most k .
- ▶ If G has a k -separation for some k , then **Tutte connectivity** of G is $\min\{j : G \text{ has a } j \text{ separation}\}$, and ∞ if no k -separation exists.

Example 5.11

1-separations:



2-separations:




Tutte Connectivity

Definition 5.10


- ▶ A **k -separation** of a graph G is a pair of subgraphs $\{A, B\}$ of G such that each of A and B has size at least k , $A \neq G$, $B \neq G$, $A \cup B = G$, and $A \cap B$ is trivial of order at most k .
- ▶ If G has a k -separation for some k , then **Tutte connectivity** of G is $\min\{j : G \text{ has a } j \text{ separation}\}$, and ∞ if no k -separation exists.

Example 5.11

1-separations:



2-separations:



∞ -te Tutte connectivity:


Tutte Connectivity

Definition 5.10


- ▶ A **k -separation** of a graph G is a pair of subgraphs $\{A, B\}$ of G such that each of A and B has size at least k , $A \neq G$, $B \neq G$, $A \cup B = G$, and $A \cap B$ is trivial of order at most k .
- ▶ If G has a k -separation for some k , then **Tutte connectivity** of G is $\min\{j : G \text{ has a } j \text{ separation}\}$, and ∞ if no k -separation exists.

Example 5.11

1-separations:



2-separations:



∞ -te Tutte connectivity: \emptyset


Tutte Connectivity

Definition 5.10

- ▶ A **k -separation** of a graph G is a pair of subgraphs $\{A, B\}$ of G such that each of A and B has size at least k , $A \neq G$, $B \neq G$, $A \cup B = G$, and $A \cap B$ is trivial of order at most k .
- ▶ If G has a k -separation for some k , then **Tutte connectivity** of G is $\min\{j : G \text{ has a } j \text{ separation}\}$, and ∞ if no k -separation exists.

Example 5.11

1-separations:



2-separations:



∞ -te Tutte connectivity: \emptyset •


Tutte Connectivity

Definition 5.10


- ▶ A **k -separation** of a graph G is a pair of subgraphs $\{A, B\}$ of G such that each of A and B has size at least k , $A \neq G$, $B \neq G$, $A \cup B = G$, and $A \cap B$ is trivial of order at most k .
- ▶ If G has a k -separation for some k , then **Tutte connectivity** of G is $\min\{j : G \text{ has a } j \text{ separation}\}$, and ∞ if no k -separation exists.

Example 5.11

1-separations:



2-separations:



∞ -te Tutte connectivity: \emptyset




Tutte Connectivity

Definition 5.10


- ▶ A **k -separation** of a graph G is a pair of subgraphs $\{A, B\}$ of G such that each of A and B has size at least k , $A \neq G$, $B \neq G$, $A \cup B = G$, and $A \cap B$ is trivial of order at most k .
- ▶ If G has a k -separation for some k , then **Tutte connectivity** of G is $\min\{j : G \text{ has a } j \text{ separation}\}$, and ∞ if no k -separation exists.

Example 5.11

1-separations:



2-separations:



∞ -te Tutte connectivity: \emptyset




Tutte Connectivity

Definition 5.10


- ▶ A **k -separation** of a graph G is a pair of subgraphs $\{A, B\}$ of G such that each of A and B has size at least k , $A \neq G$, $B \neq G$, $A \cup B = G$, and $A \cap B$ is trivial of order at most k .
- ▶ If G has a k -separation for some k , then **Tutte connectivity** of G is $\min\{j : G \text{ has a } j \text{ separation}\}$, and ∞ if no k -separation exists.

Example 5.11

1-separations:



2-separations:



∞ -te Tutte connectivity: \emptyset

•




Tutte Connectivity

Definition 5.10


- ▶ A **k -separation** of a graph G is a pair of subgraphs $\{A, B\}$ of G such that each of A and B has size at least k , $A \neq G$, $B \neq G$, $A \cup B = G$, and $A \cap B$ is trivial of order at most k .
- ▶ If G has a k -separation for some k , then **Tutte connectivity** of G is $\min\{j : G \text{ has a } j \text{ separation}\}$, and ∞ if no k -separation exists.

Example 5.11

1-separations:



2-separations:



∞ -te Tutte connectivity: \emptyset

•




Tutte Connectivity

Definition 5.10


- ▶ A **k -separation** of a graph G is a pair of subgraphs $\{A, B\}$ of G such that each of A and B has size at least k , $A \neq G$, $B \neq G$, $A \cup B = G$, and $A \cap B$ is trivial of order at most k .
- ▶ If G has a k -separation for some k , then **Tutte connectivity** of G is $\min\{j : G \text{ has a } j \text{ separation}\}$, and ∞ if no k -separation exists.

Example 5.11

1-separations:



2-separations:



∞ -te Tutte connectivity: \emptyset



Tutte Connectivity vs. Vertex Connectivity

Theorem 5.12

*If G is a graph on at least 3 vertices and $G \not\cong K_3$, then the Tutte connectivity of G is $\min(\kappa(G), g(G))$, where $g(G)$ is the **girth** of G , that is, the length of a shortest cycle in G .*

Tutte Connectivity vs. Vertex Connectivity

Theorem 5.12

*If G is a graph on at least 3 vertices and $G \not\cong K_3$, then the Tutte connectivity of G is $\min(\kappa(G), g(G))$, where $g(G)$ is the **girth** of G , that is, the length of a shortest cycle in G .*

Tutte Connectivity vs. Vertex Connectivity

Theorem 5.12

*If G is a graph on at least 3 vertices and $G \not\cong K_3$, then the Tutte connectivity of G is $\min(\kappa(G), g(G))$, where $g(G)$ is the **girth** of G , that is, the length of a shortest cycle in G .*

Definition 5.13

- ▶ A **component** of a graph G is a maximal subgraph of G that has Tutte connectivity at least 1.

Tutte Connectivity vs. Vertex Connectivity

Theorem 5.12

*If G is a graph on at least 3 vertices and $G \not\cong K_3$, then the Tutte connectivity of G is $\min(\kappa(G), g(G))$, where $g(G)$ is the **girth** of G , that is, the length of a shortest cycle in G .*

Definition 5.13

- ▶ A **component** of a graph G is a maximal subgraph of G that has Tutte connectivity at least 1.
- ▶ A **block** of a graph G is a maximal subgraph of G that has Tutte connectivity at least 2.

Tutte Connectivity vs. Vertex Connectivity

Theorem 5.12

*If G is a graph on at least 3 vertices and $G \not\cong K_3$, then the Tutte connectivity of G is $\min(\kappa(G), g(G))$, where $g(G)$ is the **girth** of G , that is, the length of a shortest cycle in G .*

Definition 5.13

- ▶ A **component** of a graph G is a maximal subgraph of G that has Tutte connectivity at least 1.
- ▶ A **block** of a graph G is a maximal subgraph of G that has Tutte connectivity at least 2.

Note 5.14

A block of a non-empty graph is an isolated vertex, a loop-graph, a graph on two vertices with a positive number of edges between those vertices, or is vertex-2-connected.

Note 5.15

Two distinct blocks in a graph share at most one vertex since otherwise their union would be Tutte-2-connected.

Block Tree

Note 5.15

Two distinct blocks in a graph share at most one vertex since otherwise their union would be Tutte-2-connected.

Definition 5.16

The **block-tree** of a connected graph G is a tree T whose vertex set is the disjoint union of the blocks of G

Note 5.15

Two distinct blocks in a graph share at most one vertex since otherwise their union would be Tutte-2-connected.

Definition 5.16

The **block-tree** of a connected graph G is a tree T whose vertex set is the disjoint union of the blocks of G and those vertices of G that belong to more than one block.

Note 5.15

Two distinct blocks in a graph share at most one vertex since otherwise their union would be Tutte-2-connected.

Definition 5.16

The **block-tree** of a connected graph G is a tree T whose vertex set is the disjoint union of the blocks of G and those vertices of G that belong to more than one block. The only edges in T are those that join vertices of G to blocks that contain them.

Block Tree


Note 5.15

Two distinct blocks in a graph share at most one vertex since otherwise their union would be Tutte-2-connected.

Definition 5.16

The **block-tree** of a connected graph G is a tree T whose vertex set is the disjoint union of the blocks of G and those vertices of G that belong to more than one block. The only edges in T are those that join vertices of G to blocks that contain them.

Example 5.17



Block Tree


Note 5.15

Two distinct blocks in a graph share at most one vertex since otherwise their union would be Tutte-2-connected.

Definition 5.16

The **block-tree** of a connected graph G is a tree T whose vertex set is the disjoint union of the blocks of G and those vertices of G that belong to more than one block. The only edges in T are those that join vertices of G to blocks that contain them.

Example 5.17



Block Tree


Note 5.15

Two distinct blocks in a graph share at most one vertex since otherwise their union would be Tutte-2-connected.

Definition 5.16

The **block-tree** of a connected graph G is a tree T whose vertex set is the disjoint union of the blocks of G and those vertices of G that belong to more than one block. The only edges in T are those that join vertices of G to blocks that contain them.

Example 5.17



Block Tree


Note 5.15

Two distinct blocks in a graph share at most one vertex since otherwise their union would be Tutte-2-connected.

Definition 5.16

The **block-tree** of a connected graph G is a tree T whose vertex set is the disjoint union of the blocks of G and those vertices of G that belong to more than one block. The only edges in T are those that join vertices of G to blocks that contain them.

Example 5.17



Block Tree


Note 5.15

Two distinct blocks in a graph share at most one vertex since otherwise their union would be Tutte-2-connected.

Definition 5.16

The **block-tree** of a connected graph G is a tree T whose vertex set is the disjoint union of the blocks of G and those vertices of G that belong to more than one block. The only edges in T are those that join vertices of G to blocks that contain them.

Example 5.17



Whitney's Characterization of 2-Connected Graphs

Definition 5.18

Two paths are **internally-disjoint** if neither contains a non-endpoint of the other.

Whitney's Characterization of 2-Connected Graphs

Definition 5.18

Two paths are **internally-disjoint** if neither contains a non-endpoint of the other.

Theorem 5.19 (Whitney)

A graph with at least three vertices is 2-connected if and only if each pair u and v of vertices is connected by a pair internally-disjoint uv -paths.

Expansion Lemma

Lemma 5.20 (Expansion Lemma)

If G is a k -connected graph and G' is obtained from G by adding a new vertex y adjacent to at least k vertices of G , then G' is also k -connected.

Expansion Lemma

Lemma 5.20 (Expansion Lemma)

If G is a k -connected graph and G' is obtained from G by adding a new vertex y adjacent to at least k vertices of G , then G' is also k -connected.

Proof.

Suppose S is a separating set of G' .



Expansion Lemma

Lemma 5.20 (Expansion Lemma)

If G is a k -connected graph and G' is obtained from G by adding a new vertex y adjacent to at least k vertices of G , then G' is also k -connected.

Proof.

Suppose S is a separating set of G' . If $y \in S$, then $S - y$ separates G , so $|S| \geq k + 1$.



Expansion Lemma

Lemma 5.20 (Expansion Lemma)

If G is a k -connected graph and G' is obtained from G by adding a new vertex y adjacent to at least k vertices of G , then G' is also k -connected.

Proof.

Suppose S is a separating set of G' . If $y \in S$, then $S - y$ separates G , so $|S| \geq k + 1$. If $y \notin S$ and $N(y) \subseteq S$, then $|S| \geq k$.



Expansion Lemma

Lemma 5.20 (Expansion Lemma)

If G is a k -connected graph and G' is obtained from G by adding a new vertex y adjacent to at least k vertices of G , then G' is also k -connected.

Proof.

Suppose S is a separating set of G' . If $y \in S$, then $S - y$ separates G , so $|S| \geq k + 1$. If $y \notin S$ and $N(y) \subseteq S$, then $|S| \geq k$. Otherwise, S must separate G , and again $|S| \geq k$. □

Characterization of 2-Connected Graphs

Theorem 5.21

If G is simple and $|G| \geq 3$, then the following are equivalent (and characterize simple 2-connected graphs):

Characterization of 2-Connected Graphs

Theorem 5.21

If G is simple and $|G| \geq 3$, then the following are equivalent (and characterize simple 2-connected graphs):

(A) *G is connected and has no cut-vertex;*

Characterization of 2-Connected Graphs

Theorem 5.21

If G is simple and $|G| \geq 3$, then the following are equivalent (and characterize simple 2-connected graphs):

- (A) *G is connected and has no cut-vertex;*
- (B) *For every two vertices x and y of G , there are two internally-disjoint xy -paths;*

Characterization of 2-Connected Graphs

Theorem 5.21

If G is simple and $|G| \geq 3$, then the following are equivalent (and characterize simple 2-connected graphs):

- (A) *G is connected and has no cut-vertex;*
- (B) *For every two vertices x and y of G , there are two internally-disjoint xy -paths;*
- (C) *For every two vertices x and y of G , there is a cycle through x and y .*

Characterization of 2-Connected Graphs

Theorem 5.21

If G is simple and $|G| \geq 3$, then the following are equivalent (and characterize simple 2-connected graphs):

- (A) G is connected and has no cut-vertex;*
- (B) For every two vertices x and y of G , there are two internally-disjoint xy -paths;*
- (C) For every two vertices x and y of G , there is a cycle through x and y .*
- (D) $\delta \geq 1$ and every pair of edges of G lies on a common cycle.*

Definition 5.22

- ▶ **Subdividing** an edge uv of a graph G is the operation of deleting uv and adding a path uwv through a new vertex w .

Subdivisions

Definition 5.22

- ▶ **Subdividing** an edge uv of a graph G is the operation of deleting uv and adding a path uvw through a new vertex w .
- ▶ A graph G is a **subdivision** of a graph H if G can be obtained from H by successively subdividing (zero or more) edges.

Definition 5.22

- ▶ **Subdividing** an edge uv of a graph G is the operation of deleting uv and adding a path uvw through a new vertex w .
- ▶ A graph G is a **subdivision** of a graph H if G can be obtained from H by successively subdividing (zero or more) edges.
- ▶ A graph H is a **topological minor** of G , written $H \leq_t G$, if a subgraph of G is a subdivision of H .

Definition 5.22

- ▶ **Subdividing** an edge uv of a graph G is the operation of deleting uv and adding a path uvw through a new vertex w .
- ▶ A graph G is a **subdivision** of a graph H if G can be obtained from H by successively subdividing (zero or more) edges.
- ▶ A graph H is a **topological minor** of G , written $H \leq_t G$, if a subgraph of G is a subdivision of H .
- ▶ A graph is a **topological minor** of G if it can be obtained from G by a sequence of operations each of which is one of the following:

Definition 5.22

- ▶ **Subdividing** an edge uv of a graph G is the operation of deleting uv and adding a path uvw through a new vertex w .
- ▶ A graph G is a **subdivision** of a graph H if G can be obtained from H by successively subdividing (zero or more) edges.
- ▶ A graph H is a **topological minor** of G , written $H \leq_t G$, if a subgraph of G is a subdivision of H .
- ▶ A graph is a **topological minor** of G if it can be obtained from G by a sequence of operations each of which is one of the following:
 - ▶ deleting an edge;

Definition 5.22

- ▶ **Subdividing** an edge uv of a graph G is the operation of deleting uv and adding a path uvw through a new vertex w .
- ▶ A graph G is a **subdivision** of a graph H if G can be obtained from H by successively subdividing (zero or more) edges.
- ▶ A graph H is a **topological minor** of G , written $H \leq_t G$, if a subgraph of G is a subdivision of H .
- ▶ A graph is a **topological minor** of G if it can be obtained from G by a sequence of operations each of which is one of the following:
 - ▶ deleting an edge;
 - ▶ deleting a vertex; and

Definition 5.22

- ▶ **Subdividing** an edge uv of a graph G is the operation of deleting uv and adding a path uvw through a new vertex w .
- ▶ A graph G is a **subdivision** of a graph H if G can be obtained from H by successively subdividing (zero or more) edges.
- ▶ A graph H is a **topological minor** of G , written $H \leq_t G$, if a subgraph of G is a subdivision of H .
- ▶ A graph is a **topological minor** of G if it can be obtained from G by a sequence of operations each of which is one of the following:
 - ▶ deleting an edge;
 - ▶ deleting a vertex; and
 - ▶ contracting an edge incident with a vertex of degree two (un-subdividing an edge).

Subdivisions and 2-Connectedness

Corollary 5.23

A subdivision of a 2-connected graph is also 2-connected.

Definition 5.24

- ▶ A **path addition** to G is the addition to G of a path of length $\ell \geq 1$ between two vertices of G , introducing $\ell - 1$ new vertices.

Definition 5.24

- ▶ A **path addition** to G is the addition to G of a path of length $\ell \geq 1$ between two vertices of G , introducing $\ell - 1$ new vertices.
- ▶ The added path is an **ear**.


Definition 5.24

- ▶ A **path addition** to G is the addition to G of a path of length $\ell \geq 1$ between two vertices of G , introducing $\ell - 1$ new vertices.
- ▶ The added path is an **ear**.
- ▶ An **ear decomposition** is a partition of $E(G)$ into sets R_0, R_1, \dots, R_k so that $C = R_0$ is a cycle, and R_i , for $i > 0$, is a path addition to the graph $R_0 \cup R_1 \cup \dots \cup R_{i-1}$.

Definition 5.24

- ▶ A **path addition** to G is the addition to G of a path of length $\ell \geq 1$ between two vertices of G , introducing $\ell - 1$ new vertices.
- ▶ The added path is an **ear**.
- ▶ An **ear decomposition** is a partition of $E(G)$ into sets R_0, R_1, \dots, R_k so that $C = R_0$ is a cycle, and R_i , for $i > 0$, is a path addition to the graph $R_0 \cup R_1 \cup \dots \cup R_{i-1}$.



Example 5.25



Definition 5.24

- ▶ A **path addition** to G is the addition to G of a path of length $\ell \geq 1$ between two vertices of G , introducing $\ell - 1$ new vertices.
- ▶ The added path is an **ear**.
- ▶ An **ear decomposition** is a partition of $E(G)$ into sets R_0, R_1, \dots, R_k so that $C = R_0$ is a cycle, and R_i , for $i > 0$, is a path addition to the graph $R_0 \cup R_1 \cup \dots \cup R_{i-1}$.



Example 5.25



Definition 5.24

- ▶ A **path addition** to G is the addition to G of a path of length $\ell \geq 1$ between two vertices of G , introducing $\ell - 1$ new vertices.
- ▶ The added path is an **ear**.
- ▶ An **ear decomposition** is a partition of $E(G)$ into sets R_0, R_1, \dots, R_k so that $C = R_0$ is a cycle, and R_i , for $i > 0$, is a path addition to the graph $R_0 \cup R_1 \cup \dots \cup R_{i-1}$.



Example 5.25



Definition 5.24

- ▶ A **path addition** to G is the addition to G of a path of length $\ell \geq 1$ between two vertices of G , introducing $\ell - 1$ new vertices.
- ▶ The added path is an **ear**.
- ▶ An **ear decomposition** is a partition of $E(G)$ into sets R_0, R_1, \dots, R_k so that $C = R_0$ is a cycle, and R_i , for $i > 0$, is a path addition to the graph $R_0 \cup R_1 \cup \dots \cup R_{i-1}$.



Example 5.25



Definition 5.24

- ▶ A **path addition** to G is the addition to G of a path of length $\ell \geq 1$ between two vertices of G , introducing $\ell - 1$ new vertices.
- ▶ The added path is an **ear**.
- ▶ An **ear decomposition** is a partition of $E(G)$ into sets R_0, R_1, \dots, R_k so that $C = R_0$ is a cycle, and R_i , for $i > 0$, is a path addition to the graph $R_0 \cup R_1 \cup \dots \cup R_{i-1}$.



Example 5.25



Definition 5.24

- ▶ A **path addition** to G is the addition to G of a path of length $\ell \geq 1$ between two vertices of G , introducing $\ell - 1$ new vertices.
- ▶ The added path is an **ear**.
- ▶ An **ear decomposition** is a partition of $E(G)$ into sets R_0, R_1, \dots, R_k so that $C = R_0$ is a cycle, and R_i , for $i > 0$, is a path addition to the graph $R_0 \cup R_1 \cup \dots \cup R_{i-1}$.

Example 5.25



Whitney's Ear Decomposition

Theorem 5.26 (Whitney's Ear Decomposition)

A simple graph is 2-connected if and only if it has an ear decomposition.

Whitney's Ear Decomposition

Theorem 5.26 (Whitney's Ear Decomposition)

*A simple graph is 2-connected if and only if it has an ear decomposition.
Furthermore, every cycle in a 2-connected graph is the initial cycle of some ear decomposition.*

Closed-Ear Decomposition

Definition 5.27

A **closed-ear decomposition** of a graph G is a partition of $E(G)$ into sets R_0, R_1, \dots, R_k such that R_0 is a cycle and R_i for $i > 0$ is either a path addition

Closed-Ear Decomposition

Definition 5.27

A **closed-ear decomposition** of a graph G is a partition of $E(G)$ into sets R_0, R_1, \dots, R_k such that R_0 is a cycle and R_i for $i > 0$ is either a path addition or a cycle with exactly one vertex in $R_0 \cup R_1 \cup \dots \cup R_{i-1}$ (**closed ear**).

Closed-Ear Decomposition

Definition 5.27

A **closed-ear decomposition** of a graph G is a partition of $E(G)$ into sets R_0, R_1, \dots, R_k such that R_0 is a cycle and R_i for $i > 0$ is either a path addition or a cycle with exactly one vertex in $R_0 \cup R_1 \cup \dots \cup R_{i-1}$ (**closed ear**).

Theorem 5.28

A simple graph is 2-edge-connected if and only if it has a closed-ear decomposition.

Closed-Ear Decomposition

Definition 5.27

A **closed-ear decomposition** of a graph G is a partition of $E(G)$ into sets R_0, R_1, \dots, R_k such that R_0 is a cycle and R_i for $i > 0$ is either a path addition or a cycle with exactly one vertex in $R_0 \cup R_1 \cup \dots \cup R_{i-1}$ (**closed ear**).

Theorem 5.28

A simple graph is 2-edge-connected if and only if it has a closed-ear decomposition. Moreover, every cycle in a 2-edge-connected graph is the initial cycle in some closed-ear decomposition.

The Menger Theorem

Theorem 5.29 (Menger 1927)

If x and y are non-adjacent distinct vertices of a graph G , then the minimum size of a vertex-cut separating x from y equals the maximum number of pairwise internally-disjoint xy -paths.

The Edge Version of Menger's Theorem

Theorem 5.30 (Edge Version of Menger's Theorem)

If x and y are distinct vertices of a graph, then the minimum size $\kappa'(x, y)$ of the set of edges that separate x from y equals the maximum number $\lambda'(x, y)$ of pairwise edge-disjoint xy -paths.

The Edge Version of Menger's Theorem

Theorem 5.30 (Edge Version of Menger's Theorem)

If x and y are distinct vertices of a graph, then the minimum size $\kappa'(x, y)$ of the set of edges that separate x from y equals the maximum number $\lambda'(x, y)$ of pairwise edge-disjoint xy -paths.

Definition 5.31

The **line graph** of a graph G , written $L(G)$, is a simple graph whose vertex set is $E(G)$ with two vertices adjacent if the corresponding edges are adjacent in G .

The Edge Version of Menger's Theorem


Theorem 5.30 (Edge Version of Menger's Theorem)

If x and y are distinct vertices of a graph, then the minimum size $\kappa'(x, y)$ of the set of edges that separate x from y equals the maximum number $\lambda'(x, y)$ of pairwise edge-disjoint xy -paths.

Definition 5.31

The **line graph** of a graph G , written $L(G)$, is a simple graph whose vertex set is $E(G)$ with two vertices adjacent if the corresponding edges are adjacent in G .

Example 5.32



The Edge Version of Menger's Theorem


Theorem 5.30 (Edge Version of Menger's Theorem)

If x and y are distinct vertices of a graph, then the minimum size $\kappa'(x, y)$ of the set of edges that separate x from y equals the maximum number $\lambda'(x, y)$ of pairwise edge-disjoint xy -paths.

Definition 5.31

The **line graph** of a graph G , written $L(G)$, is a simple graph whose vertex set is $E(G)$ with two vertices adjacent if the corresponding edges are adjacent in G .

Example 5.32



The Edge Version of Menger's Theorem



Theorem 5.30 (Edge Version of Menger's Theorem)

If x and y are distinct vertices of a graph, then the minimum size $\kappa'(x, y)$ of the set of edges that separate x from y equals the maximum number $\lambda'(x, y)$ of pairwise edge-disjoint xy -paths.

Definition 5.31

The **line graph** of a graph G , written $L(G)$, is a simple graph whose vertex set is $E(G)$ with two vertices adjacent if the corresponding edges are adjacent in G .

Example 5.32



Tutte's Wheel Theorem

Theorem 5.34 (Tutte's Wheel Theorem)

If G is a Tutte-3-connected graph on at least four vertices that is not a wheel, then there is an edge e of G such that at least one of G/e and $G \setminus e$ is also Tutte-3-connected.

Tutte's Wheel Theorem

Theorem 5.34 (Tutte's Wheel Theorem)

If G is a Tutte-3-connected graph on at least four vertices that is not a wheel, then there is an edge e of G such that at least one of G/e and $G \setminus e$ is also Tutte-3-connected.

Lemma 5.35 (Thomassen 1980)

Every 3-connected graph G on at least five vertices has an edge e such that G/e is 3-connected.