

Linear Algebra, Bernard Kolman

Q1

$$\begin{array}{l} x+2y=8 \longrightarrow \textcircled{1} \\ 3x-4y=4 \longrightarrow \textcircled{2} \end{array}$$

xing equation $\textcircled{1}$ by 3 and subtract from $\textcircled{2}$

$$\begin{array}{r} 3x+6y=24 \\ 3x-4y=4 \\ \hline 10y=20 \Rightarrow y=2 \end{array}$$

$$\text{From } \textcircled{1} \Rightarrow x+2(2)=8 \Rightarrow x=8-4=4 \Rightarrow \boxed{x=4}$$

Hence $(x,y) = (4,2)$ which is unique solution.

Q3

$$\begin{array}{l} 3x+2y+z=2 \longrightarrow \text{(i)} \\ 4x+2y+2z=8 \longrightarrow \text{(ii)} \\ x-y+z=4 \longrightarrow \text{(iii)} \end{array}$$

Solving eq(i) & (iii)

xing eq(iii) by 3 then subtract from (i)

$$\begin{array}{r} 3x-3y+3z=12 \\ 3x+2y+z=8 \\ \hline -5y+2z=10 \end{array} \quad \text{(iv)}$$

(2)
Ex: 1.1

Name Solving equation (iii) + (ii)

Xing equation (iii) by 4 then subtract from (ii)

$$\begin{array}{r} 4x - 4y + 4z = 16 \\ 4x + 2y + 2z = 8 \\ \hline -6y + 2z = 8 \end{array}$$

Now solve eq (iv) & (v).

(iv) - (v)

$$\begin{array}{r} -8y + 2z = 10 \\ -6y + 2z = 8 \\ \hline -2y = 2 \end{array}$$

$$2y = 2 \Rightarrow y = 1$$

$$\Rightarrow 4y = 3z + z \Rightarrow y = z/4 \quad (\text{iii})$$

where z is any real no then $\bar{z} = z$

$$\sqrt{(\text{iii})} \Rightarrow y = z/4$$

$$\sqrt{(\text{ii})} \Rightarrow x = 12 - 4y + z$$

$$x = 12 - 4(1) + z$$

$$x = 12 - 3z - \frac{4z}{4} + z$$

$$x = 12 - 3z - z + z$$

$$x = -20$$

$$3x + 2(z) + 10 = 2$$

$$3x = 2 - 14 = -12$$

$$\frac{3x}{3} = \frac{-12}{3} = \boxed{x = -4}$$

$$\text{Hence } (x, y, z) = (-4, 1, 10) \text{ which is unique solution.}$$

$$\begin{aligned} x + 2y - z &= 12 \quad (\text{i}) \\ 3x + 8y - 2z &= 4 \quad (\text{ii}) \end{aligned}$$

Xing equ. (i) by 3 then sub. from (ii)

$$\begin{array}{r} 3x + 12y - 3z = 36 \\ 3x + 8y - 2z = 4 \\ \hline 4y - z = 32 \end{array}$$

Now solving eq (iv) & (v).

(iv) - (v)

$$\begin{array}{r} -8y + 2z = 10 \\ -6y + 2z = 8 \\ \hline -2y = 2 \end{array}$$

$$\boxed{y = 1}$$

$$\begin{array}{r} -8y + 2z = 10 \\ -8 + 2z = 10 \\ \hline 2z = 18 \end{array}$$

$$2z = 18 \Rightarrow z = 9$$

Putting the value of $y + z$ in (i)

$$3x + 2(z) + 10 = 2$$

$$3x = 2 - 14 = -12$$

$$\frac{3x}{3} = \frac{-12}{3} = \boxed{x = -4}$$

$$\text{Hence } (x, y, z) = (-4, 1, 10) \text{ which is unique solution.}$$

$$\begin{aligned}x+y+3z &= 12 \quad (i) \\2x+2y+6z &= 6 \quad (ii)\end{aligned}$$

$$\begin{aligned}10x+15y &= 65 \\10x+20y &= 54\end{aligned}$$

$$\frac{10x+20y}{10x+15y} = \frac{54}{65}$$

Xing eq (i) by (2) then subst. from (ii)

$$\begin{aligned}2x+2y+6z &= 24 \\2x+2y+6z &= 6 \\0 &= 18\end{aligned}$$

No solution.

Q

$$\begin{aligned}2x+3y &= 13 \quad (i) \\x-2y &= 3 \quad (ii)\end{aligned}$$

$$\begin{aligned}5x+2y &= 27 \quad (iii) \\5x+2y &= 27\end{aligned}$$

Xing eq (ii) by (3) and subst. from (i)

$$\begin{aligned}2x+3y &= 13 \\2x-4y &= 6 \\6y &= 19 \\y &= 1\end{aligned}$$

Xing eq (ii) by (3) then subst. from (iii)

$$\begin{aligned}5x-10y &= 15 \\5x+2y &= 27 \\-12y &= -12 \\y &= 1\end{aligned}$$

$$\begin{aligned}7y &= 7 \Rightarrow \boxed{y=1} \\I f z=4 &\text{ then } \text{eq (iii)} \Rightarrow 3y+4=7 \Rightarrow 3y=7-4=3\end{aligned}$$

$$\begin{aligned}3y &= 3 \Rightarrow \boxed{y=1} \\2x+1-2(4) &= -5 \\2x-7 &= -5 \\2x &= 2 \\x &= 1\end{aligned}$$

$$\begin{aligned}\text{Hence } x=1, y=1 &\text{ and } z=4 \text{ Ans.}\end{aligned}$$

Q17 without using the method of elimination solve the linear system

$$\begin{aligned}2x+y-2z &= -5 \quad (i) \\3y+z &= 7 \quad (ii) \\z &= 4 \quad (iii)\end{aligned}$$

Hence $y=1$ & $x=5$ which is unique solution

Put $y=1$ in (i) then

$$2x+3(1) = 13 \Rightarrow 2x = 13-3 = 10 \Rightarrow \boxed{x=5}$$

$$\begin{aligned}3y &= 3 \Rightarrow \boxed{y=1} \\I f z=4 &\text{ then } \text{eq (iii)} \Rightarrow 3y+4=7 \Rightarrow 3y=7-4=3\end{aligned}$$

Xing eq (i) by (5) & eq (iii) by 2 then subst:

Ex: 11

Q23 Let x_1 denotes lower-sulfur & x_2 denotes high-sulfur in each ton

$$5x_1 + 4x_2 = 3 \times 60 = 180$$

$$4x_1 + 2x_2 = 2 \times 60 = 120$$

$$2x_1 + 3x_2 = 11 \quad \text{--- (i)}$$

$$x_1 + 2x_2 = 7 \quad \text{--- (ii)}$$

$$4x_1 + 2x_2 = 12 \quad \text{--- (iii)}$$

Given that $x_1 = 1, x_2 = 2, x_3 = 1$

$$\sqrt{(i) \Rightarrow 2(1) + 3(2) - x_3 = 11}$$

$$2 + 6 - 1 = 11 \Rightarrow -x_3 = 11 - 8 = 3 \Rightarrow \boxed{x_3 = -3}$$

$$\text{Eq (ii) } \Rightarrow 1 - 2 + 2x_2 = -7$$

$$-1 + 2x_2 = -7 \Rightarrow 2x_2 = -7 + 1 = -6$$

$$\boxed{x_2 = -3}$$

$$\begin{array}{l} 5x_1 + 4x_2 = 180 \\ -8x_1 + 2x_2 = 240 \end{array}$$

$$-3x_1 = -60 \Rightarrow x_1 = \frac{-60}{-3} = 20$$

$$\text{Put } x_1 = 20 \text{ in Eq (i)}$$

$$5x_1 + 4x_2 = 180$$

$$5 \times 20 + 4x_2 = 180$$

$$100 + 4x_2 = 180$$

$$4x_2 = 180 - 100 = 80$$

$$x_2 = \frac{80}{4} = 20$$

$$\boxed{x_2 = -3}$$

$$\boxed{\text{Hence } x_2 = -3}$$

Ans.

$$\boxed{x_1 = 20 \text{ tons}}$$

unique solution

(24) Let x denote regular plastic ton

π_2 denotes special plastic tom

$$2x_1 + 2x_2 = 8 \quad \text{--- (ii)}$$

Solving (i) & (ii)

卷之三

Q25
Let x = Number ofounce A
P

~~Y₂ denser at Dunc B
Y₃ denser at Dunc C~~

$$\begin{array}{ccc} A & B & C \\ 2x_1 + 2x_2 + 3x_3 = 25 & \end{array} \quad (ii)$$

$$3x_1 + 2x_2 + 3x_3 = 24 \quad \text{--- (ii)} \\ 4x_1 + x_2 + 2x_3 = 21 \quad \text{--- (iii)}$$

Xing eq (iii) by 2 from Subj: from (ii)

$$3x_1 + 2x_2 + 3x_3 = 24$$

$$-5x_1 - 3x_3 = -18$$

Xing qy (iii) by 3 from Subhi: *Lamai*

$$2x_1 + 3x_2 + 3x_3 = 25$$

$$\frac{1}{2}x_1 + \frac{3}{2}x_2 + 6x_3 = 63$$

$$\Rightarrow 10x_1 + 3x_3 = 38. \quad (\text{v})$$

Xinggu (in) bag 2

$$\begin{array}{l} 10x_1 + 20x_3 = 36 \\ 10x_1 + 30x_3 = 38 \end{array}$$

$$15x_1 + 3x_3 = 54$$

$$\frac{2x_1 + 2x_2}{2} = \frac{x_1 + x_2}{1}$$

$$e \cdot S = 12$$

Put $x_1 = 3.2$ & $x_3 = 2$ in eq(i)

$$\sqrt{5} = \sqrt{2(3+2) + 3\lambda_2 + 3(\lambda_2)} = 2\sqrt{5}$$

$$3x_2 = 25 - 12 \cdot 3 = 12 \cdot 6$$

$$x_2 = 4 \cdot 2$$

Ex. 1.1

6

Ques Let X_1 denotes tons of 2-minute
 X_2 denotes tons of 6-minute
 X_3 denotes tons of 9-minute

+ X_3 denotes tons of 9-minute

$$\text{Now } 6X_1 + 12X_2 + 12X_3 = 10 \times 60 = 600$$

$$24X_1 + 12X_2 + 12X_3 = 16 \times 60 = 960$$

N

$$6X_1 + 12X_2 + 12X_3 = 600$$

$$24X_1 + 12X_2 + 12X_3 = 960$$

$$X_1 + 2X_2 + 2X_3 = 100 \quad \text{--- (i)}$$

$$2X_1 + X_2 + X_3 = 80 \quad \text{--- (ii)}$$

X_1 \vee X_2 \vee X_3 by 2 then solve from (i)

$$X_1 + 2X_2 + 2X_3 = 100$$

$$\underline{6X_1 + 2X_2 + 2X_3 = 160}$$

$$\frac{-3X_1 = -60}{\boxed{X_1 = 20}}$$

X_1 \vee X_2 \vee X_3 by 2 then solve from (ii)

$$2X_1 + 4X_2 + 4X_3 = 200$$

$$\underline{2X_1 + X_2 + X_3 = 80}$$

$$\frac{3X_2 + 3X_3 = 120}{3X_2 + 3X_3 = 120}$$

$$\Rightarrow X_2 + X_3 = 40 \quad \text{--- (iii)}$$


Let $X_3 = Y$ any real no: $A + mY$.
From (iii) $\Rightarrow X_2 + Y = 40 \Rightarrow \boxed{X_2 = 40 - Y}$.

$$\boxed{X_1 = 20}, \boxed{X_2 = 40 - Y}, \boxed{X_3 = Y}$$

Ques Given that $P_1(w) = \frac{(w+4)(w-5)}{w}$

$$P_2(w) = (-1, 1)$$

$$P_3(w) = (2, 7)$$



+ Parabolic $P(w) = aw^2 + bw + c$.

$$(a) \quad P_1(w) = y = aw^2 + bw + c$$

$$-1 = a(-1)^2 + b(-1) + c$$

$$-5 = a + b + c \quad \text{--- (i)}$$

$$P_2(w) = y = aw^2 + bw + c$$

$$1 = a(2)^2 + b(2) + c$$

$$1 = 4a + 2b + c \quad \text{--- (ii)}$$

$$P_3(w) = y = aw^2 + bw + c$$

$$7 = 4a + 2b + c \quad \text{--- (iii)}$$

$$\alpha_1 \otimes \Rightarrow Z_1 = 24000 - 3X_1$$

$$Z_1 = 24000 - 3(7000)$$

$$Z_1 = 24000 - 21000$$

$$\boxed{Z_1 = 3000}$$

M. Jamal Nabi

Lecturer U.E.T Peshawar.

Spring - 2012 Solved Depth

$$\text{Q1} \quad \text{if } \begin{bmatrix} a+b & c+d \\ c-d & a-b \end{bmatrix} = \begin{bmatrix} 4 & 6 \\ 10 & 2 \end{bmatrix} \quad \text{find } a, b, c, d.$$

Sol:

$$a+b = 4 \quad \text{(i)}$$

$$c+d = 6 \quad \text{(ii)}$$

$$c-d = 10 \quad \text{(iii)}$$

$$a-b = 2 \quad \text{(iv)}$$

(i) + (iv)

$$a+b = 4$$

$$a-b = 2$$

$$2a = 6 \Rightarrow \boxed{a = 3}$$

$$\alpha_1 \text{ (i)} \Rightarrow 3+b = 4 \Rightarrow b = 4-3 = 1 \Rightarrow \boxed{b = 1}$$

(ii) + (iii)

$$c+d = 6$$

$$c-d = 10$$

$$2c = 16 \Rightarrow \boxed{c = 8}$$

$$8+d = 6 \Rightarrow d = 6-8 = -2 \Rightarrow \boxed{d = -2}$$

$$\text{Q1} \quad \text{if } A = \begin{bmatrix} 2 & -3 & 5 \\ 6 & -5 & 4 \end{bmatrix}, \quad B = \begin{bmatrix} 4 & 3 \\ 6 & 5 \end{bmatrix} \quad \text{and} \quad C = \begin{bmatrix} 7 & 3 & 2 \\ 6 & 3 & 5 \end{bmatrix}$$

$$a_{12} = -3, \quad a_{22} = -5, \quad a_{23} = 4$$

$$b_{11} = 4, \quad b_{31} = 5$$

$$c_{13} = 2, \quad c_{31} = 6, \quad c_{33} = -1$$

Qatar Shabab
0300 5829015
Asstt. 0345 9289401
Ph. 051-5610987

Jan Laser Photostate

Matrix to M.A.
Notes are available

Shopping centre Bazaar Maria college Peshawar University

Q3 is similarly b Q2.

In exercise 4 through 7, let

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 1 & 4 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & 0 \\ 2 & 3 \\ 3 & 2 \end{bmatrix}, \quad C = \begin{bmatrix} 3 & -1 & 3 \\ 4 & 1 & -1 \\ 2 & -1 & 2 \end{bmatrix}$$

$$D = \begin{bmatrix} 3 & -2 \\ 2 & 4 \end{bmatrix}, \quad E = \begin{bmatrix} 2 & -4 & 5 \\ 0 & 1 & 4 \\ 3 & 2 & 1 \end{bmatrix}, \quad F = \begin{bmatrix} -1 & 3 \\ 2 & 3 \end{bmatrix}$$

and $O = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$.

Q4 a) $C+E$ & $E+C$

$$C+E = \begin{bmatrix} 3 & -1 & 3 \\ 4 & 1 & -1 \\ 2 & 1 & 3 \end{bmatrix} + \begin{bmatrix} 2 & -4 & 5 \\ 0 & 1 & 4 \\ 3 & 2 & 1 \end{bmatrix} = \begin{bmatrix} 5 & -5 & 8 \\ 4 & 2 & 9 \\ 3 & 4 & 4 \end{bmatrix}$$

$$+ E+C = \begin{bmatrix} 2 & -4 & 5 \\ 0 & 1 & 4 \\ 3 & 2 & 1 \end{bmatrix} + \begin{bmatrix} 3 & -1 & 3 \\ 4 & 1 & -1 \\ 2 & 1 & 3 \end{bmatrix} = \begin{bmatrix} 5 & -5 & 8 \\ 4 & 2 & 9 \\ 3 & 4 & 4 \end{bmatrix}.$$

commutative addition.

b)

$$A+B = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 1 & 4 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 2 & 1 \\ 3 & 2 \end{bmatrix}$$

addition is not possible because the order of matrices is not same.

$$= \begin{bmatrix} 1 & 4 \\ 10 & 18 \end{bmatrix}$$

(2) $D-F = \begin{bmatrix} 3 & -2 \\ 2 & 4 \end{bmatrix} - \begin{bmatrix} -1 & 5 \\ 2 & 3 \end{bmatrix} = \begin{bmatrix} 3+4 & -2-5 \\ 2-2 & 4-3 \end{bmatrix} = \begin{bmatrix} 7 & -7 \\ 0 & 1 \end{bmatrix}$

(3) $-3C+5O = -3 \begin{bmatrix} 3 & -1 & 3 \\ 4 & 1 & -1 \\ 2 & -1 & 2 \end{bmatrix} + 5 \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$
 $= \begin{bmatrix} -9 & 3 & -9 \\ -12 & -3 & -15 \\ -6 & -3 & -9 \end{bmatrix} + \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$
 $= \begin{bmatrix} -9 & 3 & -9 \\ -12 & -3 & -15 \\ -6 & -3 & -9 \end{bmatrix} \text{ & }$

(4) $2B+F = 2 \begin{bmatrix} 1 & 0 \\ 2 & 1 \\ 3 & 2 \end{bmatrix} + \begin{bmatrix} -1 & 5 \\ 2 & 3 \end{bmatrix}$

$$= \begin{bmatrix} 2 & 0 \\ 4 & 2 \\ 6 & 4 \end{bmatrix} + \begin{bmatrix} -1 & 5 \\ 2 & 3 \end{bmatrix}$$

addition is not possible

because the order of matrices is not same.

Q5 It's possible compute the indicated linear combination

(5) $3D+2F = 3 \begin{bmatrix} 3 & -2 \\ 2 & 4 \end{bmatrix} + 2 \begin{bmatrix} -1 & 5 \\ 2 & 3 \end{bmatrix} = \begin{bmatrix} 9 & -6 \\ 6 & 12 \end{bmatrix} + \begin{bmatrix} -8 & 10 \\ 4 & 6 \end{bmatrix}$

$(C+E)^T$ and $C^T + E^T$.

$$3(2A) \text{ and } 6A \\ 3(2A) = 6A \Rightarrow 6 \begin{bmatrix} 1 & 2 & 3 \\ 2 & 1 & 4 \end{bmatrix} = \begin{bmatrix} 6 & 12 & 18 \\ 12 & 6 & 24 \end{bmatrix}$$

$$(C+E)^t = \begin{bmatrix} 5 & -5 & 8 \\ 5 & 2 & 0 \\ 5 & 0 & 2 \end{bmatrix} = \begin{bmatrix} 5 & 4 & 5 \\ 5 & 0 & 2 \\ 5 & 2 & -1 \end{bmatrix}.$$

$$\textcircled{B} \quad 3(B+D) = 3 \left(\begin{bmatrix} 1 & 0 \\ 3 & 2 \end{bmatrix} + \begin{bmatrix} 3 & -2 \\ 2 & 4 \end{bmatrix} \right) \text{ addition is not} \\ \text{possible because the order of matrices is not same.}$$

$$C^t = \begin{bmatrix} 3 & 4 & 2 \\ 1 & 5 & 1 \\ 3 & 5 & 3 \end{bmatrix} \text{ and } E^t = \begin{bmatrix} 2 & 0 & 3 \\ 2 & 5 & 4 \\ 2 & 4 & -1 \end{bmatrix}.$$

Q6 If possible, compute:

a) A^T and $(A^t)^t$

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 1 & 4 \end{bmatrix}$$

$$A^t = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 1 & 4 \end{bmatrix}^t = \begin{bmatrix} 1 & 2 \\ 2 & 1 & 4 \end{bmatrix}.$$

$$d(A^t)^t = \begin{bmatrix} 1 & 2 \\ 2 & 1 & 4 \end{bmatrix}^t = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 1 & 4 \end{bmatrix}.$$

$$\therefore A^t = (A^t)^t.$$

Q7 is similarly to Q6.

Q8 Is the matrix $\begin{bmatrix} 4 & 1 \\ 0 & -3 \end{bmatrix}$ a linear combination of the matrices $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ and $\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$? Justify your answer.

$$\begin{bmatrix} 4 & 1 \\ 0 & -3 \end{bmatrix} = c_1 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + c_2 \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} c_1 & 0 \\ 0 & c_1 \end{bmatrix} + \begin{bmatrix} c_2 & 0 \\ 0 & 0 \end{bmatrix}.$$

$$c_1 + c_2 = 4 \quad c_1 = -3 \quad c_2 = 7$$

$$\text{So } \begin{bmatrix} 4 & 1 \\ 0 & -3 \end{bmatrix} \neq \begin{bmatrix} 4 & 0 \\ 0 & -3 \end{bmatrix}$$

So it's not a linear combination.

Q3 Let $A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ bit matrices.

Q1 Find B so that $A+B = \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$

$$\begin{aligned} B &= \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} - \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ 0 & 0 \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \text{ bit.} \end{aligned}$$

Q2 Find C so that $A+C = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$

$$\begin{aligned} C &= \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} - A \\ &= \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix}. \end{aligned}$$

Given that $a \cdot b = 17$ — (i)

Let $V = [a, b, c, d]$

$$\text{Then } [0 \ 1 \ 0 \ 1] + [a, b, c, d] = [1 \ 1 \ 1 \ 1]$$

$$[a \ b+1 \ c \ d+1] = [1 \ 1 \ 1 \ 1]$$

$$\boxed{a=1} \quad b+1=1 \Rightarrow \boxed{b=0} \quad \boxed{c=1} \quad d+1=1 \Rightarrow \boxed{d=0}$$

$$\text{So } V = [a \ b \ c \ d] = [1 \ 0 \ 1 \ 0].$$

Q1 (a) $a = [1 \ 1 \ 2], b = \begin{bmatrix} 4 \\ 1 \end{bmatrix}$. Find $a \cdot b$
 Sol: $a \cdot b = [1 \ 2] \cdot \begin{bmatrix} 4 \\ 1 \end{bmatrix}$
 $= 1 \times 4 + 2 \times 1$
 $= 4 + 2 = 6$ Ans.

Q2 is similarly to Q1

Q3 Let $a = [-3, 2 \ \pi]$ and $b = \begin{bmatrix} -3 \\ \pi \end{bmatrix}$ if $a \cdot b = 17$ find π .
 Sol: Given that $a \cdot b = 17$ — (i)
 $a \cdot b = [-3 \ 2 \ \pi] \cdot \begin{bmatrix} -3 \\ \pi \end{bmatrix}$
 $= -3 \times -3 + 2(\pi) + \pi \cdot \pi$
 $= 9 + 4\pi + \pi^2$

$$a \cdot b = 17 + \pi^2 \quad \text{--- (ii)}$$

Comparing eq (i) and eq (ii)

$$17 = 17 + \pi^2$$

$$\Rightarrow \pi^2 = 17 - 17 = 0$$

$$\pi^2 = 0$$

$$\boxed{\pi = \pm 2}$$

(2)

$$\sum x = 1.3$$

12

Q4 Let $W = \begin{bmatrix} \sin\theta \\ \cos\theta \end{bmatrix}$. compute $W \cdot W$

$$W \cdot W = \begin{bmatrix} \sin\theta \\ \cos\theta \end{bmatrix} \cdot \begin{bmatrix} \sin\theta \\ \cos\theta \end{bmatrix}$$

$$= \sin^2\theta + \cos^2\theta$$

$$[W \cdot W = 1] \text{ Ans}$$

Q5 Find all values of x so that $V \cdot V = 1$ where $V = \begin{bmatrix} V_1 \\ -V_2 \\ x \end{bmatrix}$.

Sol:

$$V \cdot V = 1$$

$$\begin{bmatrix} V_1 \\ -V_2 \\ x \end{bmatrix} \cdot \begin{bmatrix} V_1 \\ -V_2 \\ x \end{bmatrix} = 1$$

$$\left(\frac{1}{2}\right)(V_1) + (-V_2)(-V_2) + x \cdot x = 1$$

$$\frac{1}{2}V_1 + V_2 + x^2 = 1$$

$$\frac{1+1}{4} - 1 = -x^2$$

$$\frac{2}{4} - 1 = -x^2$$

$$\frac{1-2}{2} = -x^2$$

$$\cancel{\frac{1}{2}}V_2 = \cancel{\frac{1}{2}}x^2$$

$$x^2 = V_2 = \frac{2}{4}$$

$$x = \pm \frac{\sqrt{2}}{2} \text{ Ans}$$

$$Q6 \quad \text{If } A = \begin{bmatrix} 1 & 2 & x \\ 3 & -1 & 2 \end{bmatrix} \text{ and } B = \begin{bmatrix} y \\ z \\ x \end{bmatrix} \text{ if } AB = \begin{bmatrix} 6 \\ 8 \end{bmatrix} \text{ find } x+y.$$

Given that $AB = \begin{bmatrix} 6 \\ 8 \end{bmatrix}$ — (i)

$$\text{Now } A \cdot B = \begin{bmatrix} 1 & 2 & x \\ 3 & -1 & 2 \end{bmatrix} \begin{bmatrix} y \\ z \\ x \end{bmatrix}$$

$$= \begin{bmatrix} y+2x+x \\ 3y-x+2 \end{bmatrix}$$

$$A \cdot B = \begin{bmatrix} y+3x \\ 3y-x+2 \end{bmatrix} — (ii)$$

Comparing eq (i) and eq (ii)

$$\begin{bmatrix} y+3x \\ 3y-x+2 \end{bmatrix} = \begin{bmatrix} 6 \\ 8 \end{bmatrix}$$

$$y+3x = 6 — (i) \quad 3y-x+2 = 8 — (ii)$$

Xing eq (ii) by 3 from subtract from (ii)

$$3y+9x = 18$$

$$\frac{-3y-x}{10x = 12} \Rightarrow \boxed{x = 6/5}$$

$$2y = 12 \Rightarrow y + 3(6/5) = 6$$

$$y = 6 - 18/5 = \frac{30-18}{5} = 12/5$$

$$\boxed{y = 12/5}$$

212

5

卷之三

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} + B = \begin{bmatrix} 0 & 2 \\ 3 & 4 \end{bmatrix}$$

$S_{\text{max}}(f)$ $AB \neq BA$

$$\begin{array}{r} \text{A. } \\ \text{B. } \\ \text{C. } \\ \text{D. } \\ \text{E. } \\ \text{F. } \\ \text{G. } \\ \text{H. } \\ \text{I. } \\ \text{J. } \\ \text{K. } \\ \text{L. } \\ \text{M. } \\ \text{N. } \\ \text{O. } \\ \text{P. } \\ \text{Q. } \\ \text{R. } \\ \text{S. } \\ \text{T. } \\ \text{U. } \\ \text{V. } \\ \text{W. } \\ \text{X. } \\ \text{Y. } \\ \text{Z. } \end{array}$$

C₁ Col₁A₁ C₂ Col₂A₁ C₃ Col₃A₁

$$\frac{1}{\sqrt{1-\frac{v^2}{c^2}}}$$

OK is similarly 013

6

1

182

$$B \cdot A = \begin{bmatrix} 9 & -1 \\ -3 & 4 \\ 4 & 3 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix} = \begin{bmatrix} 3-3 & 4-2 \\ -3+8 & -6+8 \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 5 & 2 \end{bmatrix}$$

$$\frac{1}{1 - e^x} = \frac{e^x}{e^x - 1}$$

$$A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad B = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$

$$A = \begin{bmatrix} 1 & 2 & -1 \\ 3 & 1 & 4 \end{bmatrix}_{2 \times 3}$$

$$A_0 = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}_{2 \times 2}$$

$$B = \begin{bmatrix} 1 & -3 \\ 3 & -2 \\ -1 & 2 \\ 4w-5 & 9 \end{bmatrix}$$

$$col_2(AB) = A col_2(B)$$

$$\frac{1}{w} \left(\frac{1}{w} - \frac{1}{w_0} \right) + \frac{1}{w_0} \left(\frac{1}{w_0} - \frac{1}{w} \right)$$

$$\begin{array}{r}
 90 \Delta w - \\
 - 60 w \\
 \hline
 30 w \Delta 2 \\
 \hline
 130 \\
 \hline
 11 \\
 \hline
 1-3+8 \\
 3+6+16 \\
 4-6+12 \\
 2+3+20 \\
 \hline
 11 \\
 \hline
 6 \\
 25 \\
 10 \\
 \hline
 25
 \end{array}$$

) The 3rd column.

$$\begin{array}{r} \boxed{24w} \\ - \boxed{-9w^2w} \\ \hline \boxed{1w^4+9w} \\ \hline \boxed{w^3+w^2} \end{array}$$

$$A = \begin{bmatrix} 2 & -3 & 1 \\ -1 & 2 & 4 \end{bmatrix} \text{ and } B = \begin{bmatrix} 3 \\ 2 \\ 5 \end{bmatrix}$$

Scanned by CamScanner

a) Verify $AB = 3a_1 + 5a_2 + 2a_3$.

$$AB = \begin{bmatrix} 2 & -3 & 1 \\ 1 & 2 & 4 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \\ 2 \end{bmatrix}$$

$$= \begin{bmatrix} 6-15+2 \\ 3+10+8 \end{bmatrix} = \begin{bmatrix} -7 \\ 21 \end{bmatrix}. \quad \textcircled{1}$$

Now $3a_1 + 5a_2 + 2a_3 = 3\begin{bmatrix} 2 \end{bmatrix} + 5\begin{bmatrix} -3 \\ 2 \end{bmatrix} + 2\begin{bmatrix} 1 \end{bmatrix}$.

$$= \begin{bmatrix} 6 \end{bmatrix} + \begin{bmatrix} -15 \\ 10 \end{bmatrix} + \begin{bmatrix} 2 \\ 2 \end{bmatrix} = \begin{bmatrix} 6-15+2 \\ 3+10+8 \end{bmatrix} = \begin{bmatrix} -7 \\ 21 \end{bmatrix}$$

from Q1 & Q2 we have

$AB = 3a_1 + 5a_2 + 2a_3$ Hence Verified.

b) Verify that $AB = \begin{bmatrix} (\text{Row}_1(A))B \\ (\text{Row}_2(A))B \end{bmatrix}$.

from Part a) $AB = \begin{bmatrix} -7 \\ 21 \end{bmatrix}$. $\textcircled{2}$

$$\{ \text{Row}_1(A) \} B = \begin{bmatrix} 2 & -3 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \\ 2 \end{bmatrix} = \begin{bmatrix} 6-15+2 \end{bmatrix} = \begin{bmatrix} -7 \end{bmatrix}$$

$$\{ \text{Row}_2(A) \} B = \begin{bmatrix} 1 & 2 & 4 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \\ 2 \end{bmatrix} = \begin{bmatrix} 3+10+8 \end{bmatrix} = \begin{bmatrix} 21 \end{bmatrix}$$

$$\text{Now } \begin{bmatrix} (\text{Row}_1(A))B \\ (\text{Row}_2(A))B \end{bmatrix} = \begin{bmatrix} -7 \\ 21 \end{bmatrix} \rightarrow \textcircled{2}$$

from Q1 & Q2 Hence Verified.

Q21 Soln. $\begin{bmatrix} 2 & 0 & 0 & 1 & -7 \\ 2 & 3 & 2 & 0 & 1 \\ 1 & 0 & 3 & 0 & 1 \\ 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 1 & 5 \end{bmatrix}$

Q22 Soln. $\begin{bmatrix} 2 & 0 & 0 & 1 & -7 \\ 2 & 3 & 2 & 0 & 1 \\ 1 & 0 & 3 & 0 & 1 \\ 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 1 & 5 \end{bmatrix}$

Q23 Soln. $\begin{bmatrix} 2 & 0 & 0 & 1 & -7 \\ 2 & 3 & 2 & 0 & 1 \\ 1 & 0 & 3 & 0 & 1 \\ 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 1 & 5 \end{bmatrix}$

which is augmented matrix

Q24 Soln. $\begin{bmatrix} 2 & 0 & 0 & 1 & -7 \\ 2 & 3 & 2 & 0 & 1 \\ 1 & 0 & 3 & 0 & 1 \\ 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 1 & 5 \end{bmatrix}$

which is linear system in

matrix form:

$$\begin{bmatrix} 2 & 0 & 0 & 1 & -7 \\ 2 & 3 & 2 & 0 & 1 \\ 1 & 0 & 3 & 0 & 1 \\ 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 1 & 5 \end{bmatrix}$$

which is augmented matrix.

Q25 a) Soln.

$$\begin{aligned} -2x - y + 0z + 4w &= 5 \\ -3x + 2y + 7z + 8w &= 3 \\ x + 0y + 0z + 2w &= 4 \\ 3x + 0y + z + 3w &= 6 \end{aligned}$$

Q21 Similarly Q20
 Q22 Similarly Q19.

Q23 Sol: These are equivalent as third row have 3rd col only.

$$\text{Q24. a)} \begin{bmatrix} 1 & 2 & 0 \\ 2 & 5 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$$\text{b)} \begin{bmatrix} 1 & 2 & 1 \\ 1 & 1 & 2 \\ 2 & 0 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\text{Q25 a)} \begin{bmatrix} 1 & 2 \\ 2 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 3 \\ 5 \end{bmatrix} \quad \text{⑤} \quad \begin{bmatrix} 2 & -3 & 5 \\ 1 & 4 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$$

$$x \begin{bmatrix} 1 \\ 2 \end{bmatrix} + y \begin{bmatrix} 2 \\ -1 \end{bmatrix} = \begin{bmatrix} 3 \\ 5 \end{bmatrix}$$

$$x \begin{bmatrix} 2 \\ 1 \end{bmatrix} + y \begin{bmatrix} -3 \\ 4 \end{bmatrix} + z \begin{bmatrix} 5 \\ -1 \end{bmatrix} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$$

Q26 where AB gives total cost of producing each kind of product in each city.

$$AB = \begin{bmatrix} S.D & N.O & P.M \\ 300 & 100 & 150 \end{bmatrix} \text{ Product P}$$

$$\begin{bmatrix} 200 & 250 & 400 \end{bmatrix} \text{ Product Q}$$

$$B = \begin{bmatrix} \text{Plank} & \text{Plasty} \\ 8 & 12 \\ 7 & 9 \\ 15 & 10 \end{bmatrix} \begin{matrix} S.D \\ N.O \\ P.M \end{matrix}$$

- Q26
- comes say nothing.
 - can say nothing.

$$\text{Q27 a)} AB^t = 0$$

$$[Y \ 1 \ -2] \begin{bmatrix} 1 \\ 3 \\ -1 \end{bmatrix} = 0$$

$$1(Y) + 3(1) - 1(-2) = 0$$

$$\boxed{Y = -5}$$

Q28 Similarly Q27.

$$\begin{array}{c|cc} A.P & F.P & S.L.C \\ \hline A = \begin{bmatrix} 2 & 2 \\ 3 & 4 \end{bmatrix} & \text{chart} & B = \begin{bmatrix} 9 & 10 \\ 10 & 12 \end{bmatrix} \text{ F.P} \end{array}$$

S.L.C chart
S.L.C chart
A.P
F.P

Q29 where AB gives each product quantity in each city.

$$A \cdot B = \begin{bmatrix} 300 & 100 & 150 \\ 200 & 250 & 400 \end{bmatrix} \begin{bmatrix} 8 & 12 \\ 7 & 9 \\ 15 & 10 \end{bmatrix}$$

$$= \begin{bmatrix} \text{Plank} & \text{Plasty} \\ 5350 & 6000 \\ 9350 & 8650 \end{bmatrix} \text{ Product P, " Q.}$$

$$\text{Q33} \quad A = \begin{bmatrix} 33 \\ 80 & 120 \\ 100 & 200 \end{bmatrix} \begin{array}{l} \text{Adult} \\ \text{Child} \\ \text{Male} \\ \text{Female} \end{array}$$

$$\text{Q34} \quad \begin{bmatrix} P & F & C \\ 20 & 20 & 20 \\ 10 & 20 & 30 \end{bmatrix} \begin{array}{l} \text{Adult} \\ \text{Child} \\ \text{Male} \end{array}$$

$$\text{Q35} \quad a.b = \begin{bmatrix} 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} = 0$$

$$\text{Q36} \quad a.b = \begin{bmatrix} 1 & 0 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \end{bmatrix} = 1$$

$$\text{Q37} \quad 80 \times 20 + 10 \times 120 = 2800 \text{g}$$

$$\text{Q38} \quad \begin{bmatrix} 20 \\ 10 \\ 20 \end{bmatrix} \begin{bmatrix} 100 & 200 \end{bmatrix} = 20 \times 100 + 20 \times 200 = 6000 \text{g}$$

$$\text{Q39} \quad \begin{bmatrix} 220 \\ 550 \\ 120 \end{bmatrix} \begin{bmatrix} 200 & 150 & 120 \\ 120 \end{bmatrix} = 103400$$

$$\text{Q40} \quad \begin{bmatrix} 50 & 40 & 25 \end{bmatrix} \begin{bmatrix} 100 \\ 120 \\ 250 \end{bmatrix} = 16050$$

$$= 1\{x\} + 1\{x\} + 0\{1\} = 0$$

$$= x+x=0 \quad 2x=0 \Rightarrow \boxed{x=0}$$

$$\text{Q41} \quad a = \begin{bmatrix} 1 & x & 0 \end{bmatrix} \quad b = \begin{bmatrix} x \\ 1 \end{bmatrix} \quad \text{Find } x \quad \Delta a.b = 0$$

$$S_1 = \begin{bmatrix} 18.95 & 14.75 & 8.98 \end{bmatrix}$$

$$S_2 = \begin{bmatrix} 17.80 & 13.50 & 10.79 \end{bmatrix}$$

$$\psi = \begin{bmatrix} 18.95 & 14.75 & 8.98 \\ 17.80 & 13.50 & 10.79 \end{bmatrix}$$

b) 20% Reduced.

$$\psi = \begin{bmatrix} 18.95 \times 80\% & 14.75 \times 80\% & 8.98 \times 80\% \\ 17.80 \times 80\% & 13.50 \times 80\% & 10.79 \times 80\% \end{bmatrix}$$

$$\begin{bmatrix} 15.16 & 11.8 & 7.18 \\ 14.24 & 10.8 & 8.62 \end{bmatrix}$$

$$\text{Q42} \quad AB = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} a+c & b+d \\ c & d \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\boxed{c=0}, \boxed{d=1}$$

$$a+c=1 \Rightarrow a+0=1 \Rightarrow \boxed{a=1}$$

$$b+d=0 \Rightarrow b+1=0 \Rightarrow b=-1=1 \quad (\text{Ans})$$

$$B = \begin{bmatrix} 1 & +1 \\ 0 & 1 \end{bmatrix}$$

$\rightarrow X = \dots$

(b)

$$A+(B+C) = (A+B)+C.$$

$$B+C = \begin{bmatrix} -2 & -6 & 2 \\ 5 & 1 & 5 \end{bmatrix}$$

$$A+(B+C) = \begin{bmatrix} -1 & -4 & 0 \\ 8 & 5 & 10 \end{bmatrix}.$$

Similarly

$$A+B = \begin{bmatrix} 3 & 2 & -1 \\ 6 & 8 & 10 \end{bmatrix}$$

$$(A+B)+C = \begin{bmatrix} -1 & -4 & 0 \\ 8 & 5 & 10 \end{bmatrix}.$$

$$\text{Hence } A+(B+C) = (A+B)+C.$$

Verifying theorem 1.1
 (a) $A+B = B+A$

$$A+B = \begin{bmatrix} 3 & 2 & -1 \\ 6 & 8 & 10 \end{bmatrix} \text{ and } B+A = \begin{bmatrix} 3 & 2 & -1 \\ 6 & 8 & 10 \end{bmatrix}.$$

Mr. Tawar Nasir
Lecturer U.E.T Peshawar


Qaiser Shahab
0300 5829015
Arif: 0345 9289401
Ph: 091-5610987

Matrix to M.A.
Notes are available

Shop#58 coffee shop bazar Islamia college Peshawar university

Jan Laser Photostate


Matrix to M.A.
Notes are available

Qaiser Shahab
0300 5829015
Arif: 0345 9289401
Ph: 091-5610987

Shop#58 coffee shop bazar Islamia college Peshawar university

(2)

19

Q2 Verify (a) of theorem 1.2 for
 $A(BC) = (AB)C$.

$$\text{Sol: } BC = \begin{bmatrix} -1 & 3 & 2 \\ 1 & -3 & 4 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 3 & -1 \\ 1 & 2 \end{bmatrix} = \begin{bmatrix} -1+9+2 & -3+4 \\ 1-9+4 & 3+8 \end{bmatrix} = \begin{bmatrix} 10 & 1 \\ -2 & 11 \end{bmatrix}$$

$$A(BC) = \begin{bmatrix} 1 & 3 \\ 2 & -1 \end{bmatrix} \begin{bmatrix} 10 & 1 \\ -2 & 11 \end{bmatrix} = \begin{bmatrix} 10-12 & 1+33 \\ 20+4 & 2-11 \end{bmatrix} = \begin{bmatrix} -2 & 34 \\ 24 & -9 \end{bmatrix}$$

~~Ansatz~~

(a) $\check{Y}(SA) = (\check{Y}S)A$

$$SA = -2 \begin{bmatrix} 4 & 2 \\ 1 & -3 \end{bmatrix} = \begin{bmatrix} -8 & -4 \\ -2 & 6 \end{bmatrix}.$$

$$\check{Y}(SA) = 6 \begin{bmatrix} -8 & -4 \\ -2 & 6 \end{bmatrix} = \begin{bmatrix} -48 & -24 \\ -12 & 36 \end{bmatrix}.$$

$$(\check{Y}S) = (-2)(6) = 12$$

$$(\check{Y}S)A = -12 \begin{bmatrix} 4 & 2 \\ 1 & -3 \end{bmatrix} = \begin{bmatrix} -48 & -24 \\ -12 & 36 \end{bmatrix}.$$

Hence $\check{Y}(SA) = (\check{Y}S)A$ Verified.

Q3 Verify (b) of theorem 1.2

$$A(BC) = AB + AC$$

S.Y.S.

Verify (a), (b) and (c) of theorem 1.3 for $Y=6$, $S=-2$.

(a) $\check{Y}(SA) = (\check{Y}S)A$

(b) $(\check{Y}+S)A = \check{Y}A + SA$

(c) $\check{Y}(A+B) = \check{Y}A + \check{Y}B$

$$AB = \begin{bmatrix} 1 & 3 \\ 2 & -1 \end{bmatrix} \begin{bmatrix} -1 & 3 \\ 1 & -2 \end{bmatrix} = \begin{bmatrix} 9 & -6 \\ -2 & 9 \end{bmatrix}.$$

$$= \begin{bmatrix} -1+3 & 3-9 & 2+12 \\ -2-1 & 6+3 & 4-4 \end{bmatrix} = \begin{bmatrix} 9 & -6 & 14 \\ -3 & 9 & 0 \end{bmatrix}.$$

Scanned by CamScanner

$$Ax = y$$

$$A \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 2 & 1 & 1 \\ 1 & 2 & 1 \\ 1 & 1 & 2 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 2+1 \\ 1+2 \\ 1+2 \end{pmatrix} = \begin{pmatrix} 3 \\ 3 \\ 3 \end{pmatrix}$$

Q5 Verify (d) of Thm: 1.3 for $y = -3$
 $A(yB) = y(AB) = yA^T B$

S.Y.S.

Q6 Verify (b) & (d) of Thm: 1.4 for $y = -4$

Given $Ax = y$
 $\begin{pmatrix} 3 \\ 3 \\ 3 \end{pmatrix} = \begin{pmatrix} y \\ y \\ y \end{pmatrix} \Rightarrow \boxed{y = 3}$

(b) $(A+B)^T = A^T + B^T$
(c) $(yA)^T = yA^T$

S.Y.S.

Q7 Verify (c) of Thm: 1.4 for

(c) $(AB)^T = B^T A^T$

S.Y.S.

Q10, Q11 S.Y.S

Q12 If $A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ show $A^2 = I_2$

$$\frac{K^2 = 1}{\sqrt{K}} \\ K = \pm \sqrt{1}$$

$$A \cdot A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = I_2$$

Ans. $A^2 = I_2$

Q13 & Q14 S.Y.S.



Jan Laser Photostate
 Matrix to M.A.
 Notes are available

Qader Shahab
 0300 3820015
 Aarif: 0345 9289401
 Ph: 091-5610987

$$\text{Q18} \quad \text{a) } A\chi_0 = \begin{bmatrix} \frac{1}{3} & \frac{2}{3} \\ \frac{2}{3} & \frac{3}{5} \end{bmatrix} \begin{bmatrix} \frac{1}{3} \\ \frac{1}{5} \end{bmatrix}.$$

$$A\chi_0 = \begin{bmatrix} \frac{1}{3} + \frac{2}{5} \\ \frac{2}{3} + \frac{3}{5} \end{bmatrix}.$$

b) Find state distribution

$$A\chi_0 = \chi_0$$

$$\begin{bmatrix} \frac{1}{3} & \frac{2}{5} \\ \frac{2}{3} & \frac{3}{5} \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} a \\ b \end{bmatrix}$$

$$= \begin{bmatrix} \frac{1}{3}a + \frac{2}{5}b \\ \frac{2}{3}a + \frac{3}{5}b \end{bmatrix} = \begin{bmatrix} a \\ b \end{bmatrix}.$$

$$\begin{aligned} \frac{1}{3}a + \frac{2}{5}b &= a \quad \text{--- (i)} \\ \frac{2}{3}a + \frac{3}{5}b &= b \quad \text{--- (ii)} \end{aligned}$$

$$\gamma_0 \Rightarrow \frac{1}{3}a - a + \frac{2}{5}b = 0 \Rightarrow \frac{2}{5}b = 0$$

$$-\frac{2}{3}a + \frac{2}{5}b = 0$$

$$-10a + 6b = 0 \quad \text{--- (iii)}$$

$$\gamma_0 \Rightarrow \frac{2}{3}a + \frac{2}{5}b = 0$$

$$\Rightarrow 10a - 6b = 0 \quad \text{--- (iv)}$$

Solving (iii), (iv)

$$a = \frac{3}{8}, b = \frac{5}{8} \quad \chi_0 = \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} \frac{3}{8} \\ \frac{5}{8} \end{bmatrix}.$$

$$\text{Q19} \quad A = \begin{bmatrix} R & S & T \\ S & T & R \\ T & R & S \end{bmatrix} \quad \chi_0 = \begin{bmatrix} R \\ S \\ T \end{bmatrix}.$$

$$\text{Q19} \quad A\chi_0 = \begin{bmatrix} \frac{1}{3} & \frac{1}{2} & \frac{1}{4} \\ \frac{1}{2} & \frac{1}{4} & \frac{1}{2} \\ \frac{1}{4} & \frac{1}{2} & \frac{1}{3} \end{bmatrix} \begin{bmatrix} R \\ S \\ T \end{bmatrix} = \begin{bmatrix} \frac{13}{12} \\ \frac{17}{12} \\ \frac{11}{12} \end{bmatrix}.$$

$$A\chi_0 = \chi_0$$

$$\begin{bmatrix} \frac{1}{3} & \frac{1}{2} & \frac{1}{4} \\ \frac{1}{2} & \frac{1}{4} & \frac{1}{2} \\ \frac{1}{4} & \frac{1}{2} & \frac{1}{3} \end{bmatrix} \begin{bmatrix} R \\ S \\ T \end{bmatrix} = \begin{bmatrix} R \\ S \\ T \end{bmatrix}.$$

S. VS

$$\alpha = \frac{3}{4}R + \frac{1}{4}S = \frac{4}{7}R$$

$$\chi_0 = \begin{bmatrix} R \\ S \\ T \end{bmatrix} = \begin{bmatrix} \frac{4}{7}R \\ \frac{3}{7}R \\ \frac{1}{7}R \end{bmatrix}.$$

$$\text{Q19} \quad \text{Q19} \quad \chi_0 = \begin{bmatrix} R \\ S \\ T \end{bmatrix} \quad A = \begin{bmatrix} R & S & T \\ S & T & R \\ T & R & S \end{bmatrix}.$$

$$A\chi_0 = \begin{bmatrix} \frac{1}{3} & \frac{1}{2} & \frac{1}{4} \\ \frac{1}{2} & \frac{1}{4} & \frac{1}{2} \\ \frac{1}{4} & \frac{1}{2} & \frac{1}{3} \end{bmatrix} \begin{bmatrix} R \\ S \\ T \end{bmatrix} = \begin{bmatrix} \frac{13}{12} \\ \frac{17}{12} \\ \frac{11}{12} \end{bmatrix}.$$

after 2 years

$$\begin{aligned} \mathbf{A}\mathbf{x}_1 &= \mathbf{A}(\mathbf{A}\mathbf{x}_0) \\ &= \begin{bmatrix} \frac{1}{3} & \frac{1}{2} & \frac{1}{4} \\ \frac{2}{3} & \frac{1}{4} & \frac{1}{2} \\ 0 & \frac{1}{4} & \frac{1}{4} \end{bmatrix} \begin{bmatrix} 13/168 \\ 17/168 \\ 23/168 \end{bmatrix} = \begin{bmatrix} 43/168 \\ 191/168 \\ 23/168 \end{bmatrix}. \end{aligned}$$

$$\begin{aligned} \text{⑥ } \mathbf{A}\mathbf{x}_0 &= \mathbf{x}_0, \quad \mathbf{x}_0 = \begin{bmatrix} q \\ b \\ c \end{bmatrix}, \\ \text{⑦ } q+b+c &= 1 \\ \begin{bmatrix} \frac{1}{3} & \frac{1}{2} & \frac{1}{4} \\ \frac{2}{3} & \frac{1}{4} & \frac{1}{2} \\ 0 & \frac{1}{4} & \frac{1}{4} \end{bmatrix} \begin{bmatrix} q \\ b \\ c \end{bmatrix} &= \begin{bmatrix} q \\ b \\ c \end{bmatrix}, \\ \frac{1}{3}q + \frac{1}{2}b + \frac{1}{4}c &= q \quad \text{⑧} \\ \frac{2}{3}q + \frac{1}{4}b + \frac{1}{2}c &= b \quad \text{⑨} \\ \frac{1}{4}b + \frac{1}{4}c &= c \quad \text{⑩} \end{aligned}$$

$$\begin{aligned} \text{⑪ } q \text{ ④} &\Rightarrow b+c=4c \\ &\Rightarrow b-3c=0 \quad \text{⑪} \\ &\quad b=3c \end{aligned}$$

$$\begin{aligned} \text{⑫ } q \text{ ⑤} &\Rightarrow -8q+6c+3c=0 \\ &\Rightarrow -8q+9c=0 \quad \text{⑫} \\ &\Rightarrow q=\frac{9c}{8} \quad \text{⑬} \end{aligned}$$

$$\begin{aligned} \text{⑭ } q \text{ ⑩} &\Rightarrow \frac{21}{8}c + 3c + c = 1 \\ &\Rightarrow c = \frac{8}{21} \quad \text{⑭} \end{aligned}$$

$$\begin{aligned} \text{⑮ } q \text{ ⑧} &\Rightarrow q = \frac{21}{8} \times \frac{8}{21} = q \\ &\Rightarrow q = \frac{21}{21} = q \quad \text{⑯} \end{aligned}$$

$$\begin{aligned} \text{⑰ } q \text{ ⑪} &\Rightarrow b = 3 \cdot \frac{8}{21} = \frac{24}{21} = b \\ &\Rightarrow b = \frac{24}{21} \quad \text{⑰} \end{aligned}$$

$$\begin{aligned} \text{⑱ } q \text{ ⑭} &\Rightarrow \mathbf{x}_0 = \begin{bmatrix} q \\ b \\ c \end{bmatrix} = \begin{bmatrix} \frac{21}{21} \\ \frac{24}{21} \\ \frac{8}{21} \end{bmatrix} \quad \text{⑲} \end{aligned}$$

$$\text{⑲ } \mathbf{x}_1 = \mathbf{A}\mathbf{x}_0$$

$$\begin{aligned} \text{⑳ } q \text{ ⑩} &\Rightarrow 4q+6b+3c=12q \\ &\Rightarrow -8q+6b+3c=0 \quad \text{⑳} \end{aligned}$$

$$\mathbf{x}_1 = \begin{bmatrix} 0.4 & 0 & 0.4 \\ 0 & 0.5 & 0.4 \\ 0.6 & 0.5 & 0.2 \end{bmatrix}$$

$$\mathbf{x}_1 = \begin{bmatrix} \frac{4}{3} \\ \frac{2}{3} \\ \frac{1}{3} \end{bmatrix}$$

$$\begin{aligned} \text{㉑ } q \text{ ③} &\Rightarrow \frac{8q+3b+6c}{12} = b \\ &\Rightarrow 8q+3b+6c=12b \\ &\Rightarrow 8q+9b+6c=0 \quad \text{㉑} \end{aligned}$$

After 2 years

$$A\mathbf{x}_1 = A(A\mathbf{x}_0)$$

$$= \begin{bmatrix} 0.4 & 0 & 0.4 \\ 0 & 0.5 & 0.4 \\ 0.6 & 0.5 & 0.2 \end{bmatrix} \begin{bmatrix} 415 \\ 310 \\ 1350 \end{bmatrix}$$

$$= \begin{bmatrix} 84/500 \\ 97/500 \\ 119/500 \end{bmatrix}.$$

(b)

$$A\mathbf{x}_0 = \mathbf{x}_0 \quad \mathbf{x}_0 = \begin{bmatrix} 9 \\ 2 \\ 1 \end{bmatrix}$$

$$\text{Ansatz } A + b + c = 1 \quad \text{①}$$

$$\begin{bmatrix} 0.4 & 0 & 0.4 \\ 0 & 0.5 & 0.4 \\ 0.6 & 0.5 & 0.2 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

S.V.S ab ② 20.


$$\mathbf{x}_0 = \begin{bmatrix} 9 \\ 2 \\ 1 \end{bmatrix} = \begin{bmatrix} 10/500 \\ 11/500 \\ 1/500 \end{bmatrix}.$$

In exercises 1 through 8, sketch \mathbf{u} and its image under the given transformation.

$f: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by

$$f\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}; \quad \mathbf{u} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}.$$

$$\text{Given } \mathbf{u} = \begin{bmatrix} 2 \\ 3 \end{bmatrix} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$$




$$\text{Now } f(\mathbf{u}) = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} 2 \\ 3 \end{bmatrix} = \begin{bmatrix} 2 \\ -3 \end{bmatrix} = \begin{bmatrix} 2 \\ -3 \end{bmatrix}$$


$f: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ (reflection w.r.t. y-axis) defined by

$$f\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}, \quad \mathbf{u} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}.$$

$$\text{Given } \mathbf{u} = \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}.$$



$$\text{Now } f(\mathbf{u}) = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} -1 \\ 2 \end{bmatrix}.$$




Q3 $f: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a counter-clockwise rotation through 30° , w.r.t.

$$f(u) = \begin{bmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix},$$

$$\begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} 3 \\ 2 \end{bmatrix} = \begin{bmatrix} -3 \\ -2 \end{bmatrix}.$$

$$= \begin{bmatrix} \cos 30^\circ & -\sin 30^\circ \\ \sin 30^\circ & \cos 30^\circ \end{bmatrix} \begin{bmatrix} -1 \\ 3 \end{bmatrix}.$$

$$= \begin{bmatrix} \cos 30^\circ - 3 \sin 30^\circ \\ -\sin 30^\circ + 3 \cos 30^\circ \end{bmatrix} = \begin{bmatrix} -\frac{\sqrt{3}}{2} - \frac{3}{2} \\ -\frac{1}{2} + \frac{3\sqrt{3}}{2} \end{bmatrix} = \begin{bmatrix} -2.366 \\ 2.098 \end{bmatrix}.$$




Q4 $f: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a counter-clockwise rotation through $2\pi/3$

$$\text{various } u = \begin{bmatrix} -2 \\ -3 \end{bmatrix}$$

$$f(u) = \begin{bmatrix} \cos 2\pi/3 & -\sin 2\pi/3 \\ \sin 2\pi/3 & \cos 2\pi/3 \end{bmatrix} \begin{bmatrix} -2 \\ -3 \end{bmatrix}.$$

sketch

$$= \begin{bmatrix} -2 \cos 2\pi/3 + 3 \sin 2\pi/3 \\ -2 \sin 2\pi/3 - 3 \cos 2\pi/3 \end{bmatrix} = \begin{bmatrix} 3.6 \\ -0.23 \end{bmatrix}$$



Q5

θ_6 is similarly to θ_2 .


$f: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ defined by

$$f\left(\begin{bmatrix} x \\ y \\ z \end{bmatrix}\right) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix}, \quad u = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix}.$$


$$\text{given } u = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

sketch

$$u = (2, 1, 3)$$



$$\text{Now } f(u) \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \\ 0 \end{bmatrix} = \begin{bmatrix} 2 \\ 2 \\ 0 \end{bmatrix}.$$



Q8
f: $\mathbb{R}^3 \rightarrow \mathbb{R}^3$ defined by


$$f\left(\begin{bmatrix} x \\ y \\ z \end{bmatrix}\right) = \begin{bmatrix} 1 & 0 & 1 \\ -1 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix}; \quad u = \begin{bmatrix} 0 \\ -2 \\ 4 \end{bmatrix}$$

$$f(u) = \begin{bmatrix} 1 & 0 & 1 \\ -1 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 0 \\ -2 \\ 4 \end{bmatrix} = \begin{bmatrix} 4 \\ -2 \\ 4 \end{bmatrix}$$

$$\begin{bmatrix} 7 \\ 3 \end{bmatrix} = \begin{bmatrix} 7 \\ 3 \end{bmatrix}$$

Line $W = \begin{bmatrix} 7 \\ 3 \end{bmatrix}$ is in range of f

Q10 + Q11 is similarly to Q9



Q9
f: $\mathbb{R}^2 \rightarrow \mathbb{R}^3$.

Given that $f(u) = Ax + A = \begin{bmatrix} 1 & 3 \\ -1 & 2 \end{bmatrix}x + \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$ and $W = \begin{bmatrix} 7 \\ 3 \end{bmatrix}$.

Now $Ax = f(x) = W$

$$\begin{bmatrix} 1 & 3 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 7 \\ 3 \end{bmatrix} \quad \textcircled{*}$$

$$\begin{bmatrix} x+3y \\ -x+2y \end{bmatrix} = \begin{bmatrix} 7 \\ 3 \end{bmatrix}$$

$$\begin{aligned} x+3y &= 7 & (i) \\ -x+2y &= 3 & (ii) \end{aligned}$$

$$\text{Solving } (i) \text{ and } (ii) \text{ we get } \boxed{x=1} + \boxed{y=2}$$

$$q(iii) = \boxed{z=-1} \quad q(iii) =, x=2+1=3 = \boxed{x=3}$$

$$\begin{bmatrix} 1 & 3 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 7 \\ 3 \end{bmatrix}$$

Q9
Given that $f(u) = Ax + A = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}x + \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ and $W = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$.

Now $Ax = f(x) = W$, $x = \begin{bmatrix} x \\ y \end{bmatrix}$.

$$\begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix} \quad \textcircled{*}$$

$$\begin{aligned} 1 &= x+2y & (i) \\ -1 &= y & (ii) \\ 2 &= x+y & (iii) \end{aligned}$$

$$\alpha_A(x) = \gamma$$

$$\begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 2 & -2 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$$

$$\lim_{N \rightarrow \infty} N^{-\frac{1}{2}} \sin \left[\frac{\pi}{N} \right] = 0$$

$$\text{Q13} \quad \text{Given } w = \begin{bmatrix} 1 \\ 1 \end{bmatrix} + x \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$$\text{Now } Ax = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

$$\text{Q13} \quad \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$$

$$\text{Now } Ax = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$$

and λ


Mrs. Jameel Nasli

Lecturer U.E.T Peshawar

Q5

(a) $A = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}$ & $u = \begin{bmatrix} x \\ y \end{bmatrix}$


$$Au = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -x \\ y \end{bmatrix} = \begin{bmatrix} -x \\ y \end{bmatrix}$$



Reflection about
y-axis.

(b) $A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$, $u = \begin{bmatrix} x \\ y \end{bmatrix}$


$$Au = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0-y \\ x+0 \end{bmatrix} = \begin{bmatrix} -y \\ x \end{bmatrix}$$



Rotation counter clockwise through $\pi/2$.

(c) $A = \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix}$, $u = \begin{bmatrix} x \\ y \end{bmatrix}$.


$$Au = \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -y \\ -x \end{bmatrix}.$$



Reflection about line
 $y = x$.

(d) $A = \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix}$, $u = \begin{bmatrix} x \\ y \end{bmatrix}$.

$$Au = \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -y \\ -x \end{bmatrix}.$$



Reflection about line
 $y = -x$.

Q7 (a) $A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$, $u = \begin{bmatrix} x \\ y \end{bmatrix}$ $Au = \begin{bmatrix} x \\ 0 \end{bmatrix}$.

Projection onto x-axis.

(b) $A = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$, $u = \begin{bmatrix} x \\ y \end{bmatrix}$ Then $Au = \begin{bmatrix} 0 \\ y \end{bmatrix}$.

Projection onto y-axis.

Q8(a) $A = \begin{bmatrix} 1 & 2 & 0 \\ 0 & 1 & -1 \end{bmatrix}$, $\mathbf{w} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$.

~~check~~ $A\mathbf{x} = \mathbf{w}$

$$\begin{bmatrix} 1 & 2 & 0 \\ 0 & 1 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

$$x+2y=0$$

$$y-z=-1$$

$$(2)$$

$$x=-2y \quad (1)$$

$$y+z=0$$

Let $z=y$ be real no.

$$\text{Let } z=0 \text{ then } \boxed{y=-1} + \boxed{y=2}$$

$$\text{If } z=1, \quad y=0, \quad x=0$$

$$\text{So } \mathbf{u} = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \quad \mathbf{v} = \begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix}.$$

⑥ $A = \begin{bmatrix} 2 & 1 & 0 \\ 0 & 2 & -1 \end{bmatrix}$, $\mathbf{w} = \begin{bmatrix} 9 \\ 4 \end{bmatrix}$

$$A\mathbf{x} = \mathbf{w}$$

$$\begin{bmatrix} 2 & 1 & 0 \\ 0 & 2 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 9 \\ 4 \end{bmatrix} \Rightarrow 2x+y=9 \Rightarrow \boxed{x=4-\frac{y}{2}}$$

$$2y-2=4 \Rightarrow \boxed{y=\frac{4+2}{2}}$$

$$\text{If } z=4, \quad y=4, \quad x=0$$

$$\text{If } z=0, \quad y=2, \quad x=1$$

$$\text{So } \mathbf{u} = \begin{bmatrix} 0 \\ 4 \\ 4 \end{bmatrix}, \quad \mathbf{v} = \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}$$

Q1 Reduced row echelon form.
Q2 Neither Q3 Reduced row echelon form

Q4 Neither Q5 Row echelon form Q6 Neither

Q7 Neither Q8 Neither

Q9 $A = \begin{bmatrix} 1 & 0 & 3 \\ 5 & -1 & 4 \\ 5 & 2 & 5 \\ -3 & 1 & 2 \end{bmatrix}$



a) Interchanging the second and fourth rows.

$$A = \begin{bmatrix} 1 & 0 & 3 \\ 5 & -1 & 4 \\ 5 & 2 & 5 \\ -3 & 1 & 2 \end{bmatrix}$$

b) Multiplying the third row by 3.

$$A = \begin{bmatrix} 1 & 0 & 3 \\ -3 & 1 & 4 \\ 15 & -3 & 12 \\ -3 & 1 & 2 \end{bmatrix}$$

(c) Adding (-3) times the 1st row to the 4th row

R.W
-3 0 -9 (-3)

$$\text{Q10} \quad \begin{bmatrix} 1 & 0 & 3 \\ -3 & 1 & 4 \\ 4 & 2 & 2 \\ 2 & -1 & -4 \end{bmatrix}$$

$$A = \begin{bmatrix} 1 & 0 & 3 \\ -3 & 1 & 4 \\ 4 & 2 & 2 \\ 2 & -1 & -4 \end{bmatrix}$$

$$\text{Q10} \quad \begin{bmatrix} 1 & 0 & 3 \\ -3 & 1 & 4 \\ 4 & 2 & 2 \\ 2 & -1 & -4 \end{bmatrix}$$

Q10 is similar to Q9.

Q11 Find three matrices that are row equivalent to

$$A = \begin{bmatrix} 2 & -1 & 3 & 4 \\ 0 & 1 & 2 & -1 \\ 5 & 2 & -3 & 4 \end{bmatrix}$$

1st Possibility

$$(a) \quad \begin{bmatrix} 2 & -1 & 3 & 4 \\ 5 & 2 & -3 & 4 \\ 0 & 1 & 2 & -1 \end{bmatrix} \xrightarrow{\text{R}_2 \leftrightarrow \text{R}_3}$$

3rd Possibility

$$(b) \quad \begin{bmatrix} 2 & -1 & 3 & 4 \\ 0 & 1 & 2 & -1 \\ 7 & 1 & 0 & 8 \end{bmatrix} \xrightarrow{\text{R}_1 + \text{R}_2}$$

$$(b) \quad \begin{bmatrix} 4 & -2 & 6 & 8 \\ 5 & 1 & 2 & -1 \\ 2 & -3 & 4 \end{bmatrix} \xrightarrow{\text{R}_1 + \text{R}_2}$$

In Exercise 13 through 16, find a row echelon form of the given matrix.

$$\begin{bmatrix} 0 & 0 & -1 & 3 & -1 \\ 3 & 0 & 1 & 2 & 4 \\ 2 & 1 & 4 & 2 & 3 \\ 4 & -1 & 2 & 1 & 3 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & -1 & 3 & -1 \\ 0 & 1 & 2 & 4 & 2 \\ 0 & -1 & 2 & 1 & 3 \\ 0 & 4 & -1 & 2 & 1 \end{bmatrix}$$

$\xrightarrow{\text{R}_1 + \text{R}_2}$ Row of dashes

$$\begin{bmatrix} 1 & 0 & -1 & 3 & -1 \\ 0 & 1 & 2 & 4 & 2 \\ 0 & -1 & 2 & 1 & 3 \\ 0 & 4 & -1 & 2 & 1 \end{bmatrix} \xrightarrow{\text{R}_2 + 2\text{R}_1}$$

$$\begin{bmatrix} 1 & 0 & -1 & 3 & -1 \\ 0 & 1 & 2 & 4 & 2 \\ 0 & -1 & 2 & 1 & 3 \\ 0 & 4 & -1 & 2 & 1 \end{bmatrix} \xrightarrow{\text{R}_4 - 3\text{R}_1}$$

$$\begin{bmatrix} 1 & 0 & -1 & 3 & -1 \\ 0 & 1 & 2 & 4 & 2 \\ 0 & -1 & 2 & 1 & 3 \\ 0 & 7 & 7 & -5 & 1 \end{bmatrix} \xrightarrow{\text{R}_{23}}$$

$$\begin{bmatrix} 1 & 0 & -1 & 3 & -1 \\ 0 & 1 & 2 & 4 & 2 \\ 0 & -1 & 2 & 1 & 3 \\ 0 & 7 & 7 & -5 & 1 \end{bmatrix} \xrightarrow{\text{R}_4 - 3\text{R}_1}$$

$$\begin{bmatrix} 1 & 0 & -1 & 3 & -1 \\ 0 & 1 & 2 & 4 & 2 \\ 0 & -1 & 2 & 1 & 3 \\ 0 & 7 & 7 & -5 & 1 \end{bmatrix} \xrightarrow{\text{R}_4 - 3\text{R}_1}$$

$$\begin{bmatrix} 1 & 0 & -1 & 3 & -1 \\ 0 & 1 & 2 & 4 & 2 \\ 0 & -1 & 2 & 1 & 3 \\ 0 & 7 & 7 & -5 & 1 \end{bmatrix} \xrightarrow{\text{R}_4 - 3\text{R}_1}$$

OK

$$\left[\begin{array}{cccc} 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & -1 \\ 0 & 0 & 2 & 1 \\ 0 & 0 & 0 & -2 \end{array} \right] \left[\begin{array}{c} 1 \\ -1 \\ 2 \\ -2 \end{array} \right] \left[\begin{array}{c} R_1 \\ R_2 \\ R_3 \\ R_4 \end{array} \right]$$

$$\left[\begin{array}{cccc} 1 & -1 & 2 & -2 \\ 0 & 0 & 1 & 2 \\ 0 & 2 & 1 & 0 \\ -2 & 1 & 0 & 1 \end{array} \right] \left[\begin{array}{c} 1 \\ -1 \\ 2 \\ -2 \end{array} \right]$$

$$\left[\begin{array}{cccc} 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 3 \\ 0 & -2 & 1 & 2 \\ -1 & 3 & 2 & 7 \end{array} \right] \left[\begin{array}{c} 1 \\ -1 \\ 2 \\ -2 \end{array} \right] \left[\begin{array}{c} R_1 \\ R_2 \\ R_3 \\ R_4 \end{array} \right]$$

$$\left[\begin{array}{cccc} 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 3 \\ 0 & 1 & 2 & 1 \\ 1 & 2 & 1 & 0 \end{array} \right] \left[\begin{array}{c} 1 \\ -1 \\ 2 \\ -2 \end{array} \right] \left[\begin{array}{c} R_1 \\ R_2 \\ R_3 \\ R_4 \end{array} \right]$$

$$\left[\begin{array}{cccc} 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 2 \\ 0 & -1 & 0 & 0 \\ 0 & -2 & 1 & 2 \end{array} \right] \left[\begin{array}{c} 1 \\ -1 \\ 2 \\ -2 \\ 1 \end{array} \right] \left[\begin{array}{c} R_1 \\ R_2 \\ R_3 \\ R_4 \\ R_5 \end{array} \right]$$

$$\left[\begin{array}{cccc} 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 2 \\ 0 & -1 & 0 & 0 \\ 0 & -2 & 1 & 2 \end{array} \right] \left[\begin{array}{c} 1 \\ -1 \\ 2 \\ -2 \\ 1 \end{array} \right] \left[\begin{array}{c} R_1 \\ R_2 \\ R_3 \\ R_4 \\ R_5 \end{array} \right]$$

$$\left[\begin{array}{cccc} 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 2 \\ -1 & 2 & 1 & 7 \end{array} \right] \left[\begin{array}{c} R_1 \\ R_2 \\ R_3 \\ R_4 \\ R_5 \end{array} \right]$$

$$\left[\begin{array}{cccc} 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 2 \\ -1 & 2 & 1 & 7 \end{array} \right] \left[\begin{array}{c} R_1 \\ R_2 \\ R_3 \\ R_4 \\ R_5 \end{array} \right]$$

$$\left[\begin{array}{cccc} 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 2 \\ -1 & 2 & 1 & 7 \end{array} \right] \left[\begin{array}{c} R_1 \\ R_2 \\ R_3 \\ R_4 \\ R_5 \end{array} \right]$$

Ex: 1.6

30

Q8 Let $A = \begin{bmatrix} 1 & 2 & 1 \\ -1 & 1 & 2 \\ 2 & 1 & -2 \end{bmatrix}$ & $Ax = b$.

$$(a) X = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}; b = 0$$

JAN

$$\text{Now } Ax = b$$

$$\begin{bmatrix} 1 & 2 & 1 \\ -1 & 1 & 2 \\ 2 & 1 & -2 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

PHOTOSTAT

$$\Rightarrow \begin{pmatrix} 1+4+3 \\ -1+2+6 \\ 2+1-6 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

$$\Rightarrow \begin{bmatrix} 8 \\ 7 \\ -2 \end{bmatrix} \neq \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \text{ No Solution.}$$

ADDRESS:

JAN PHOTOSTAT, COFFEE SHOP MARKET,
UNIVERSITY OF PESHAWAR,

PROPRIETOR:

QAISAR SHABAB, ARIF JAN

$$Ax = b$$

$$(b) X = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}; b = 0$$

$$\begin{bmatrix} 1 & 2 & 1 \\ -1 & 1 & 2 \\ 2 & 1 & -2 \end{bmatrix} \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 0+0+0 \\ 0+0+0 \\ 0+0+0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \Rightarrow \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Solution exist. / 0c + 0c
Similarly.

Phone No: 091-5610987

Mobile No: 0300-5829015, 0345-9165401
Arif Jan: 0345-9289401

In extends to through 22, find all solutions to R.

Linear system

$$\text{Ques a) } x+y+2z = -1$$

$$x-2y+2z = -5$$

$$3x+y+2z = 3$$

$$q^{(iii)} \Rightarrow z = \frac{26}{-13} \Rightarrow \boxed{z = -2}$$

$$q^{(ii)} \Rightarrow y - 4(-2) = 10$$

$$y = 10 - 8$$

$$\boxed{y = 2}$$

$$q^{(i)} \Rightarrow x + (-2) + 2(-2) = -1$$

$$x + 2 - 4 - 1 \Rightarrow \boxed{x = 1}$$

$\theta_b, \theta_c, \theta_d$ similarly

$$\text{Ques c) } x+y+2z+3w = 13$$

$$x-2y+2z+w = 8$$

$$3x+y+2z-w = 1$$

$$\left[\begin{array}{cccc|c} 1 & 1 & 2 & 1 & -1 \\ 1 & -2 & 1 & 1 & -5 \\ 0 & 3 & 1 & -1 & 4 \\ 0 & -2 & -5 & 1 & 6 \end{array} \right] \xrightarrow{\text{R}_2+R_1} \left[\begin{array}{cccc|c} 1 & 1 & 2 & 1 & -1 \\ 0 & -3 & 1 & -4 & -5 \\ 0 & -2 & -5 & 1 & 6 \\ 0 & 3 & 1 & -1 & 4 \end{array} \right] \xrightarrow{\text{R}_3+3\text{R}_1} \left[\begin{array}{cccc|c} 1 & 1 & 2 & 1 & -1 \\ 0 & -3 & 1 & -4 & -5 \\ 0 & -2 & -5 & 1 & 6 \\ 0 & 0 & -14 & 2 & 13 \end{array} \right] \xrightarrow{\text{R}_4+14\text{R}_3} \left[\begin{array}{cccc|c} 1 & 1 & 2 & 1 & -1 \\ 0 & -3 & 1 & -4 & -5 \\ 0 & -2 & -5 & 1 & 6 \\ 0 & 0 & 0 & 1 & 26 \end{array} \right]$$

$$\xrightarrow{\text{R}_2+R_1} \left[\begin{array}{cccc|c} 1 & 1 & 2 & 1 & -1 \\ 0 & -2 & 1 & -5 & -5 \\ 0 & -2 & -5 & 1 & 6 \\ 0 & 0 & -14 & 2 & 13 \end{array} \right]$$

$$\xrightarrow{\text{R}_3+2\text{R}_2} \left[\begin{array}{cccc|c} 1 & 1 & 2 & 1 & -1 \\ 0 & -2 & 1 & -5 & -5 \\ 0 & 0 & -1 & 1 & 1 \\ 0 & 0 & -1 & 1 & 26 \end{array} \right]$$

$$\xrightarrow{\text{R}_4+\text{R}_3} \left[\begin{array}{cccc|c} 1 & 1 & 2 & 1 & -1 \\ 0 & -2 & 1 & -5 & -5 \\ 0 & 0 & -1 & 1 & 1 \\ 0 & 0 & 0 & 2 & 27 \end{array} \right]$$

(6)

Ex 1.6

33

$$\left[\begin{array}{ccc|c} 1 & 1 & 2 & 13 \\ 0 & -3 & -1 & -5 \\ 0 & -2 & -5 & -10 \\ 0 & -2 & -5 & -10 \end{array} \right] \xrightarrow{\substack{R_1-R_3 \\ R_3-3R_1}} \left[\begin{array}{ccc|c} 1 & 1 & 2 & 13 \\ 0 & 3 & 1 & 5 \\ 0 & -2 & -5 & -10 \\ 0 & -2 & -5 & -10 \end{array} \right]$$

$$\left[\begin{array}{ccc|c} 1 & 1 & 2 & 13 \\ 0 & 3 & 1 & 5 \\ 0 & -2 & -5 & -10 \\ 0 & -2 & -5 & -10 \end{array} \right] \xrightarrow{(-1)R_2} \left[\begin{array}{ccc|c} 1 & 1 & 2 & 13 \\ 0 & -3 & -1 & -5 \\ 0 & -2 & -5 & -10 \\ 0 & -2 & -5 & -10 \end{array} \right]$$

$$\left[\begin{array}{ccc|c} 1 & 1 & 2 & 13 \\ 0 & -3 & -1 & -5 \\ 0 & -2 & -5 & -10 \\ 0 & -2 & -5 & -10 \end{array} \right] \xrightarrow{R_3+R_2} \left[\begin{array}{ccc|c} 1 & 1 & 2 & 13 \\ 0 & -3 & -1 & -5 \\ 0 & -2 & -5 & -10 \\ 0 & -2 & -5 & -10 \end{array} \right]$$

$$\left[\begin{array}{ccc|c} 1 & 1 & 2 & 13 \\ 0 & -3 & -1 & -5 \\ 0 & -2 & -5 & -10 \\ 0 & -2 & -5 & -10 \end{array} \right] \xrightarrow{R_2+R_3} \left[\begin{array}{ccc|c} 1 & 1 & 2 & 13 \\ 0 & 0 & -4 & -33 \\ 0 & -2 & -5 & -10 \\ 0 & -2 & -5 & -10 \end{array} \right]$$

$$\boxed{\omega w = y}$$

$$\sqrt{(ii)} \Rightarrow -13z - 26y = 64$$

$$-13(z+wy) = -164$$

$$\boxed{\frac{z+wy=8}{z=8-wy}}$$

$$\begin{aligned} \sqrt{(i')} &\Rightarrow y - 4(8-wy) - 8y = -33 \\ y - 32 + 8wy - 8y &= -33 \\ y = -33 + 32 &= -1 \Rightarrow \boxed{y = -1} \end{aligned}$$

$$\begin{aligned} \sqrt{(i)} &\Rightarrow x + (-1) + 2(8-wy) + 3y = 13 \\ x - 1 + 16 - 4y + 3y &= 13 \end{aligned}$$

$$x + 15 - y = 13$$

$$x = 13 - 15 + y$$

$$\boxed{x = y - 2}$$

$\boxed{\text{Qb, Qd, similarly}}$

$\underbrace{\quad}_{0}$

$$\text{Qb}(C) \left[\begin{array}{ccccc} 2 & 1 & 1 & -2 & 1 \\ 3 & -2 & 1 & -6 & 1 \\ 1 & -1 & -1 & -1 & -1 \\ 6 & 0 & 1 & -9 & 1 \\ 5 & -1 & 2 & -8 & 1 \end{array} \right]$$

$$\left[\begin{array}{ccccc} 2 & 1 & 1 & -2 & 1 \\ 3 & -2 & 1 & -6 & 1 \\ 1 & -1 & -1 & -1 & -1 \\ 6 & 0 & 1 & -9 & 1 \\ 5 & -1 & 2 & -8 & 1 \end{array} \right]$$

$$\left[\begin{array}{ccccc} 1 & -1 & -1 & -1 & -1 \\ 3 & -2 & 1 & -6 & 1 \\ 1 & -1 & -1 & -1 & -1 \\ 6 & 0 & 1 & -9 & 1 \\ 5 & -1 & 2 & -8 & 1 \end{array} \right] \xrightarrow{R_3}$$

$$\begin{bmatrix} 1 & -1 & -1 & -1 \\ 0 & 1 & 4 & -3 \\ 0 & -6 & 3 & 0 \\ 0 & 7 & -3 & -3 \\ \hline 0 & -6 & 7 & -3 \\ 0 & -6 & 7 & -3 \\ \hline R_2 - 3R_1 & R_3 - 2R_1 & R_4 - 6R_1 & R_5 - 5R_1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -1 & -1 & -1 \\ 0 & 0 & -5 & -1 \\ 0 & -6 & 4 & -3 \\ 0 & 7 & -3 & 0 \\ \hline 0 & -6 & 7 & -3 \\ 0 & -6 & 7 & -3 \\ \hline -R_3 & R_{32} \end{bmatrix}$$

$$\begin{bmatrix} 1 & -1 & -1 & -1 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & -1 \\ \hline 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & -1 \\ \hline R_3 + SR_2 & R_4 + 6R_2 & R_5 + 6R_2 \end{bmatrix}$$

$$\sim \begin{bmatrix} 1 & -1 & -1 & | & 2 \\ 0 & 1 & 0 & | & 1 \\ 0 & 0 & \alpha^2 & | & \alpha \\ \hline 0 & 0 & \alpha^2 & | & 2 \\ 0 & 0 & \alpha^2 & | & 3 \\ \hline R_2 - R_1 & R_3 - R_1 \end{bmatrix}$$

a) If $\alpha = -2$ no solution

(b) If $\alpha \neq \pm 2$ then unique solution

(c) If $\alpha = 2$ infinite many soln.

$$\begin{bmatrix} 1 & -1 & -1 & | & 2 \\ 0 & 0 & 0 & | & 1 \\ 0 & 0 & 0 & | & 1 \\ 0 & 0 & 0 & | & 1 \\ \hline 0 & 0 & 0 & | & 1 \\ 0 & 0 & 0 & | & 1 \\ \hline -\frac{1}{10}(R_3) \end{bmatrix}$$

$$\begin{bmatrix} 1 & -1 & -1 & | & 2 \\ 0 & 0 & 0 & | & 1 \\ 0 & 0 & 0 & | & 1 \\ 0 & 0 & 0 & | & 1 \\ \hline 0 & 0 & 0 & | & 1 \\ 0 & 0 & 0 & | & 1 \\ \hline R_4 + R_3 & R_5 + 6R_3 \end{bmatrix}$$

No solution.

$$\begin{array}{l} x+y-z=2 \\ x+2y+z=3 \\ x+y+(x^2-5)z=a \end{array}$$

$$\text{If } \alpha^2 - 4 = 0 \Rightarrow \alpha^2 = 4 \Rightarrow (\alpha = \pm 2)$$

$$\text{If } \alpha^2 - 4 = \alpha^2 - 2^2 = 0 = -4 \text{ not soln}$$

$$\text{If } \alpha = 2 \quad 4 - 4 = 2 - 2 = 0 = 0 \text{ Infnt sol.}$$

If $\alpha \neq \pm 2$ unk soln.

$$\text{Q24: } \left[\begin{array}{ccc|c} 1 & -1 & 1 & 2 \\ 2 & 3 & 2 & 5 \\ 2 & 3 & (a^2-1) & a^4+1 \end{array} \right]$$

$$\sim \left[\begin{array}{ccc|c} 1 & 1 & 2 & R_2-2R_1 \\ 0 & 1 & 1 & R_3-2R_1 \\ 0 & 1 & a^2-3 & a-3 \end{array} \right]$$

$$\sim \left[\begin{array}{ccc|c} 1 & 1 & 2 & \\ 0 & 1 & 1 & \\ 0 & 0 & a^2-3 & a-4 \end{array} \right]$$

- a) $\alpha = \pm\sqrt{3}$ (b) $\alpha \neq \pm\sqrt{3}$, (c) None.

Comparing (ii) & (iii) we get

$$\left[\begin{array}{ccc|c} 4 & 1 & 3 & 4 \\ 2 & -1 & 3 & 5 \\ 2 & 2 & 0 & 1 \end{array} \right] \xrightarrow{\text{R}_1 \leftrightarrow \text{R}_2} \left[\begin{array}{ccc|c} 4 & 1 & 3 & 4 \\ 2 & 1 & 3 & 5 \\ 2 & 2 & 0 & 1 \end{array} \right]$$

$$\left[\begin{array}{ccc|c} 4x+y+3z & 4 \\ 2x-y+3z & 5 \\ 2x+2y & 1 \end{array} \right] = \left[\begin{array}{c} 4 \\ 5 \\ 1 \end{array} \right].$$

By augmented matrix

$$\left[\begin{array}{ccc|c} 4 & 1 & 3 & 4 \\ 2 & -1 & 3 & 5 \\ 2 & 2 & 0 & 1 \end{array} \right]$$

$$\sim \left[\begin{array}{ccc|c} 1 & 1 & 2 & R_2-R_1 \\ 0 & 0 & 1 & R_3-R_1 \\ 0 & 0 & a^2-6 & a-2 \end{array} \right]$$

- a) $\alpha = \pm\sqrt{6}$, (b) $\alpha \neq \pm\sqrt{6}$ (c) None.

$$\text{Q25: } \left[\begin{array}{ccc|c} 1 & 1 & 3 & 1 \\ 1 & a^2-8 & a & 3 \\ 1 & a^2-8 & a & 3 \end{array} \right] \sim \left[\begin{array}{ccc|c} 1 & 1 & 3 & 1 \\ 0 & a^2-9 & a-3 & 3 \\ 0 & a^2-9 & a-3 & 3 \end{array} \right] \xrightarrow{\text{R}_2-\text{R}_1}$$

- (a) $\alpha = -3$, (b) $\alpha \neq \pm 3$, (c) $\alpha = 3$.

$$\text{Q26: } \left[\begin{array}{ccc|c} -1 & -1 & 0 & -1/2 \\ 2 & -1 & 3 & 4 \\ -1 & 3 & -1 & 5 \end{array} \right] \xrightarrow{\text{R}_3 \leftrightarrow \text{R}_1} \left[\begin{array}{ccc|c} 4 & -1 & 3 & 4 \\ 2 & -1 & 3 & 5 \\ 2 & 2 & 0 & -1 \end{array} \right] \xrightarrow{\text{R}_3 \leftrightarrow \text{R}_2} \left[\begin{array}{ccc|c} 4 & -1 & 3 & 4 \\ 2 & -1 & 3 & 5 \\ 2 & 2 & 0 & -1 \end{array} \right]$$

$$\text{Q27 to Q30 similarly to Q20}$$

Ex 16

$$\sim \left[\begin{array}{ccc|cc} 1 & -1 & 0 & -1 & -\frac{1}{2} \\ 0 & -3 & 3 & 1 & \frac{6}{6} \\ 0 & -3 & 3 & 1 & 6 \end{array} \right] R_2 - 2R_1$$

$$\sim \left[\begin{array}{ccc|cc} 1 & -1 & 0 & -1 & -\frac{1}{2} \\ 0 & 1 & -1 & -2 & \frac{1}{2} \\ 0 & -3 & 3 & 1 & 6 \end{array} \right]$$

$$\sim \left[\begin{array}{ccc|cc} 1 & -1 & 0 & -1 & -\frac{1}{2} \\ 0 & 1 & -1 & -2 & \frac{1}{2} \\ 0 & 0 & 0 & 1 & 0 \end{array} \right] - R_3$$

$$\sim \left[\begin{array}{ccc|cc} 1 & -1 & 0 & -1 & -\frac{1}{2} \\ 0 & 1 & -1 & -2 & \frac{1}{2} \\ 0 & 0 & 0 & 1 & 0 \end{array} \right] R_3 + 3R_2$$

$$\text{Now } x+y = -\frac{1}{2} \quad (i)$$

$$y-z = -2 \quad (ii)$$

$$\text{Let } z=t$$

$$y(t) \Rightarrow y = -2+t$$

$$x(t) \Rightarrow x + (-2+t) = -\frac{1}{2}$$

$$x = -\frac{1}{2} + t$$

$$\boxed{x = \frac{3}{2}t - t}$$

$$\text{M.V. } \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} \frac{3}{2}t - t \\ -2+t \\ t \end{pmatrix}$$

$$\text{Lehrbuch U.E.T Pohlhausen}$$

$$\text{Soln:- } \left[\begin{array}{ccc|cc} 4 & -1 & 3 & a \\ 2 & -1 & 3 & b \\ 2 & -2 & 0 & c \end{array} \right] \left[\begin{array}{c} x \\ y \\ z \end{array} \right] = \left[\begin{array}{c} a \\ b \\ c \end{array} \right]$$

A.M

$$\left[\begin{array}{ccc|cc} 4x+y+3z & a \\ 2x-y+3z & b \\ 2x+2y & c \end{array} \right] = \left[\begin{array}{c} a \\ b \\ c \end{array} \right] = \left[\begin{array}{c} 4 & -1 & 3 & a \\ 2 & -1 & 3 & b \\ 2 & -2 & 0 & c \end{array} \right]$$

Augmented

$$\sim \left[\begin{array}{ccc|cc} 2 & 2 & 0 & 1 & a \\ 2 & -1 & 3 & b \\ 4 & -1 & 3 & c \end{array} \right] R_3$$

$$\sim \left[\begin{array}{ccc|cc} 1 & -1 & 0 & \frac{1}{2} & a \\ 2 & -1 & 3 & b \\ 4 & -1 & 3 & c \end{array} \right] (\frac{1}{2})R_1$$

$$\sim \left[\begin{array}{ccc|cc} 1 & -1 & 0 & \frac{1}{2} & a \\ 0 & -3 & 3 & b & a-2c \\ 0 & -3 & 3 & b & a-2c \end{array} \right] R_2 - 2R_1$$

$$\sim \left[\begin{array}{ccc|cc} 1 & -1 & 0 & \frac{1}{2} & a \\ 0 & 1 & -1 & \frac{1}{2}(b-a) & a-2c \\ 0 & -3 & 3 & b & a-2c \end{array} \right] - R_3$$

$$\sim \left[\begin{array}{ccc|cc} 1 & -1 & 0 & \frac{1}{2} & a \\ 0 & 1 & -1 & \frac{1}{2}(b-a) & a-2c \\ 0 & 0 & 0 & a-c-b & a-2c \end{array} \right] R_3 + 3R_2$$

Solution is possible if

$$\cancel{\text{if } a-c-b=0}$$

$$\boxed{-a+c+b=0}$$

Q34 is similarly to Q33

$$\text{Q35} \\ A = \begin{bmatrix} 1 & -1 \\ 2 & 3 \end{bmatrix}, b_1 = \begin{bmatrix} 1 \\ -8 \end{bmatrix}, b_2 = \begin{bmatrix} 5 \\ -5 \end{bmatrix}$$

$$Ax = b_1$$

$$\begin{bmatrix} 1 & -1 \\ 2 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1 \\ -8 \end{bmatrix}$$

$$\begin{bmatrix} x-y \\ 2x+3y \end{bmatrix} = \begin{bmatrix} 1 \\ -8 \end{bmatrix}$$

New Augmented matrix

$$\begin{bmatrix} 1 & -1 & 1 \\ 2 & 3 & -8 \end{bmatrix} \Rightarrow \sim \begin{bmatrix} 1 & -1 & 1 \\ 0 & 5 & -10 \end{bmatrix} R_2 - 2R_1$$

$$\therefore 5y = -10 \Rightarrow y = -2$$

$$x - y = 1 \Rightarrow x = -2 + 1 = -1$$

$$\boxed{x = -1, y = -2}$$

2nd part
Ax = b₂

$$\begin{bmatrix} 1 & -1 & 1 \\ 2 & 3 & -8 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 5 \\ -5 \end{bmatrix}, \Rightarrow \begin{bmatrix} x-y \\ 2x+3y \end{bmatrix} = \begin{bmatrix} 5 \\ -5 \end{bmatrix}$$

A.M

$$\begin{bmatrix} 1 & -1 & 5 \\ 2 & 3 & -5 \end{bmatrix}$$

3rd part $[A : b_1, b_2]$

$$\begin{bmatrix} 1 & -1 & 1 & 5 \\ 0 & 5 & -10 & -15 \end{bmatrix}$$

$$\sim \begin{bmatrix} 1 & -1 & 1 & 5 \\ 0 & 5 & -10 & -15 \end{bmatrix} R_2 - 2R_1$$

Same results

Q36 is similarly to Q35.

$$\sim \begin{bmatrix} 1 & -1 & 5 \\ 0 & 5 & -15 \end{bmatrix} R_2 - 2R_1$$

$$5y = -15 \Rightarrow y = -3$$

$$x - y = 5 \Rightarrow x = 5 - (-3) = 8$$

$$\boxed{x = 8, y = -3}$$

$$\underline{\text{Q37}} \quad A = \begin{bmatrix} 1 & 0 & 5 \\ 0 & 1 & -4 \\ 0 & 0 & 1 \end{bmatrix} \quad (-4T_3 - A)x = 0$$

$$T_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} -1 & 5 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix} R_2 + SR_1$$

$$-4T_3 = \begin{bmatrix} -1 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & -4 \end{bmatrix}$$

$$-4T_3 - A = \begin{bmatrix} -4 & 0 & 0 \\ 0 & -4 & 0 \\ 0 & 0 & -4 \end{bmatrix} - \begin{bmatrix} 1 & 0 & 5 \\ 0 & 1 & -4 \\ 0 & 1 & -4 \end{bmatrix} = \begin{bmatrix} -5 & 0 & 0 \\ 0 & -5 & 0 \\ 0 & 1 & 0 \end{bmatrix}$$

$$N.D.W. (-4T_3 - A)x = 0$$

$$\Rightarrow \begin{bmatrix} -5 & 0 & 0 \\ 0 & -5 & 0 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$x+5y+z=0 \quad (i) \\ y=0 \quad (ii)$$

Let $x=y$ (real no.)

$$\sqrt{(i)} \Rightarrow x+y+z=0 \Rightarrow [x=-y]$$

$$\begin{bmatrix} -5x+0y-5z \\ -x-5y-1z \\ 0x-1y+0z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

A.M.

$$\begin{bmatrix} -5 & 0 & -5 \\ 0 & -5 & -1 \\ 0 & -1 & 0 \end{bmatrix}$$

$$\sim \begin{bmatrix} 1 & 5 & -1 \\ 0 & 0 & 5 \\ 0 & 1 & 0 \end{bmatrix} R_{2,1}$$

$$\begin{bmatrix} 1 & 5 & -1 \\ 0 & 0 & 5 \\ 0 & 1 & 0 \end{bmatrix} R_{2,1}$$



1039

Augmented matrix

$$\begin{array}{r} 1 \\ 2 \\ 3 \\ \hline -6 \\ \hline 6 \end{array}$$

$$\left[\begin{array}{ccc|c} 1 & 2 & -3 & a \\ 0 & 1 & q & b-2a \\ 0 & 0 & 1 & c-3a \end{array} \right] \xrightarrow{\text{R}_2-2\text{R}_1} \left[\begin{array}{ccc|c} 1 & 2 & -3 & a \\ 0 & 1 & q & b-2a \\ 0 & 0 & 1 & c-3a \end{array} \right] \xrightarrow{\text{R}_3-5\text{R}_1}$$

$$3q - b + c = 0$$

θ_{40} is similarly to θ_{39}

A vertical line on the left, with three curly braces on the right side, each containing three small circles.

Jan Laser Photostate

Matric. to M.A.
Notes are available.

Qasir Shabbab
0300 5829015

Aamir 0345 2939401
Ph: 091-5610987

Qasir Shabbab
0300 5829015

Aamir 0345 9239401
Ph: 091-5610984

$$\begin{cases} 0 \\ 0 \\ -1 \\ 0 \end{cases} \left[\begin{matrix} 0 & 0 \end{matrix} \right] R_2 - 2R_1$$

$$\overline{00}$$

$$D_2 + K_2 = \begin{cases} 0 & x=0 \\ 0 & x>0 \end{cases}$$

$$\int_0^{\infty} \frac{1}{x^2} \left(\int_0^x u dy \right) dx = \int_0^{\infty} 0 du = 0$$

$$\begin{bmatrix} 4 & -4 & 0 & -1 \\ 0 & 0 & 4 & -2 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\left(\begin{bmatrix} 4 & 0 \\ 0 & 4 \end{bmatrix} - \begin{bmatrix} 4 & 1 \\ 0 & 2 \end{bmatrix} \right) \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}.$$

$$\text{Now } \mathcal{L}X - AX = 0$$

$$(4T_2 - A)X = 0$$

$$\underline{Q41} \quad A = \begin{bmatrix} 4 & 1 \\ 0 & 2 \end{bmatrix}, \quad X = \begin{bmatrix} x \\ y \end{bmatrix}$$

$$A \mathbf{x} = \begin{bmatrix} 4 & 1 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = 4 \begin{bmatrix} x \\ y \end{bmatrix} = 4\mathbf{x}$$

A.M

Matric to M.A.
Notes are available



Matric to M.A.
Notes are available



$$y=0, \text{ for } x \in X \text{ (rest of)} \\ x = [y] = [0].$$

ایران و اسلام

Shop #153 Coffee shop 153

Scanned by CamScanner

$$(3\bar{I}_3 - A)x = 0$$

$$\text{Q4.3} \quad A = \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix} \quad d(3\bar{I}_3 - A)x = 0$$

$$\text{Now } 3x - Ax = 0$$

$$(3\bar{I}_3 - A)x = 0$$

$$\left(3 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} - \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix} \right) \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\left[\begin{bmatrix} 3 & 0 \\ 0 & 3 \end{bmatrix} - \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix} \right] \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 3-2 & 0-1 \\ 0-1 & 3-2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \Rightarrow \begin{bmatrix} x-y \\ -x+y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

A.M

$$\begin{bmatrix} 1 & -1 & 0 \\ -1 & 1 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 2 & -2 & 1 \\ -1 & 3 & -1 \\ -4 & 4 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -3 & 1 & 0 \\ 0 & 2 & -2 & 1 \\ -4 & 4 & -2 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

A.M

$$\begin{bmatrix} 2 & -2 & 1 \\ -1 & 3 & -1 \\ -4 & 4 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 2 & -2 & 1 \\ -1 & 3 & -1 \\ -4 & 4 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 3 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 3 \end{bmatrix} \begin{bmatrix} 1 & 2 & -1 \\ 1 & 0 & 1 \\ 4 & -4 & 5 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 2 & -2 & 1 \\ -1 & 3 & -1 \\ -4 & 4 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\sim \begin{bmatrix} 1 & -1 & 0 \\ 0 & 0 & 0 \end{bmatrix} R_{21} + R_1$$

$$\text{Q4.4} \quad x \neq y = 0 \quad \text{but} \quad x = y$$

$$\sim \begin{bmatrix} 1 & -3 & 1 & 0 \\ 0 & 1 & -4 & 1 \\ 0 & 0 & 2 & 0 \end{bmatrix} R_2 \leftrightarrow R_1$$

$$y = y$$

$$X = \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} y \\ y \end{bmatrix} \quad \text{d.}$$

$$\sim \left[\begin{array}{ccc|c} 1 & -3 & 1 & 0 \\ 0 & 1 & -\frac{1}{4} & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] R_3 + 8R_2$$

$$\lambda - 3y + z = 0 \quad (i)$$

$$y - \frac{1}{4}z = 0 \quad (ii)$$

$$\text{Let } \boxed{z = y} \text{ (real no only)}$$

$$\Rightarrow \boxed{y = y}$$

$$\text{Eqn} \Rightarrow \lambda - \frac{3}{4}y + y = 0$$

$$\lambda - \frac{3y+4z}{4} = 0 \Rightarrow \lambda + \frac{y}{4} = 0 \Rightarrow \boxed{\lambda = -\frac{y}{4}}$$

$$\text{From: } \boxed{\lambda = -\frac{y}{4}, \quad y = \frac{1}{4}\lambda, \quad z = \lambda}$$

Q44 is similarly to Q43

$$\sim \left[\begin{array}{cccc|c} 1 & 2 & -1 & -2 & 2 \\ 0 & -3 & 0 & 7 & -2 \\ 0 & 0 & 4 & 6 & 3 \\ 0 & -3 & 0 & 7 & -2 \end{array} \right] \xrightarrow[R_2+2R_1]{R_3-R_1, R_4-4R_1}$$

$$\sim \left[\begin{array}{cccc|c} 1 & 2 & -1 & -2 & 2 \\ 0 & 1 & 0 & -7/3 & 2/3 \\ 0 & 0 & 4 & 6 & 3 \\ 0 & -3 & 0 & 7 & -2 \end{array} \right] \xrightarrow[R_4-4R_2]{R_3-R_2}$$

Mr. Jannal Nasir

Lecturer UET Peshawar.

$$\begin{aligned} \lambda + 2y - z - 2w &= 2 \quad (i) \\ y - \frac{7}{3}w &= \frac{2}{3} \quad (ii) \\ 4z + 6w &= 3 \quad (iii) \end{aligned}$$

$$\text{Let } w = \lambda \text{ (any real no.)}$$

Q45 Given $X = X_p + X_h$,
where X_p is particular solution
& X_h is a solution to the associated homogeneous system.
A.M

$$\left[\begin{array}{cccc|c} 1 & 2 & -1 & -2 & 2 \\ 2 & 1 & -2 & 3 & 1 \\ 4 & 2 & 3 & 4 & 5 \\ 5 & -4 & 1 & 6 \end{array} \right]$$

$$\sqrt{(\text{iii})} \Rightarrow 4\bar{x} + \delta y = 3$$

$$4\bar{x} = 3 - \delta y \\ \bar{x} = \frac{3}{4} - \frac{\delta}{4}y \Rightarrow \boxed{\bar{x} = \frac{3}{4} - \frac{3}{2}y}$$

$$\sqrt{(\text{ii})} \Rightarrow \boxed{y = \frac{3}{2}x + \frac{7}{3}}$$

$$\sqrt{(\text{i})} \Rightarrow y + 2(\frac{3}{2}x + \frac{7}{3}) - (\frac{3}{4}x - \frac{3}{2}y) - 2y = 2$$

$$\stackrel{(3,3)}{\Rightarrow} 3 = 9a_2 + 3a_1 + a_0 \quad \text{--- (i)}$$

$$x = \frac{17}{12}y - \frac{5}{12}$$

$$x = \frac{17}{12}y - \frac{25}{6}$$

$$x = x_p + x_{f_k}$$

$$\begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} = \begin{bmatrix} 17a_2 \\ -25a_1 \\ 7a_2 \\ -3a_1 \end{bmatrix} + \begin{bmatrix} 7/3 \\ 1 \\ 0 \\ y \end{bmatrix} \quad \text{Ans.}$$

(1,2) Point

$$y \otimes \Rightarrow y = a_2 + a_1 + a_0 \quad \text{--- (ii)}$$

Since quadratic polynomials $a_2x^2 + a_1x + a_0 = y$ \Rightarrow $\boxed{y = a_2x^2 + a_1x + a_0}$ --- (iii)

$$(1,2)(3,3)(5,8)$$

Q_{12} is similar to Q_{13}

$$\begin{bmatrix} 1 & 1 & 1 & 2 \\ 9 & 3 & 1 & 3 \\ 25 & 5 & 1 & 8 \end{bmatrix}$$

in A.M

$$\stackrel{\text{F. P. of } (5,3)}{\Rightarrow} 8 = 25a_2 + 5a_1 + a_0 \quad \text{--- (iv)}$$

in A.M

$$\begin{bmatrix} 1 & 1 & 1 & 2 \\ 0 & -6 & -8 & -15 \\ 0 & -20 & -24 & -42 \end{bmatrix}$$

Q48, Q49, Q50 similarly to Q47

$$\sim \left[\begin{array}{ccc|c} 1 & 1 & 1 & 2 \\ 0 & 1 & 4/3 & 4/2 \\ 0 & -20 & -24 & -42 \end{array} \right] - \frac{1}{6} R_2$$

$$10x_1 + 12x_2 + 15x_3 = 16 \times 60 = 960$$

$$6x_1 + 8x_2 + 12x_3 = 11 \times 60 = 660$$

$$12x_1 + 12x_2 + 18x_3 = 18 \times 60 = 1080$$

Aug:M

$$\left[\begin{array}{ccc|c} 10 & 12 & 15 & 960 \\ 6 & 8 & 12 & 660 \\ 12 & 12 & 18 & 1080 \end{array} \right]$$

S.Y.S.

$$a_2 + a_1 + a_0 = 2 \quad (a)$$

$$a_1 + 4/3 a_0 = 5/2 \quad (b)$$

$$8/3 a_0 = 8 \quad (c)$$

$$a_1 = 8 \times 3/8 = \boxed{a_0 = 3}$$

$$a_1 = a_0 + 4/3 a_0 = 5/2$$

$$a_1 = 5/2 - 4 = \frac{5-8}{2} = -3/2$$

$$\boxed{a_1 = -3/2}$$

$$a_1 = a_0 + (-3/2) + 3 = 2$$

$$a_1 = 2 - 3 + 3/2 = \frac{4-6+3}{2} = 1/2$$

$$\boxed{a_2 = 1/2}$$

$$\sqrt{4} \Rightarrow \boxed{y = \sqrt{x^2 - 3/2 x + 3}}$$

Ans.

Qasir Shahab
0300 5829015

Aarif: 0345 9289401
Ph: 091-5610987

Matrix to M.A.
Notes are available



Jall Laser Photostate

Shop #58 coffee shop bazaar Islamia college Peshawar University

$$\begin{array}{l} \text{Q3} \\ \begin{aligned} & X_1 + 2X_2 + 3X_3 = 6 \times 60 = 360 \\ & X_1 + 4X_2 + 5X_3 = 11 \times 60 = 660 \\ & 2X_1 + 4X_2 + 5X_3 = 2 \times 60 = 120 \end{aligned} \end{array}$$

Ans: Matrix

$$\begin{bmatrix} 1 & 2 & 3 & | & 360 \\ 2 & 4 & 5 & | & 660 \end{bmatrix}$$

$$\sim \left[\begin{array}{cccc} 1 & 2 & 3 & | & 360 \\ 0 & 0 & -1 & | & -60 \end{array} \right] R_2 - 2R_1$$

$$X_3 = 60 \quad (\text{deluxe binding})$$

$$X_1 + 2X_2 + 3X_3 = 360 \quad \text{--- C}$$

$$\text{Let } X_2 = y \quad (\text{any real no.})$$

$$X_1 + 2y + 3(60) = 360$$

$$\begin{aligned} \text{Q4} \Rightarrow P(m) &= am^2 + b \\ P'(0) &= 2a(0) + b \Rightarrow P'(0) = b. \end{aligned}$$

$$\text{Q5} \Rightarrow f'(m) = 2e^{2m}$$

$$f'(0) = 2e^0 = 2$$

$$\text{But Given } f'(0) = P'(0)$$

$$\boxed{X_1 = 180 - 2y, \quad X_2 = y, \quad X_3 = 60}$$

$$\text{Similarly } \boxed{a = 2}$$

$$\boxed{2 = b}$$

$$\text{Given } P(m) = am^2 + bm + c \quad \text{--- Q5}$$

$$P(0) = f(0), \quad P'(0) = f'(0), \quad P''(0) = f''(0) \quad \text{And } f(m) = e^{2m}$$

$$f(m) = e^{2m} - c$$

$$f(0) = e^0 = 1 \Rightarrow \boxed{f(0) = 1}$$

$$\sqrt{a} \Rightarrow P(0) = a0 + b0 + c$$

$$\boxed{P(0) = c}$$

$$\text{But Given } \boxed{P(0) = f(0)}$$

$$\boxed{c = 1.}$$

Now Aug. Matrix

$$\left[\begin{array}{cccc|c} 4 & -1 & -1 & 0 & 80 \\ -1 & 4 & 0 & -1 & 80 \\ -1 & 0 & 4 & -1 & 80 \\ 0 & -1 & -1 & 4 & 80 \end{array} \right] S.V.S$$

θS_4 is similarly do. θS_3

0 0 0 0

$$T_1 = \frac{30 + 50 + T_2 + T_3}{4}$$

$$\angle T_1 = 80^\circ + T_2 + T_3$$

$$\angle T_1 - T_2 - T_3 = 80^\circ \quad (i)$$

$$T_2 = \frac{30 + 50 + T_1 + T_4}{4}$$



$$\angle T_2 - T_1 - T_4 = 80^\circ \quad (ii)$$

$$T_3 = \frac{T_1 + T_4 + 50^\circ + 0^\circ}{4}$$

$$\angle T_3 - T_1 - T_4 = 50^\circ \quad (iii)$$

$$T_4 = \frac{T_3 + T_2 + 50^\circ + 0}{4}$$

$$\angle T_4 - T_3 - T_2 = 50^\circ \quad (iv)$$

 Jan Laser Photostate Matrix to M.A. Notes are available 	 Jan Laser Photostate Matrix to M.A. Notes are available 
--	--

Qasir Shahab
0300 5829015

Aarif: 0345 9289401
Ph: 091-5610987

Qasir Shahab
0300 5829015

Aarif: 0345 9289401
Ph: 091-5610987

Shop 55 coffee shop bazar Islamia college Peshawar university

BIT Matrices (optional) P-17

Table 5

$$\begin{array}{|c|c|c|} \hline + & 0 & 1 \\ \hline 0 & 0 & 0 \\ \hline 1 & 0 & 0 \\ \hline \end{array}$$

$$\begin{array}{|c|c|c|} \hline \times & 0 & 1 \\ \hline 0 & 0 & 0 \\ \hline 0 & 0 & 0 \\ \hline \end{array}$$

$$\begin{array}{|c|c|c|} \hline - & 1 & 0 & 1 & 0 \\ \hline 0 & 0 & 1 & 0 & 1 \\ \hline 0 & 1 & 1 & 0 & 1 \\ \hline \end{array} R_2, R_3$$

Aug: M

(S.P.ca)

2x16

46

Table

$$\text{Exp} \quad A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$

$$A+B = \begin{bmatrix} 1+1 & 0+1 \\ 0+0 & 1+0 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$

$$\begin{array}{l} x+y+z=0 \quad \text{(i)} \\ y+z=1 \quad \text{(ii)} \\ x=0 \end{array}$$

$$x=-y-z$$

$$\text{Exp} \Rightarrow y=1-x \quad \left[\text{Exp(i)} \quad x=-y \right]$$

$$x=-(1-x)$$

$$y=1-x$$

$$x=1$$

$$y=0$$

$$z=1$$

$$y=1-1=0$$

$$x=-(1-1)$$

$$x=0$$

Note:- A bit is also known as binary digit or Boolean value.

A bit matrix is also called a Boolean matrix.

$$A-B = A + (-B) = A + B$$

$$A-B = A+B$$

Note:- We know zero and one's. Thus the addition of two bits will give us all the additive inverse of 1's & 0's. Now to complete the study of bit matrices $A+B$ we proceed as follows:

$$\begin{aligned} A+B &= A + (\text{inverse of } B) = A + \bar{B} = A+B \\ A-B &= A + B \end{aligned}$$

(b)

$$\begin{bmatrix} 1 & 1 & 0 & 0 & 0 \\ 1 & 1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 1 & 0 \end{bmatrix}$$

$$\sim \begin{bmatrix} 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 & 0 \end{bmatrix} R_2 + R_1$$

$$\sim \begin{bmatrix} 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 & 0 \end{bmatrix} R_3 + R_1$$

$$x+y=0 \Rightarrow x=-y$$

$$x=1 \quad \left\{ \begin{array}{l} y=0 \\ y=1 \end{array} \right.$$

$$x=1 \quad \left\{ \begin{array}{l} y=0 \\ y=1 \end{array} \right.$$

$$\sim \begin{bmatrix} 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 & 0 \end{bmatrix}$$

$\sim \begin{bmatrix} 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 & 0 \end{bmatrix}$

MN. Jammal Nasrin

Lecturer U.E.T Peshawar

$$\text{(a)} \quad Ax = C$$

$$\begin{bmatrix} -1 & 1 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ -1 \end{bmatrix}$$

$$\begin{bmatrix} x+y+z+w \\ x-y+z+w \\ x+y+z-w \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ -1 \end{bmatrix}$$

Aug: M

$$\begin{bmatrix} -1 & 1 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 1 & 0 & 0 \end{bmatrix}$$

$$\sim \begin{bmatrix} 1 & -1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 \end{bmatrix} R_3 + R_1$$

$$x+y=0 \Leftrightarrow x=-y \Rightarrow x=1$$

$$x=0 \quad \left\{ \begin{array}{l} y=0 \\ y=1 \end{array} \right.$$

$$\begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$\sim \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$

Q1 Show that $\begin{bmatrix} 2 & 1 \\ -2 & 3 \end{bmatrix}$ is nonsingular.

Sol. Let $A = \begin{bmatrix} 2 & 1 \\ -2 & 3 \end{bmatrix}$ To find \bar{A}^1 , we let $\bar{A}^1 = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. $\text{---} \textcircled{1}$

Then we must have

$$A\bar{A}^1 = \begin{bmatrix} 2 & 1 \\ -2 & 3 \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} = T_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

JAN

PHOTOSTAT

So $\text{---} \textcircled{1}$

$$\begin{bmatrix} 2a+c & 2b+d \\ -2a+3c & -2b+3d \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\begin{aligned} 2a+c &= 1 \text{ ---} \textcircled{1} & -2a+3c &= 0 \text{ ---} \textcircled{2} \\ 2b+d &= 0 \text{ ---} \textcircled{3} & -2b+3d &= 1 \text{ ---} \textcircled{4} \end{aligned}$$

$\textcircled{1} + \textcircled{2}$

$$\{c = 1 \Rightarrow \boxed{c = 1/4}$$

$$\text{---} \textcircled{3} + \textcircled{4} \Rightarrow 2a = 1 - \frac{1}{4} = \frac{3}{4} \Rightarrow \boxed{a = \frac{3}{8}}$$

$$\{d = 1 \Rightarrow \boxed{d = 1/4}$$

$$\text{---} \textcircled{3} + \textcircled{4} \Rightarrow 2b = -\frac{1}{4} \Rightarrow \boxed{b = -\frac{1}{8}}$$

Phone No: 091-5610987

Mobile No: 0300-5829015, 0345-9165902
Arif Jan: 0345-9229401

$\therefore \sqrt{\text{---} \textcircled{1}} \Rightarrow \bar{A}^1 = \begin{bmatrix} 3/8 & -1/8 \\ 1/4 & 1/4 \end{bmatrix}$ We conclude \bar{A}^1 is exist
so A is nonsingular.

Q_3 is similar to $Q_2 + Q_1$

$$A = \begin{bmatrix} 1 & a & -1 \\ 3 & 2 & 3 \\ 2 & 2 & 1 \end{bmatrix} \quad A' = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$$

Let $A = \begin{bmatrix} 2 & 1 \\ -4 & -2 \end{bmatrix}$, let $A' = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$.

Now $A A' = I_2$

$$\begin{bmatrix} 2 & 1 \\ -4 & -2 \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

$$= \begin{bmatrix} 2a+c & 2b+d \\ -4a+2c & -4b+2d \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

$$2a+c=1 \quad (1)$$

$$-4a+2c=0 \quad (2)$$

$$2b+d=0 \quad (3)$$

$$-4b+2d=1 \quad (4)$$

④ by ③ adding eqv ④

$$4a+2c=2$$

$$-4a+2c=0$$

$$\underline{0=2}$$
 impossible

A' doesn't exist so it is singular

Now $A A' = I_3$

$$\begin{bmatrix} 1 & 2 & -1 \\ 3 & 2 & 3 \\ 2 & 2 & 1 \end{bmatrix} \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

$$= \begin{bmatrix} a+2d-g & b+2e-h & c+2f-i \\ 3a+2d+g & 3b+2e+h & 3c+2f+i \\ 2a+2d+g & 2b+2e+h & 2c+2f+i \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

$$a+2d-g=1 \quad (1)$$

$$b+2e-h=0 \quad (4)$$

$$c+2f-i=0 \quad (7)$$

$$3a+2d+g=0 \quad (2)$$

$$3b+2e+h=1 \quad (5)$$

$$3c+2f+i=0 \quad (8)$$

$$2a+2d+g=0 \quad (3)$$

$$2b+2e+h=0 \quad (6)$$

$$2c+2f+i=1 \quad (9)$$

$$\begin{array}{l} (1) - (2) \\ a+2d-g=1 \\ 2a+2d+g=0 \\ \hline 0=2 \text{ impossible} \end{array} \quad \begin{array}{l} (1) - (3) \\ a+2d-g=1 \\ 2a+2d+g=0 \\ \hline -a-2g=1 \end{array}$$

$$\begin{array}{l} (2) \text{ by } (3) \text{ then sub: from } 10 \\ -2a-4g=2 \\ -2a-4g=1 \\ \hline 0=1 \text{ impossible. } A \text{ doesn't exist so} \end{array}$$

$$\begin{array}{l} -2a-4g=2 \\ -2a-4g=1 \\ \hline 0=1 \text{ impossible. } A \text{ doesn't exist so} \end{array}$$

Q5(a)

$$\text{Q5(a)} \quad A = \begin{bmatrix} 1 & 3 \\ -2 & 6 \end{bmatrix} \quad \text{Let } \tilde{A} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}. \quad \text{--- (P)}$$

$$\text{Now } A\tilde{A}^{-1} = \begin{bmatrix} 1 & 3 \\ -2 & 6 \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

$$\begin{bmatrix} a-2b & 3a+6b \\ c-2d & 3c+6d \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

$$a-2b=1 \quad \text{--- (1)} \quad 3a+6b=0 \quad \text{--- (B)}$$

$$c-2d=0 \quad \text{--- (2)} \quad 3c+6d=1 \quad \text{--- (4)}$$

Solving the above equations we get

$$\boxed{a = 1/4} \quad \boxed{b = -1/4} \quad \boxed{c = 1/4} \quad \boxed{d = 1/2}$$

$$\text{Ex (P)} \Rightarrow \tilde{A} = \begin{bmatrix} 1/4 & -1/4 \\ 1/4 & 1/2 \end{bmatrix}.$$

$$\text{Q(b)} \quad A = \begin{bmatrix} 1 & 2 & 3 \\ 1 & 1 & 2 \\ 0 & 1 & 2 \end{bmatrix}.$$

We now compute the reduced row echelon form of the matrix. To find \tilde{A} , we proceed as follows:

$$\left[\begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 1 & 1 & 2 & 0 & 1 & 0 \\ 0 & 1 & 2 & 0 & 0 & 1 \end{array} \right]$$

$$\xrightarrow{\text{R}_1 \leftrightarrow \text{R}_3} \left[\begin{array}{ccc|ccc} 0 & 1 & 2 & 0 & 0 & 1 \\ 1 & 2 & 3 & 1 & 0 & 0 \\ 1 & 1 & 2 & 0 & 1 & 0 \end{array} \right] \xrightarrow{\text{R}_2 - \text{R}_1} \left[\begin{array}{ccc|ccc} 0 & 1 & 2 & 0 & 0 & 1 \\ 1 & 1 & 1 & 1 & 0 & 0 \\ 1 & 1 & 2 & 0 & 1 & 0 \end{array} \right] \xrightarrow{\text{R}_3 - \text{R}_2} \left[\begin{array}{ccc|ccc} 0 & 1 & 2 & 0 & 0 & 1 \\ 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & -1 & 1 & 0 \end{array} \right]$$

Qasim Shahab
0300 5829015
Aarif 0345 9289401
Ph: 091-5610987

Jail Laser Photostate

Matrix to M.A.
Notes are available

$$\xrightarrow{\text{R}_1 - \text{R}_2} \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & -1 & 0 & 0 \\ 0 & 1 & 1 & 1 & -1 & 0 \\ 0 & 0 & 1 & 1 & 1 & -1 \end{array} \right] \xrightarrow{\text{R}_3 - \text{R}_2} \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & -1 & 0 & 0 \\ 0 & 1 & 1 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 \end{array} \right] \xrightarrow{\text{R}_1 - \text{R}_2} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 & -1 \end{array} \right]$$

Ex 1.7

5

$$\left[\begin{array}{cccc} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & -2 \\ 1 & 2 & 1 & 1 \end{array} \right] R_1 - R_3 \\ R_2 - R_3$$

$$8 \\ A^{-1} = \left[\begin{array}{ccc} 0 & -1 & -1 \\ -2 & 1 & 1 \\ -1 & 1 & 1 \end{array} \right]$$

$$\left[\begin{array}{ccc} -2 & -3 & 0 \\ 0 & -7 & 0 \\ 0 & 0 & 0 \end{array} \right] R_2 - 2R_1 \\ R_3 + 3R_1$$

$$\left[\begin{array}{ccc} -1 & -2 & -3 \\ 0 & 1 & -1 \\ 0 & -7 & 0 \end{array} \right] 5R_2$$

$\sim \sim \sim \sim \sim \sim$

$D_6 - b = D_10 \cdot R$ similarly do D_5

All which of the following linear system have a non-homogeneous solution?

$$(b) \quad 2x + y - z = 0$$

$$x - 2y - 3z = 0$$

$$-3x - y + 2z = 0$$

Method 1
A.M

$$\left[\begin{array}{ccc|c} 2 & 1 & -1 & 0 \\ -1 & -2 & -3 & 0 \\ -3 & 1 & 2 & 0 \\ 2 & 0 & 0 & 0 \end{array} \right] R_2$$

$$\left[\begin{array}{ccc|c} 0 & -1 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] R_1$$

$$\left[\begin{array}{ccc|c} 0 & 0 & -1 & 0 \\ 0 & 0 & -2 & 0 \\ 0 & 0 & -2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] R_2 - R_1$$

$$\left[\begin{array}{ccc|c} 1 & 1/2 & -1/2 & 0 \\ 1 & -2 & -3 & 0 \\ -3 & 1 & 2 & 0 \\ 0 & 0 & 0 & 1 \end{array} \right] \frac{1}{2}R_1$$

Method 2
 $[A : I_3]$

$$A : I_3 \Rightarrow \left[\begin{array}{ccc|c} 2 & 1 & -1 & 1 \\ -1 & -2 & -3 & 0 \\ -3 & 1 & 2 & 0 \end{array} \right] R_3 + 3R_1$$

$$\left[\begin{array}{ccc|c} 2 & 1 & -1 & 1 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{array} \right] R_2 - R_1$$

An infinite solution exists so it is non-trivial.

$$\begin{array}{r} 2 \\ \hline 0 - - \\ 0 - 0.5 - 0.5 \\ 0 - - \\ \hline 0.5 & 0.5 & 0 \\ 0 & 0.4 & 0 \\ - & 0 & 0 \\ \hline R_3 - 0.5 R_2 \end{array}$$

So no inverse exists since its non-invertible.

Ω_{12} is similarly to Ω_{11}

?

$$A = \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix} \text{ und } A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \quad \text{---}$$

$$S_{\text{inv}} = \Gamma A - I_2$$

$$\begin{bmatrix} 2a+3c \\ a+4c \end{bmatrix} = \begin{bmatrix} 2b+3d \\ b+4d \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$\begin{array}{l} 2a+3c=1 \quad (1) \\ a+4c=0 \quad (2) \\ b+4d=-1 \quad (3) \end{array}$$

Solving the above equation we get

$$\boxed{a = 4/5}, \boxed{b = -3/5}, \boxed{c = -1/5}, \boxed{d = 2/5}$$

$$\sqrt{A} = \begin{pmatrix} 4/5 & -3/5 \\ -1/5 & 2/5 \end{pmatrix}$$

δ_{14} is similarly to δ_{13}

$$\sin A \bar{A} = I_3$$

$$\begin{array}{r}
 \overline{0} \\
 -0 \\
 \overline{20} \\
 \overline{0} \\
 \overline{0}
 \end{array}$$

$$= \begin{pmatrix} b+e & c+f & d+g \\ b & c & d \\ b+2e+g & c+2f+g & d+2g+j \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}$$

$$\begin{array}{l}
 b+2e = 0 \quad (7) \\
 c+2f+ai = 0 \quad (8) \\
 d+2g+ji = 1 \quad (9)
 \end{array}$$

$$\begin{array}{l} \text{Q1} \Rightarrow 0+e=1 \\ \text{Q2} \Rightarrow 1+e=0 \end{array}$$

110

$$\text{Case } \Theta = 0 \Rightarrow 0 + 2 + k_0 = 0 \Rightarrow k_0 = -2$$

$$\sigma \circ \oplus = 1 + \sigma(-) + \sigma_{ii} \Rightarrow$$

$$\text{Ansatz: } \psi = \frac{1}{\sqrt{2}}(\psi_1 + i\psi_2)$$

$$\sqrt{\oplus} \Rightarrow \tilde{A} = \begin{bmatrix} 0 & 1 & 0 \\ 1 & -1 & 0 \\ -2/a & 1/a & 1/a \end{bmatrix} \text{ when } a \neq 0.$$

$$\text{Given } A = \begin{bmatrix} 1 & -1 & 0 \\ 0 & 0 & 2 \end{bmatrix}, \text{ let } \bar{A}' = \begin{bmatrix} b & c & d \\ e & f & g \end{bmatrix}$$

Q17
Let X be input matrix given and $A = A^T$

$$\text{find } A \quad \text{Given } A^{-1} = \begin{bmatrix} g & h & f \\ g & k & l \\ g & l & i \end{bmatrix} \quad A = \begin{bmatrix} 2 & -1 & 1 \\ 3 & 2 & 1 \\ 2 & 1 & 1 \end{bmatrix}$$

$$\text{Now } AA^{-1} = I_3$$

$$\begin{bmatrix} 2 & -1 & 3 \\ 3 & 2 & -1 \\ 2 & 1 & 1 \end{bmatrix} \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad (*)$$

$$\begin{bmatrix} 2a+d+3g & 2b+e+3f & 2c+f+i+3i \\ 3a+2d-g & 3b+2e-f & 3c+2f-i \\ 2a+d+g & 2b+e+f & 2c+f+i \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$-3 + \frac{1}{2} + 3g = 1 \Rightarrow g = \frac{1}{2}$$

Put value in ③

$$2a+d+2g = 1$$

$$2(-\frac{3}{2}) + (\frac{5}{2}) + 2g = 1$$

⑤ + ⑥

$$3b+2e-f = 1$$

$$2b+e+f = 0$$

$$2c+f+i = 1$$

④ + ⑤ add

$$\begin{aligned} 2a+2d-g &= 0 \\ 2a+d+g &= 0 \end{aligned}$$

$$5a+3d = 0 \quad (1)$$

$$\text{Add 3 times } ② \text{ to } ①$$

$$\begin{aligned} 2a+d+3g &= 1 \\ 9a+6d-3g &= 0 \end{aligned}$$

$$11a+7d = 3 \quad (2)$$

$$11a+7d = 1 \quad (3)$$

$$\text{Solving } (2)' \text{ and } (1)'$$

Solving (2)' and (3)'

$$5a = -3d \Rightarrow a = -\frac{3}{5}d$$

$$11(-\frac{3}{5}d) + 7d = 1$$

$$-33d + 35d = 5 \Rightarrow d = \frac{5}{2}$$

$$d = \frac{5}{2}$$

$$5b = 1 - 3e \Rightarrow b = \frac{1-3e}{5}$$

$$11\left(\frac{1-3e}{5}\right) + 7e = 3$$

$$\Rightarrow \boxed{e = 2}$$

$$b = \frac{1-3(2)}{5} = -5/5 = -1$$

$$\boxed{b = -1}$$

$$\text{or } \textcircled{1} \Rightarrow 2b + e + 3h = 0 \\ 2(-1) + 2 + 3h = 0$$

$$\Rightarrow \boxed{h = 0}$$

$$\textcircled{2} + \textcircled{1}$$

$$\begin{matrix} 3c + 2f - i = 0 \\ 2c + f + i = 1 \\ 5c + 2f = 1 \end{matrix} \quad \text{(2)}$$

Add (3) times \textcircled{2} to \textcircled{1}

$$\begin{matrix} 2c + f + 3i = 0 \\ 9c + 6f + 3i = 0 \end{matrix}$$

$$\frac{11c + 7f = 0}{\text{(1)}}$$

Solving \textcircled{2} & \textcircled{1}

$$11c = -7f \Rightarrow c = -\frac{7}{11}f$$

$$5(-\frac{7}{11}f) + 3f = 1 \Rightarrow f = -\frac{1}{12}$$

$$C = -\frac{7}{11}f \times -\frac{1}{2} = \frac{7}{22}$$

$$\boxed{C = \frac{7}{22}}$$

$$3c + 2f - i = 0$$

$$3(\frac{7}{22}) + 2(-\frac{1}{12}) - i = 0$$

$$\Rightarrow \boxed{i = -\frac{1}{12}}$$

$$\text{or } \textcircled{3} \Rightarrow A = \begin{bmatrix} -\frac{3}{2} & -1 & \frac{7}{2} \\ \frac{9}{2} & 2 & -\frac{11}{2} \\ \frac{1}{2} & 0 & -\frac{1}{2} \end{bmatrix}$$

$$X = A^{-1}b, \quad b = \begin{bmatrix} 3 \\ 0 \\ 10 \end{bmatrix}$$

$$X = \begin{bmatrix} -\frac{3}{2} & -1 & \frac{7}{2} \\ \frac{9}{2} & 2 & -\frac{11}{2} \\ \frac{1}{2} & 0 & -\frac{1}{2} \end{bmatrix} \begin{bmatrix} 3 \\ 0 \\ 10 \end{bmatrix}$$

$$X = \begin{bmatrix} -30 \\ 60 \\ 10 \end{bmatrix} A$$

$$X = \tilde{A}^{-1}b, \quad b = \begin{bmatrix} 1 \\ 2 \\ 14 \end{bmatrix}$$

$$X = \begin{bmatrix} -\frac{3}{2} & -1 & \frac{7}{2} \\ \frac{9}{2} & 2 & -\frac{11}{2} \\ \frac{1}{2} & 0 & -\frac{1}{2} \end{bmatrix} \begin{bmatrix} 8 \\ 14 \\ 14 \end{bmatrix} = \begin{bmatrix} -33 \\ -1 \\ -1 \end{bmatrix} +$$

$$= \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \begin{bmatrix} x & y \\ z & w \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

Ques

$$A = \begin{bmatrix} 1 & 3 \\ 2 & 7 \end{bmatrix}$$

(a) A^{-1}
 Now $A\bar{A} = I_2$, $\text{let } \bar{A} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ ————— \otimes

$$\Rightarrow \begin{bmatrix} 1 & 3 \\ 2 & 7 \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} a+3c & b+3d \\ 2a+7c & 2b+7d \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\begin{aligned} a+3c &= 1 & \text{--- (1)} \\ 2a+7c &= 0 & \text{--- (2)} \end{aligned}$$

Solving the above equations we get

$$\begin{bmatrix} a=1 \\ b=-3 \\ c=-2 \\ d=1 \end{bmatrix}$$

$$q_{(a)} \Rightarrow \bar{A} = \begin{bmatrix} 1 & -3 \\ 2 & 1 \end{bmatrix} \quad \text{--- (a)}$$

(b)

$$(A^T)^{-1} = ?$$

$$A^T = \begin{bmatrix} 1 & 2 \\ 3 & 7 \end{bmatrix} \quad \text{and } (A^T)^{-1} = \begin{bmatrix} x & y \\ z & w \end{bmatrix} \quad \text{--- (b)}$$

$$(A^T)(A^T)^{-1} = I_2$$


The $(A^T)^{-1}$ exists and it is defined.

$$\text{Given } A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + B = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

$$q_{(b)} \Rightarrow (A^T)^{-1} = \begin{bmatrix} 7 & -2 \\ -3 & 1 \end{bmatrix}, \text{ Ans.}$$

$$q_{(b)} \Rightarrow (A^T)^T = \begin{bmatrix} 7 & -2 \\ -3 & 1 \end{bmatrix}$$

$$8.0 (A^T)^T = (A^T)^T$$



$$= \begin{bmatrix} 1 & 2x \\ 3x & 7x \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\begin{aligned} 1+2x &= 1 & \text{--- (1)} \\ 3x+7x &= 0 & \text{--- (2)} \\ 10x &= 0 & \text{--- (3)} \\ x &= 0 & \text{--- (4)} \end{aligned}$$

Solving all the above equations we get

$$\begin{bmatrix} x=0 \\ y=-3 \\ z=-3 \\ w=1 \end{bmatrix}$$

⑥ Yes for A non-singular & $c \neq 0$

$$(cA)^{-1} = \frac{1}{c} A^{-1}$$

$$T_n = c(A(\frac{1}{c}A)^{-1})$$

$$T_n = c(\frac{1}{c}A) \cdot A A^{-1}$$

$$T_n = T_n$$

\Rightarrow

$$(\lambda - 1)x + 2y = 0 \quad (1)$$

$$2x + (\lambda - 1)y = 0 \quad (2)$$

$$\text{from } (1) \quad x = \frac{(1-\lambda)y}{2}$$

$$x^2 + (\lambda - 1)y^2 = 0$$

$$\Rightarrow \frac{(\lambda - 1)(1-\lambda)y^2}{4} + 2y^2 = 0$$

$$\Rightarrow -\frac{(\lambda - 1)(1-\lambda)y^2}{4} + 2y^2 = 0$$

$$\Rightarrow -\frac{(\lambda - 1)^2 y^2 + 4y^2}{4} = 0$$

$$\Rightarrow -((1-\lambda)^2 + 4)y^2 = 0$$

$$\Rightarrow -((1-\lambda)^2 + 4) = 0$$

$$\Rightarrow -(\lambda^2 - 2\lambda + 1 + 4) = 0$$

$$\Rightarrow \lambda^2 - 2\lambda - 3 = 0 \quad \lambda = \frac{2 \pm \sqrt{4+12}}{2} = \frac{2 \pm 4}{2}$$

$$\boxed{\lambda = 3, -1}$$

Similarly to Q5

$$Q23 \quad \bar{A}' = \begin{bmatrix} 3 & 2 \\ 1 & 3 \end{bmatrix} \quad \bar{B}' = \begin{bmatrix} 2 & 5 \\ 3 & 2 \end{bmatrix}$$

$$f_{\text{inv}}(AB)$$

$$\text{Since } (AB)^{-1} = (\bar{B}')^{-1} \cdot (\bar{A}')^{-1} = \begin{bmatrix} 2 & 5 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} 3 & 2 \\ 1 & 3 \end{bmatrix}$$

$$= \begin{bmatrix} 6+5 & 4+15 \\ 9-2 & 6-6 \end{bmatrix} = \begin{bmatrix} 11 & 19 \\ 7 & 0 \end{bmatrix}$$

$$\Rightarrow X = \bar{A}'^{-1} b$$

$$X = \begin{bmatrix} 2 & 3 \\ 4 & 1 \end{bmatrix} \begin{bmatrix} 5 \\ 3 \end{bmatrix} = \begin{bmatrix} 10+9 \\ 20+3 \end{bmatrix} = \begin{bmatrix} 19 \\ 23 \end{bmatrix}$$

$$\Rightarrow X = \bar{A}'^{-1} b$$

$$X = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 10 & 0 \\ 0 & 0 & 0 & 10 \end{bmatrix} \begin{bmatrix} 5 \\ 3 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 5 \\ 6 \\ 10 \\ 10 \end{bmatrix}$$

$$\Rightarrow X = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 10 & 0 \\ 0 & 0 & 0 & 10 \end{bmatrix} \begin{bmatrix} 5 \\ 3 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 5 \\ 6 \\ 10 \\ 10 \end{bmatrix}$$

$$\Rightarrow X = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 10 & 0 \\ 0 & 0 & 0 & 10 \end{bmatrix} R_1 \leftarrow$$

Q29(c)

$$\begin{bmatrix} 1 & -1 & 1 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 1 \end{bmatrix} R_{2+R_1}$$

$$\begin{bmatrix} 1 & 0.4 & 0 & 0.2 & 0 & 0 \\ 0 & -0.2 & 0 & -0.6 & +0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} R_{2-3R_1}$$

$$\begin{bmatrix} 1 & 0.4 & 0 & 0.2 & 0 & 0 \\ 0 & -1 & 0 & 3 & -5 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{bmatrix} \frac{R_2}{0.2}$$

$$\begin{bmatrix} 1 & 0 & -1 & 2 & 0 & 0 \\ 0 & -1 & 0 & 3 & -5 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{bmatrix} R_{1-0.4R_2}$$

$$\begin{bmatrix} 1 & 0 & -1 & 2 & 0 & 0 \\ 0 & -1 & 0 & 3 & -5 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{bmatrix} -\frac{R_3}{4}$$

$$\text{So } \bar{A} = \begin{bmatrix} -1 & 2 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\begin{array}{c} \begin{bmatrix} 1 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ -1 & 0 & -1 & 0 & 0 & 0 \\ -1 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \end{bmatrix} R_{1+R_2} \\ \begin{bmatrix} 1 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ -1 & 0 & -1 & 0 & 0 & 0 \\ -1 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \end{bmatrix} R_{2+R_3} \\ \begin{bmatrix} 1 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ -1 & 0 & -1 & 0 & 0 & 0 \\ -1 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \end{bmatrix} R_{3+R_4} \end{array}$$

Mr. Jamal Nasir
Lecturer UET Peshawar

$$\text{Q30} \quad A' = \begin{bmatrix} 1 & 0 & -1 \\ 0 & 0 & 1 \\ -1 & 0 & 0 \\ 1 & 0 & 1 \end{bmatrix}$$

Q_{30} is similar to Q_{29} .

Q_{32} is similar to Q_{31}

$$\text{Q31(a)} \quad \begin{bmatrix} 1 & -1 & 1 & 1 \\ -1 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{bmatrix}$$

$$\sim \begin{bmatrix} 1 & -1 & 1 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 1 \end{bmatrix} R_2 + R_3$$

$$\sim \begin{bmatrix} 1 & -1 & 1 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{bmatrix} R_3 + R_1$$

As non-trivial so matrix is singular.

$$\sim \begin{bmatrix} 1 & 0 & 0 & 100 \\ 0 & 1 & 0 & 110 \\ 0 & 0 & 1 & 001 \end{bmatrix}$$

$$\sim \begin{bmatrix} 1 & 0 & 0 & 100 \\ 0 & 1 & 0 & 110 \\ 0 & 0 & 0 & 111 \end{bmatrix} R_3 + R_1$$

$$\sim \begin{bmatrix} 1 & 0 & 0 & 100 \\ 0 & 1 & 0 & 110 \\ 0 & 0 & 0 & 111 \end{bmatrix} R_{2+R_3}$$

$$x_1 = 9$$

$$x_2 = 5$$

$$x_3 = 2$$

$$\text{eq (1)} \Rightarrow UX = \bar{x}$$

$$\begin{bmatrix} 1 & -4 & 0 \\ 0 & 2 & 1 \\ 0 & 0 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 9 \\ 5 \\ 2 \end{bmatrix} = \begin{bmatrix} \bar{x}_1 \\ \bar{x}_2 \\ \bar{x}_3 \end{bmatrix}$$

Solving the above equation we get

$$2x_2 + x_3 = 5 \quad (1)$$

$$2x_3 = 2 \quad (2)$$

$$x_1 + 4x_2 = 9 \quad (1)$$

$$\sim \begin{bmatrix} 1 & 0 & 0 & 100 \\ 0 & 1 & 0 & 110 \\ 0 & 0 & 1 & 101 \end{bmatrix} R_{2+R_1} \text{ As inverse exist so it's trivial.}$$

$$\text{Q1} \quad A = \begin{bmatrix} 2 & 8 & 0 \\ 2 & 2 & -3 \\ 1 & 2 & 7 \end{bmatrix}, \quad b = \begin{bmatrix} 18 \\ 3 \\ 12 \end{bmatrix}, \quad L = \begin{bmatrix} 2 & 0 & 0 \\ 2 & -3 & 0 \\ 1 & -1 & 4 \end{bmatrix}, \quad U = \begin{bmatrix} 1 & 4 & 0 \\ 0 & 2 & -1 \\ 0 & 0 & 1 \end{bmatrix}.$$

$$\text{Now } UX = \bar{x} \quad (1) \quad L\bar{x} = b \quad (2)$$

$$\text{eq (1)} \Rightarrow \cancel{U\bar{x}} \quad L\bar{x} = b$$

$$\begin{bmatrix} 2 & 0 & 0 \\ 2 & -3 & 0 \\ 1 & -1 & 4 \end{bmatrix} \begin{bmatrix} \bar{x}_1 \\ \bar{x}_2 \\ \bar{x}_3 \end{bmatrix} = \begin{bmatrix} 18 \\ 3 \\ 12 \end{bmatrix}$$

$$2\bar{x}_1 = 18 \quad (1) \quad 2\bar{x}_1 - 3\bar{x}_2 = 3 \quad (2)$$

$$\bar{x}_1 - \bar{x}_2 + 4\bar{x}_3 = 12 \quad (3)$$

As non-trivial so matrix is singular.

From above equation we get

$$\boxed{x_1=1} \quad \boxed{x_2=2} \quad \boxed{x_3=1}$$

$$x = \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} \text{ Ans.}$$

θ_2 & θ_3 is similarly to θ_1

$$A = \begin{bmatrix} 2 & 3 & 4 \\ 4 & 5 & 10 \\ 4 & 8 & 2 \end{bmatrix}; b = \begin{bmatrix} 6 \\ 16 \\ 2 \end{bmatrix}$$

For matrix U (Upper triangular)

$$\sim \left[\begin{array}{ccc|c} 2 & 3 & 4 & R_2 - 2R_1 \\ 0 & -1 & 2 & \\ 0 & 2 & -6 & R_3 - 2R_1 \end{array} \right]$$

$$\sim \left[\begin{array}{ccc|c} 2 & 3 & 4 & \\ 0 & -1 & 2 & \\ 0 & 0 & -2 & R_3 + 2R_2 \end{array} \right]$$

$$2x_1 + 3x_2 + 4x_3 = 6 \quad (1)$$

$$-x_2 + 2x_3 = 4 \quad (2)$$

$$-2x_3 = -2 \quad (3)$$

$$L = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\text{Now } L\bar{z} = b$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} z_1 \\ z_2 \\ z_3 \end{bmatrix} = \begin{bmatrix} 6 \\ 16 \\ 2 \end{bmatrix}$$

$$z_1 = 6 \quad (1)$$

$$2z_1 + z_2 = 16 \quad (2)$$

$$2z_1 + 2z_2 + z_3 = 2 \quad (3)$$

Solving we get

$$\boxed{\bar{z}_1=6}$$

$$\boxed{\bar{z}_2=4}$$

$$\boxed{\bar{z}_3=-2}$$

Now solving

$$UX = \bar{z}$$

$$\begin{bmatrix} 2 & 3 & 4 \\ 0 & -1 & 2 \\ 0 & 0 & -2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 6 \\ 4 \\ -2 \end{bmatrix}$$

For matrix L (Lower triangular)

Ex 2.3

$$A = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}, V = \begin{bmatrix} 1 & 2 & 1 & 2 \\ 1 & 1 & 3 & 3 \end{bmatrix}, (1,1) (2,1) (1,3) (2,2)$$

Now we get

$$\boxed{X_1 = 4} \quad \boxed{X_2 = -2} \quad \boxed{X_3 = 1}$$

$$X = \begin{bmatrix} 4 \\ -2 \\ 1 \end{bmatrix}$$

$\Phi_6 = \Phi_{10}$ similarly to Φ_5


$$= \begin{bmatrix} -1x_1 + 0x_2 & -1x_2 + 0x_1 & -1x_1 + 0x_3 & -1x_2 + 0x_3 \\ 0x_1 + 1x_2 & 0x_2 + 1x_1 & 0x_1 + 1x_3 & 0x_2 + 1x_3 \end{bmatrix}$$

$$\text{Given } f(v) = AV$$

$$= \begin{bmatrix} -1 & 0 & 1 & 2 \\ 0 & 1 & 1 & 3 \end{bmatrix}.$$

$$f(v) = \begin{bmatrix} -1 & -2 & -1 & -2 \\ 1 & 1 & 3 & 3 \end{bmatrix} \text{ image.}$$

Graph image solid



MN. Jamal Nasir

Lecturer U.E.T. Peshawar

Spring Semester-2012

Electrical Engg:

Qasim Shahab
0300 5829015
Anif: 0345 92289401
Ph: 091-5610987

Jai Laser Photostate

Matrix to M.A.
Notes are available

Shop #53 coffee shop bazar Islamia college Peshawar university

MN. Jamal Nasir
Lecturer U.E.T. Peshawar
Spring Semester-2012 Electrical Engg:

Q2 Let f be the shear in the X -direction

$$f(v) = \begin{bmatrix} 1 & k \\ 0 & 1 \end{bmatrix} v.$$


$$\text{where } k=3 \quad \text{and } v = \begin{bmatrix} 1 & 1 & 3 & 3 \end{bmatrix}^T$$

$$f(v) = \begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix} v$$


$$\Rightarrow f(v) = \begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 3 & 3 \end{bmatrix}$$

$$= \begin{bmatrix} 4 & 13 & 6 & 15 \end{bmatrix}^T \text{ image}$$

Graphs



Graphs



Q3 $A = \begin{bmatrix} 1 & 0 \\ k & 1 \end{bmatrix} \quad k = -2 \quad \text{then } A = \begin{bmatrix} 1 & 0 \\ -2 & 1 \end{bmatrix}$

$$A v = \begin{bmatrix} 1 & 0 \\ -2 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 3 & 3 \end{bmatrix}^T$$

$$= \begin{bmatrix} 1 & 1 & 3 & 3 \\ -1 & 2 & -5 & -2 \end{bmatrix}^T \text{ image.}$$

Graphs

Scanned by CamScanner


$$\text{Q4(a)} \quad A = \begin{bmatrix} k & 0 \\ 0 & k \end{bmatrix}, \quad k=4$$

$$V = \begin{bmatrix} 1 & 1 & 3 & 3 \\ 1 & 4 & 1 & 4 \end{bmatrix}$$

$$f_{(W)}, \Delta V = \begin{bmatrix} 4 & 0 \\ 0 & 4 \end{bmatrix} \begin{bmatrix} 1 & 1 & 3 & 3 \\ 1 & 4 & 1 & 4 \end{bmatrix}$$

$$= \begin{bmatrix} 4 & 4 & 12 & 12 \\ 4 & 16 & 4 & 16 \end{bmatrix} \text{ image.}$$

Graph




$$\text{Q5} \quad k=2, \quad A = \begin{bmatrix} k & 0 \\ 0 & 1 \end{bmatrix} \Rightarrow A = \begin{bmatrix} 2 & 0 \\ 0 & 1 \end{bmatrix}$$

$$V = \begin{bmatrix} 0 & 0 & 1 & 1 \\ 0 & 1 & 0 & 1 \end{bmatrix}$$

$$f_{(W)} = \Delta V = \begin{bmatrix} 2 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & 0 & 1 & 1 \\ 0 & 1 & 0 & 1 \end{bmatrix}$$


$$= \begin{bmatrix} 0 & 0 & 2 & 2 \\ 0 & 1 & 0 & 1 \end{bmatrix} \text{ image.}$$

Graphs



$$\text{(b)} \quad k=1/4 \quad \begin{bmatrix} 1/4 & 0 \\ 0 & 1/4 \end{bmatrix} \begin{bmatrix} 1 & 1 & 3 & 3 \\ 1 & 4 & 1 & 4 \end{bmatrix} = \begin{bmatrix} 0.25 & 0.25 & 0.75 & 0.75 \\ 0.25 & 1 & 0.25 & 1 \end{bmatrix}$$

Graph



Q6 is similarly to Q5


$$\text{Q7} \quad f(v) = \begin{bmatrix} -2 & 1 \\ 3 & 4 \end{bmatrix} v$$

$$T = V = \begin{bmatrix} 5 & 0 & 2 \\ 0 & 3 & -1 \end{bmatrix}$$

$$f(v) = \begin{bmatrix} -2 & 1 \\ 3 & 4 \end{bmatrix} \begin{bmatrix} 5 & 0 & 2 \\ 0 & 3 & -1 \end{bmatrix}$$

$$= \begin{bmatrix} -10 & 3 & -5 \\ 15 & 12 & 2 \end{bmatrix}$$

Graph



Q8 is similar to Q7.


$$\text{Q8} \quad T = \begin{bmatrix} 1 & -3 & 2 \\ 1 & -3 & -1 \end{bmatrix} \quad A = \begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix} \quad \theta = 60^\circ$$

$$A = \begin{bmatrix} \cos 60 & -\sin 60 \\ \sin 60 & \cos 60 \end{bmatrix} = \begin{bmatrix} 0.5 & -0.866 \\ 0.866 & 0.5 \end{bmatrix} \quad (\underline{\underline{\text{Exp 9 Ex 1.5}}})$$

$$f(T) = AT = \begin{bmatrix} 0.5 & -0.866 \\ 0.866 & 0.5 \end{bmatrix} \begin{bmatrix} 1 & -3 & 2 \\ 1 & -3 & -1 \end{bmatrix}$$

$$= \begin{bmatrix} -0.366 & 1.866 \\ 1.366 & -4.098 \end{bmatrix} \begin{bmatrix} 1 & -3 & 2 \\ 1 & -3 & -1 \end{bmatrix}$$

Graph



$$(1.866, 1.366)$$

$$(-0.366, 1.866)$$

$$(1.098, -4.098)$$

Qasr Shabab
0300 5639013
Arif: 0305 9280401
Ph: 031 5610987

Jam Laser Photostate
Matrix to M.A.
Notes are available

Shop #59 coffee shop bazar Islamia college Peshawar University

$\mathcal{Q}_{10}(A \otimes g)$

$$A = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \quad (\exp s \cos i, \sin i) \quad \text{if } u = \begin{bmatrix} x \\ y \end{bmatrix}.$$

$$f_2(u) = Av = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -x \\ y \end{bmatrix}.$$

$$f_1(f_2(u)) = f_1(Av) = B(Av)$$

$$B = \begin{bmatrix} \cos \pi/2 & \sin \pi/2 \\ -\sin \pi/2 & \cos \pi/2 \end{bmatrix} \quad (f_1 \circ f_2 \text{ or } s).$$

$$B(Av) = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} y \\ -x \end{bmatrix}.$$

Now

$$f_1(u) = \begin{bmatrix} \cos \pi/2 & \sin \pi/2 \\ -\sin \pi/2 & \cos \pi/2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} y \\ -x \end{bmatrix}.$$

$$f_2(f_1(u)) = f_2(Bu) = A(Bu)$$

$$= \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} y \\ -x \end{bmatrix} = \begin{bmatrix} -y \\ x \end{bmatrix}.$$

$$\text{Hence } f_1(f_2(u)) \neq f_2(f_1(u)).$$

$$A = \begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix}, \quad T = \begin{bmatrix} 1 & -3 & 2 \\ 1 & -3 & -1 \end{bmatrix}$$

$$f(v) = Av = \begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix} \begin{bmatrix} 1 & -3 & 2 \\ 1 & -3 & -1 \end{bmatrix}$$


$$= \begin{bmatrix} 3 & -9 & 0 \\ 6 & -18 & 0 \end{bmatrix}.$$

$$\mathcal{Q}_{12} \quad A = \begin{bmatrix} f_h & 0 \\ 0 & k \end{bmatrix} + f_h = 0, \quad k = 3 \quad (\text{given})$$

$$A = \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix}, \quad V = \begin{bmatrix} 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & -1 \end{bmatrix}.$$

$$Av = \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & -1 \end{bmatrix}$$

$$= \begin{bmatrix} 0 & 2 & 2 & 0 \\ 0 & 0 & 3 & 3 \end{bmatrix}$$



\mathcal{Q}_{12} is similarly to \mathcal{Q}_{13} .

JAN

PHOTOSTAT

**FROM MATRIC TO M.A & M.SC
NOTES ARE AVAILABLE**

ADDRESS:

JAN PHOTOSTAT, COFFEE SHOP MARKET,
UNIVERSITY OF PESHAWAR,

PROPRIETOR:
QAISAR SHABAB, ARIF JAN

Phone No: 091-5610987

Mobile No: 0300-5829015, 0345-9165401
Arif Jan: 0345-9289401

The transition Probabilities arranged in n × n matrix
 $T = [t_{ij}]$ is called transition matrix of markov chain

where

$$t_{1j} + t_{2j} + t_{3j} + \dots + t_{nj} = 1 \quad \text{--- (1)}$$

Entries in each column of T are non-negative
 added upto 1 as in (1)

(a) no transition b/c $a_{11} + a_{21} = 0.3 + 0.4 \neq 1$

$$\begin{aligned} \text{(b) transition as } 0.2 + 0.8 + 0.0 &= 1 \\ 0.3 + 0.5 + 0.2 &= 1 \\ 0.1 + 0.7 + 0.2 &= 1. \end{aligned}$$

$$\begin{aligned} \text{(c) transition because } 0.65 + 0.45 &= 1 \\ 0.33 + 0.67 &= 1. \end{aligned}$$

(d) no transition because All each column are not
 equal to 1.

Probability vector

The vector $U = \begin{bmatrix} u_1 \\ u_2 \\ u_3 \end{bmatrix}$ is called probability vector if

$$u_1 + u_2 + \dots + u_n = 1$$

$$\text{Ques (a)} \quad \frac{1}{2} + \frac{1}{3} + \frac{2}{3} = \frac{3+2+2}{6} = \frac{9}{6} \neq \text{not p.}$$

$$\text{(b)} \quad 0+1+0=1 \quad \text{Prob. value}$$

$$\text{(c)} \quad \frac{1}{4} + \frac{1}{6} + \frac{1}{3} + \frac{1}{4} = \frac{3+2+4+3}{12} = \frac{12}{12} = 1 \quad \text{Prob. value}$$

$$\text{(d)} \quad \frac{1}{5} + \frac{2}{5} + \frac{1}{10} + \frac{2}{10} = \frac{2+4+1+2}{10} = \frac{9}{10} \neq 1 \quad \text{not P.V.}$$

$$\begin{array}{l} \text{For } j=3 \\ a_{13} + a_{23} + a_{33} = 1 \\ 0.3 + 0.5 + a_{33} = 1 \\ a_{33} = 1 - 0.3 - 0.5 = 0.2 \end{array}$$

j=1

$$0.2 + 0.3 + a_{13} = 1$$

$$a_{13} = 1 - 0.2 - 0.3 = 0.5$$

$$\boxed{a_{13}=0.5}$$

$$\begin{array}{l} \text{For } j=2 \\ a_{12} + a_{22} + a_{32} = 1 \end{array}$$

$$a_{22} + a_{32} = 1 - 0.1 = 0.9 \quad (\text{So these are more})$$

$$\begin{array}{l} \text{All possible values:} \\ \boxed{a_{22} + a_{32} = 0.9} \end{array}$$

$$\begin{array}{l} \text{For } j=1 \\ a_{11} + a_{21} + a_{31} = 1 \\ \square + 0.3 + \square = 1 \end{array}$$

$$a_{11} + a_{31} = 1 - 0.3 = 0.7$$

$a_{11} + a_{31} = 0.7$ So there are more than possible values. ($0.5 + 0.2$) etc.

$$\begin{array}{l} \text{For } j=2 \\ \begin{bmatrix} a_{12} & 0.4 & 0.3 \\ 0.3 & \square & 0.5 \\ \square & 0.2 & \square \end{bmatrix} \end{array}$$

$$\begin{array}{l} \text{For } j=3 \\ a_{13} + a_{23} + a_{33} = 1 \end{array}$$

$$a_{13} = 1 - 0.3 - 0.5 = 0.2$$

$$\boxed{a_{13} = 0.2}$$

$$\boxed{F_{11}=2}$$

$$\begin{array}{l} a_{12} + a_{22} + a_{32} = 1 \\ 0.4 + \boxed{a_{22}} + 0.2 = 1 \end{array}$$

$$a_{22} = 1 - 0.2 - 0.4 = 0.4 = \boxed{a_{22} = 0.4}$$

$$(5) T = \begin{bmatrix} 0.7 & 0.4 \\ 0.3 & 0.6 \end{bmatrix}$$

(a) If $\chi^{(0)} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$

$$\chi^{(1)} = \overline{T}\chi^{(0)} = \begin{bmatrix} 0.7 & 0.4 \\ 0.3 & 0.6 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 0.7 \\ 0.3 \end{bmatrix}.$$

$$\chi^{(2)} = \overline{T}\chi^{(1)} = \begin{bmatrix} 0.7 & 0.4 \\ 0.3 & 0.6 \end{bmatrix} \begin{bmatrix} 0.7 \\ 0.3 \end{bmatrix} = \begin{bmatrix} 0.572 \\ 0.39 \end{bmatrix}.$$

$$\chi^{(3)} = \overline{T}\chi^{(2)} = \begin{bmatrix} 0.7 & 0.4 \\ 0.3 & 0.6 \end{bmatrix} \begin{bmatrix} 0.572 \\ 0.39 \end{bmatrix} = \begin{bmatrix} 0.583 \\ 0.417 \end{bmatrix}.$$

It's steady state vector.

Q6 is similarly to Q5

(b) Show that T is regular.

$$\text{Now } T^2 = \overline{T} \cdot \overline{T} = \begin{bmatrix} 0.7 & 0.4 \\ 0.3 & 0.6 \end{bmatrix} \begin{bmatrix} 0.7 & 0.4 \\ 0.3 & 0.6 \end{bmatrix} = \begin{bmatrix} 0.61 & 0.52 \\ 0.32 & 0.48 \end{bmatrix}.$$

All entries are non-zero so T is regular

T is regular if all entries in some power of T are positive

Steady state vector

$$0.572, 0.417$$

So all entries in T^2 are not non-zero so T is not regular.
* Q(c) & d proved similarly to above part &

$$\chi^{(4)} = T\chi^{(3)} = \begin{bmatrix} 0.7 & 0.4 \\ 0.3 & 0.6 \end{bmatrix} \begin{bmatrix} 0.583 \\ 0.417 \end{bmatrix} = \begin{bmatrix} 0.575 \\ 0.425 \end{bmatrix}.$$

$$(5) \chi = \overline{T}\chi = \begin{bmatrix} 0.7 & 0.4 \\ 0.3 & 0.6 \end{bmatrix} \begin{bmatrix} 0.575 \\ 0.425 \end{bmatrix} = \begin{bmatrix} 0.573 \\ 0.428 \end{bmatrix}.$$

$$(6) \chi = \overline{T}\chi = \begin{bmatrix} 0.7 & 0.4 \\ 0.3 & 0.6 \end{bmatrix} \begin{bmatrix} 0.572 \\ 0.417 \end{bmatrix} = \begin{bmatrix} 0.572 \\ 0.417 \end{bmatrix}.$$

same.

$$(b) (a) \quad \overline{T}^2 = \begin{bmatrix} 0 & 1/2 \\ 1 & 1/2 \end{bmatrix} \begin{bmatrix} 0 & 1/2 \\ 1 & 1/2 \end{bmatrix} = \begin{bmatrix} 1/2 & 1/4 \\ 1/2 & 3/4 \end{bmatrix}.$$

So no entry is 0 in \overline{T}^2 so \overline{T} is regular.

$$\overline{T}^2 = \overline{T} \cdot \overline{T} = \begin{bmatrix} 1/2 & 0 & 0 \\ 0 & -1/2 & 0 \\ 0 & 0 & 1/2 \end{bmatrix} \begin{bmatrix} 1/2 & 0 & 0 \\ 0 & 1 & 1/2 \\ 1/2 & 0 & 1/2 \end{bmatrix}$$

$$= \begin{bmatrix} 1/4 & 0 & 0 \\ 1/4 & -1/4 & 0 \\ 1/2 & 0 & 1/4 \end{bmatrix}$$

So all entries in \overline{T}^2 are not non-zero so \overline{T} is not regular.

$$\underline{\underline{Q9}}$$

(a) $T^2 = T, T = \begin{bmatrix} 1/2 & 0 \\ 1/2 & 1 \end{bmatrix} \begin{bmatrix} 1/2 & 0 \\ 1/2 & 1 \end{bmatrix}$

$$= \begin{bmatrix} 1/2 & 0 \\ 3/4 & 1 \end{bmatrix}$$

if's not regular because $t_{22}=0$.

(b) $\text{Let } X^{(0)} = \begin{bmatrix} 0.2 \\ 0.8 \end{bmatrix} \text{ (initial)}$

$$X^{(1)} = T X^{(0)} = \begin{bmatrix} 1/2 & 0 \\ 1/2 & 1 \end{bmatrix} \begin{bmatrix} 0.2 \\ 0.8 \end{bmatrix} = \begin{bmatrix} 0.1 \\ 0.9 \end{bmatrix}$$

$$X^{(2)} = T X^{(1)} = \begin{bmatrix} 0.5 & 0 \\ 0.5 & 1 \end{bmatrix} \begin{bmatrix} 0.1 \\ 0.9 \end{bmatrix} = \begin{bmatrix} 0.05 \\ 0.95 \end{bmatrix}$$

$$X^{(3)} = T X^{(2)} = \begin{bmatrix} 0.5 & 0 \\ 0.5 & 1 \end{bmatrix} \begin{bmatrix} 0.05 \\ 0.95 \end{bmatrix} = \begin{bmatrix} 0.025 \\ 0.975 \end{bmatrix}$$

$$X^{(4)} = T X^{(3)} = \begin{bmatrix} 0.5 & 0 \\ 0.5 & 1 \end{bmatrix} \begin{bmatrix} 0.025 \\ 0.975 \end{bmatrix} = \begin{bmatrix} 0.0125 \\ 0.9875 \end{bmatrix}$$

$$X^{(5)} = T X^{(4)} = \begin{bmatrix} 0.5 & 0 \\ 0.5 & 1 \end{bmatrix} \begin{bmatrix} 0.0125 \\ 0.9875 \end{bmatrix} = \begin{bmatrix} 0.00625 \\ 0.99375 \end{bmatrix} \Rightarrow \begin{bmatrix} 0.000 \\ 1 \end{bmatrix}$$

(a) $T = \begin{bmatrix} 1/3 & 1/2 \\ 2/3 & 1/2 \end{bmatrix}$

$$T^2 = \begin{bmatrix} 0.333 & 0.5 \\ 0.667 & 0.5 \end{bmatrix}$$

$$T^3 = \begin{bmatrix} 0.444 & 0.4166 \\ 0.5555 & 0.5833 \end{bmatrix}$$

$$T^4 = \begin{bmatrix} 0.42859 & 0.43057 \\ 0.5714 & 0.5694 \end{bmatrix}$$

$$T^5 = \begin{bmatrix} 0.4289 & 0.4282 \\ 0.5709 & 0.5717 \end{bmatrix}$$

$$T^6 = \begin{bmatrix} 0.4284 & 0.4285 \\ 0.5715 & 0.5719 \end{bmatrix}$$

$$T^7 = \begin{bmatrix} 0.4285 & 0.4285 \\ 0.5714 & 0.5714 \end{bmatrix}$$

Same.

$$T = \begin{bmatrix} 0.4285 & 0.4285 \\ 0.5714 & 0.5714 \end{bmatrix}$$

Steady State

JAN

PHOTOSTAT

FROM MATRIC TO M.A & M.SC

NOTES ARE AVAILABLE

ADDRESS:

JAN PHOTOSTAT, COFFEE SHOP MARKET,
UNIVERSITY OF PESHAWAR

PROPRIETOR:
QAISAR SHABAB, ARIF JAN

Phone No: 091-5610987

Mobile No: 0300-5829015, 0345-9165402
Arif Jan: 0345-9289401

$$\vec{PQ} = \begin{bmatrix} -2 \\ 5 \end{bmatrix} \text{ where } P=(3,2) \quad Q=? \quad \text{let } Q(a,b)$$

$$\text{then } \vec{PQ} = (a-3, b-2) = (-2, 5)$$

$$a-3 = -2 \Rightarrow \boxed{a=1} \quad \text{and } b-2 = 5 \Rightarrow \boxed{b=7}$$

Q4 is similarly to Q5.

Q5
Find $U+V$, $U-V$, $2U$, $4U-3V$ if

$$U = (2, 3), \quad V = (-2, 5).$$

$$U+V = (2+(-2), 3+5) = (0, 8)$$

$$U-V = (2+2, 3+5) = (4, -2)$$

$$2U = 2[2, 3] = [4, 6].$$

$$3U-2V = 3[2, 3] - 2[-2, 5] = (6, 9) - (-4, 10)$$

$$= (6+4, 9-10)$$

$$= (10, -1)$$

(b) & (c) Part similarly to part (a).

Q6 is similarly to Q5.

$$U = (1, 2), \quad V = (-3, 4), \quad W = (w_1, 4) \quad \text{and } Y = (-2, y_2).$$

$$(a) \quad W = 2U$$

$$(b) \quad (w_1, 4) = 2(1, 2)$$

$\Sigma x = 4.1$

70

$$(w_1, 4) = (2, 4).$$

$$\Rightarrow w_1 = 2$$

$$(b) \quad \frac{3}{2}x = V$$

$$\frac{3}{2}(-2, x_2) = (-3, 4)$$

$$(-3, \frac{3}{2}x_2) = (-3, 4)$$

$$\Rightarrow \frac{3}{2}x_2 = 4 \quad \Rightarrow \boxed{x_2 = \frac{8}{3}}$$

$$\begin{aligned} & \text{① - ②} \\ & \boxed{C_2 = -8} \\ & \therefore 0 \Rightarrow C_1 - 24 = -5 \quad \Rightarrow \boxed{C_1 = 19} \end{aligned}$$

$$\text{eq } ② \Rightarrow (-5, 6) = 19(1, 2) - 8(3, 4)$$

$$\underline{\underline{Q_9}}(a) (1, 2)$$

$$\|U\| = \sqrt{(1)^2 + (2)^2} = \sqrt{5}$$

b, c, d is similarly to a.

Q_{10} is similarly to Q_9 .

$$Q_{11} \|u\| = \sqrt{(\beta-2)^2 + (4-3)^2} = \sqrt{1+1} = \sqrt{2}.$$

Q_{12} is similarly to Q_{11} .

$$\begin{aligned} & (-5, 6) = C_1 \begin{bmatrix} 1, 2 \end{bmatrix} + C_2 \begin{bmatrix} 3, 4 \end{bmatrix} \quad \text{--- ④} \\ & \Rightarrow C_1 + 2C_2 + (C_2, 3 + C_2, 4) \\ & (-5, 6) = (C_1 + 3C_2, 2C_1 + 4C_2) \end{aligned}$$

$$\begin{aligned} & C_1 + 3C_2 = -5 \quad \text{--- ①} \\ & 2C_1 + 4C_2 = 6 \\ & \div 2 \quad \Rightarrow C_1 + 2C_2 = 3 \quad \text{--- ②} \end{aligned}$$

$$\begin{aligned} & \text{① - ②} \\ & \boxed{C_2 = -8} \end{aligned}$$

$$\therefore 0 \Rightarrow C_1 - 24 = -5 \quad \Rightarrow \boxed{C_1 = 19}$$

$$\begin{aligned} & \underline{\underline{Q_9}}(b) \\ & C_1 \begin{bmatrix} 1 \\ 2 \end{bmatrix} + C_2 \begin{bmatrix} 3 \\ 4 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \\ & \begin{bmatrix} C_1 \\ 2C_1 \end{bmatrix} + \begin{bmatrix} 3C_2 \\ 4C_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \\ & \begin{bmatrix} C_1 + 3C_2 \\ 2C_1 + 4C_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \\ & \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix} \begin{bmatrix} C_1 \\ C_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}. \end{aligned}$$

Augmented form

$$\left[\begin{array}{ccc|c} -3 & 0 \\ 2 & 4 & 0 \end{array} \right]$$

$$R_1 \rightarrow \left[\begin{array}{ccc|c} -1 & 0 \\ 0 & 2 & 0 \end{array} \right] R_2 \rightarrow$$

$$-2R_2 \rightarrow \boxed{L_2 = 0}$$

so system is impossible.

(iii)

$$= \frac{1}{2} \left| \det \left(\begin{bmatrix} 3 & 3 & 1 \\ -1 & 1 & 1 \\ 4 & -1 & 1 \end{bmatrix} \right) \right|$$

$$= \frac{1}{2} \left| \begin{vmatrix} 3 & 1 & 1 \\ -1 & 1 & 1 \\ 4 & 1 & 1 \end{vmatrix} + \begin{vmatrix} 3 & 1 & 1 \\ 1 & 1 & 1 \\ 4 & -1 & 1 \end{vmatrix} \right|$$

$$= \frac{1}{2} \left| 4 \begin{vmatrix} 0 & 1 \\ 3 & 1 \end{vmatrix} \right|$$

$$= \frac{1}{2} \left| (4(0-3)) \right| = \frac{1}{2} | -12 | = \frac{1}{2} (12) = 6.$$


$$\text{Q.P. } \Delta_{\text{REV}} = \left| \det \left(\begin{bmatrix} 2 & 3 & 1 \\ 5 & 3 & 1 \\ 4 & 5 & 1 \end{bmatrix} \right) \right| \quad (\text{P-221})$$

$$= \frac{1}{2} \left| -6 + 2 + 16 \right|$$

$$= \frac{1}{2} | 18 - 6 |$$

$$= \frac{1}{2} | 12 | = 6$$

$$= 6$$



(iv)

Ex 41

81

71

$$\underline{\text{Q19}} \quad (a) \quad X = (3, 4)$$

$$= \frac{3+4}{\sqrt{3^2+4^2}} = \frac{3+4}{\sqrt{25}}, \quad \frac{3+4}{5} = \left(\frac{3}{5}, \frac{4}{5} \right)$$

Find c such that θ_{19} is similarly to θ_{19} .

θ_{20} is similarly to θ_{19} .

$$\underline{\text{Q20}} \quad (a) \quad C \propto \theta = \frac{U \cdot V}{\|U\| \|V\|} - C \quad U = (1, 2), \quad V = (2, -3)$$

$$\text{where } \|U\| = \sqrt{1^2 + 2^2} = \sqrt{5}$$

$$\Rightarrow \|V\| = \sqrt{(2)^2 + (-3)^2} = \sqrt{4+9} = \sqrt{13}$$

$$U \cdot V = (1, 2) \cdot (2, -3) = (1)(2) + (2)(-3) = 2 - 6 = -4$$

$$\therefore \theta_{20} = \frac{-4}{\sqrt{5} \cdot \sqrt{13}}$$

θ_{21} , θ_{22} and θ_{23} are similarly to θ_{19} .

θ_{22} is similarly to θ_{21} .

θ_{23} is S.V.S

$$(a) \quad \text{For orthogonal. } U \cdot V = 0$$

$$U_1 \cdot U_4 = ((1+2)i) \cdot (-6i+3j) = (-8+6) = 0$$

$$\text{So } (U_1, U_4) = (1+2i) \cdot (2i+j) = (-4+4) = 0$$

$$U_3 \cdot U_4 = (-2i-4j) \cdot (-2i+j) = (4+4) = 0$$

$$U_3 \cdot U_5 = (-2i-4j) \cdot (-6i+3j) = (12+12) = 0$$

$$U_4 \cdot U_5 = (-2i+4j) \cdot (2i+4j) = (-4+4) = 0$$

$$U_5 \cdot U_6 = (2i+4j) \cdot (-6i+3j) = (-12+12) = 0$$

(b) $U_4 \perp U_5$ because $U_5 = 2U_1$,

$U_4 \perp U_6$ because $U_6 = 3U_4$.

(c) $U_1 \perp U_3$ because $U_3 = -2U_1$

$U_3 \perp U_5$ because $U_3 = -U_5$.

Find slopes $m_1 = \text{slope } m_2$ (whether are parallel or slopes are equal).

$$m_1 = \frac{4}{a}, \quad m_2 = \frac{5}{2}$$

$$\therefore \frac{4}{a} = \frac{5}{2}$$

$$\boxed{\frac{8}{a} = \frac{8}{5}}$$

Q26 Vectors are orthogonal if $\cos\theta = 0$.

$$(a_{12}) \cdot (a_{-2}) = 0$$

$$a^2 + (-4) = 0$$

$$a^2 = 4$$

$$\boxed{a = \pm 2}$$

Q27 (a) $i+3j$, (b) $-2i-3j$, (c) $-2i$, (d) $3j$


$$U-V = \begin{bmatrix} 1 & -0 \\ 2 & -3 \\ -3 & +2 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}.$$

$$2U = 2 \begin{bmatrix} 1 \\ -3 \end{bmatrix} = \begin{bmatrix} 2 \\ -6 \end{bmatrix}$$

$$3U-2V = 3 \begin{bmatrix} 1 \\ 2 \\ -3 \end{bmatrix} - 2 \begin{bmatrix} 0 \\ -1 \\ 2 \end{bmatrix} = \begin{bmatrix} 3 \\ 6 \\ -9 \end{bmatrix} - \begin{bmatrix} 0 \\ -2 \\ 4 \end{bmatrix} = \begin{bmatrix} 3-0 \\ 6-(-2) \\ -9+4 \end{bmatrix} = \begin{bmatrix} 3 \\ 8 \\ -5 \end{bmatrix}.$$

Part bisects as "y"

Q2 as same as Q1



$$U = 0i - 300j + V = -400i + 0j$$

$$\begin{aligned} \overrightarrow{AC} &= \overrightarrow{AB} + \overrightarrow{BC} \\ &= \overrightarrow{V} + \overrightarrow{U} \\ &= (-400i + 0j) + (0i - 300j) \\ &= -400i - 300j \end{aligned}$$

$$|\overrightarrow{AC}| = \sqrt{(400)^2 + (300)^2} = 500$$

Q29

$$U = \begin{bmatrix} 1 \\ -2 \\ 3 \end{bmatrix}, V = \begin{bmatrix} -3 \\ 1 \\ 3 \end{bmatrix}, W = \begin{bmatrix} 9 \\ 1 \\ b \end{bmatrix}, X = \begin{bmatrix} 3 \\ c \\ 2 \end{bmatrix}.$$

Find a, b & c so that:

$$(a) W = \frac{1}{2}U \Rightarrow \begin{bmatrix} 9 \\ 1 \\ b \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 1 \\ -2 \\ 3 \end{bmatrix} = \begin{bmatrix} \frac{1}{2} \\ -1 \\ \frac{3}{2} \end{bmatrix}$$

$$\Rightarrow \begin{cases} 9 = \frac{1}{2} \\ 1 = -1 \\ b = \frac{3}{2} \end{cases} \Rightarrow \begin{cases} b = 18 \\ b = -2 \\ b = 1.5 \end{cases}$$

$$\therefore \begin{cases} a = 18 \\ b = -2 \\ c = 1.5 \end{cases}$$

$$U = (4, 5, -2, 3) \quad V = (3, -2, 0, 1), \quad C = 2, \quad D = 3$$

$$\begin{array}{l} \text{L.H.S} \\ U + V = \boxed{V + U} \end{array}$$

$$\begin{array}{l} \text{R.H.S} \\ U + V = (4+3, 5-2, -2+0, 3+1) = (7, 3, -2, 4) \\ V + U = (3+4, -2+5, 0-2, 1+3) = (7, 3, -2, 4). \end{array}$$

$$\begin{aligned} (b) \quad W + V &= U \\ \Rightarrow \begin{bmatrix} a \\ b \end{bmatrix} + \begin{bmatrix} -3 \\ 3 \end{bmatrix} &= \begin{bmatrix} 1 \\ 2 \end{bmatrix} \\ \begin{bmatrix} a-3 \\ b+3 \end{bmatrix} &= \begin{bmatrix} 1 \\ 2 \end{bmatrix} \end{aligned}$$

$$\begin{aligned} a-3=1 \Rightarrow a &= 1+3 = 4 \Rightarrow \boxed{a=4} \\ b+3=3 \Rightarrow b &= 0 \end{aligned}$$

$$(c) \quad U + V = W$$

$$\begin{array}{l} \text{L.H.S} \\ U + (V + W) \end{array}$$

$$\begin{array}{l} (4, 5, -2, 3) + [(3, -2, 0, 1) + (-3, 2, -5, 3)] \\ (4, 5, -2, 3) + (0, 0, -5, 4) \end{array}$$

$$(4, 5, -7, 7).$$

$$\text{R.H.S } (U + V) + W$$

$$[(4, 5, -2, 3) + (3, -2, 0, 1)] + (-3, 2, -5, 3).$$

$$(7, 3, -2, 4) + (-3, 2, -5, 3)$$

$$(4, 5, -7, 7).$$

$$\text{Hence } U + (V + W) = (U + V) + W$$

$$\text{Q. 4. as same as Q. 3.}$$

$$(d) \quad U + O = O + U.$$

$$\begin{array}{l} \text{L.H.S} \\ U + O = (4, 5, -2, 3) + (0, 0, 0, 0) = (4, 5, -2, 3) \\ \text{R.H.S} \quad (0, 0, 0, 0) + (4, 5, -2, 3) = (4, 5, -2, 3). \end{array}$$

(d)

$$\begin{aligned}
 & U + (-U) = 0 \\
 & (4, 5, -2, 3) + (-4, 5, -2, 3) \\
 & (4, 5, -2, 3) + (-4, 5, 2, -3) \\
 & (4-4, 5-5, -2+2, 3-3) \\
 & (0, 0, 0, 0).
 \end{aligned}$$

(e) $C(U+V) = CU+CV$

$$L.H.S = 2[(4, 5, -2, 3) + (3, -2, 0, 1)]$$

$$\begin{aligned}
 & = 2(7, 3, -2, 4) \\
 & = (14, 6, -4, 8).
 \end{aligned}$$

$$R.H.S 2(4, 5, -2, 3) + 2(3, -2, 0, 1)$$

$$= (8, 10, -4, 6) + (6, -4, 0, 2)$$

$$= (14, 6, -4, 8).$$

(f) $(C+\delta)U = CU + \delta U$


$$L.H.S (2+3)(4, 5, -2, 3) = 5(4, 5, -2, 3) = (20, 25, -10, 15)$$

$$R.H.S 2(4, 5, -2, 3) + 3(4, 5, -2, 3)$$

$$(8, 10, -4, 6) + (12, 15, -6, 9)$$

$$= (20, 25, -10, 15)$$

(g)



\vec{PQ} same as \vec{Q}_1

If (a) P (1st point), Q (Second point).

$$\vec{PQ} = \vec{Q} - \vec{P} = (0, 0, 2) - (2, 3, 1)$$

$$= (-2, -3, 3)$$

$$\vec{PQ} = \begin{bmatrix} -2 \\ -3 \\ 3 \end{bmatrix}.$$

\vec{PQ} , b , c , d same as Pack a.

Qaiser Shabab
0300 5820015

Arif: 0345 9289401
Ph: 091-5610987

Jai Laser Photostate

Matrix to M.A.
Notes are available

$$\underline{\underline{Q_9}} \quad \text{head } \beta = \beta = (1, -2, 3) \quad \vec{P}_0(\text{head}) = (3, 4, -1).$$

$$\vec{P}_{\bar{Q}} = \vec{Q} - \vec{P}$$

$$\Rightarrow \bar{Q} = \vec{P}_{\bar{Q}} + \vec{P} = (3, 4, -1) + (1, -2, 3)$$

$$\bar{Q} = (3+1, 4-2, -1+3)$$

$$\bar{Q} = (4, 2, 2).$$

$$\underline{\underline{Q_{10}}} \quad (a) \|U\| = \sqrt{(1)^2(2)^2(-3)^2} = \sqrt{14+4+9} = \sqrt{14} = \sqrt{14},$$

Part b, c, d same as part a.

$$Q \parallel \text{ same } Q_{10}.$$

$$Q_{12}(a) \cdot (1, -1, 2), (3, 0, 2),$$

$$\|U - V\| = \sqrt{(-3)^2 + (-1+0)^2 + (2-2)^2},$$

$$= \sqrt{4+1+0}$$

$$= \sqrt{5}.$$

Part b, c, d same as part a.

Q₁₃ same as Q₁₂.

$$\alpha \begin{bmatrix} 1 \\ 2 \\ -3 \end{bmatrix} + b \begin{bmatrix} -1 \\ 1 \\ 1 \end{bmatrix} + c \begin{bmatrix} -1 \\ 4 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ 2 \\ 3 \end{bmatrix}.$$

$$\begin{bmatrix} a & -b & -c \\ 2a & +b & +4c \\ -3a & +b & -c \end{bmatrix} = \begin{bmatrix} 2 \\ -2 \\ 3 \end{bmatrix}.$$

$$\begin{aligned} a - b - c &= 2 \quad (1) \\ 2a + b + 4c &= -2 \quad (2) \\ -3a + b - c &= 3 \quad (3) \end{aligned}$$

$$\begin{aligned} \text{add (2) times first to (2)} \\ -2a + 2b + 2c &= -4 \\ 2a + b + 4c &= -2 \end{aligned}$$

$$3b + 6c = -6$$

$$b + 2c = -2 \quad (2')$$

$$\text{add (2) times (1) to (3)}:$$

$$\begin{aligned} 3a - 3b - 3c &= 6 \\ -3a + b - c &= 3 \\ -2b - 4c &= 9 \end{aligned}$$

$$2b - 4c = -9 \quad (3)'$$

$$\text{add (2) times (2)' to (3)'}:$$

add (4.2) times ① to ③.

$$\begin{array}{l} 2c_1 + 2c_2 + 6c_3 = 0 \\ -c_1 - 2c_2 - 4c_3 = 0 \end{array}$$

$$\frac{c_1 + 2c_3 = 0}{D = -5}$$

which is impossible.

$$\text{Q.E.D.} \\ c_1 \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix} + c_2 \begin{bmatrix} 1 \\ 3 \\ -2 \end{bmatrix} + c_3 \begin{bmatrix} 3 \\ -4 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} c_1 \\ 2c_1 \\ -c_1 \end{bmatrix} + \begin{bmatrix} c_2 \\ 3c_2 \\ -2c_2 \end{bmatrix} + \begin{bmatrix} 3c_3 \\ -4c_3 \\ c_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} c_1 + c_2 + 3c_3 \\ 2c_1 + 3c_2 + 7c_3 \\ -c_1 - 2c_2 - 4c_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}.$$

$$\begin{aligned} c_1 + c_2 + 3c_3 &= 0 \quad \text{--- ①} \\ 2c_1 + 3c_2 + 7c_3 &= 0 \quad \text{--- ②} \\ -c_1 - 2c_2 - 4c_3 &= 0 \quad \text{--- ③} \end{aligned}$$

$$\text{Add } (-2) \text{ times ① to ②.} \\ -2c_1 - 3c_2 + 7c_3 = 0 \quad \text{--- ④} \\ -c_1 - 2c_2 - 4c_3 = 0 \quad \text{--- ⑤}$$

$$c_2 + c_3 = 0.$$

$$\boxed{c_2 = -c_3}$$

$$\text{Let } c_3 = \gamma \\ \text{Then } c_2 = -\gamma \quad \text{and } c_1 = -2\gamma$$

For possible solution put $\gamma = 1$ (unit value)

$$c_3 = 1 \rightarrow c_2 = -1, \quad c_1 = -2(1) = -2$$

$$\text{Then } \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix} = \begin{bmatrix} -2 \\ -1 \\ 1 \end{bmatrix}.$$

$$\| (1, -3, 2) \| = 5$$

$$\sqrt{(1)^2 + (3)^2 + (2)^2} = 5$$

$$\sqrt{14 + a^2} = 5$$

$$\therefore 13 + 2a^2 = 25$$

$$a^2 = 25 - 14 = 11 \Rightarrow a = \pm \sqrt{11}$$

$$\boxed{c_1 = -2c_3},$$

$$U \cdot V = -7 = V \cdot U.$$

$$(U+V) \cdot W = U \cdot W + V \cdot W.$$

$$\underline{\underline{Q.F}} \quad U = (1, 2, 1, 0) + V = (1, -1, -2, 3).$$

$$+ V \cdot V = 0.$$

$$(1, 2, 1, 0) \cdot (1, -1, -2, 3) = 0$$

$$(1^2 - 2, -2 - 3) = 0$$

$$1^2 - 3 \cdot 1 - 4 = 0$$

$$1^2 - 2, -2 - 3) = 0$$

$$1^2 - 3 \cdot 1 - 4 = 0$$

$$\alpha = \frac{-(-5) \pm \sqrt{(-5)^2 - 4(1)(-4)}}{2(1)}$$

$$\boxed{\alpha_1 = 4, -1}$$

$$\text{Ans} \quad C = 3, \quad U = (1, 2, 1), \quad V = (1, 2, -4), \quad W = (1, 0, 2).$$

$$(a, b) \geq 0$$

$$(1, 2, 3) \cdot (1, 2, 3) \geq 0$$

$$14 + 9 > 0$$

$$14 > 0$$

$$\therefore 7 \leq 7\sqrt{6} \quad \text{Ans.}$$

$$\|U\| = \sqrt{U \cdot U} = \sqrt{(1, 2, 1) \cdot (1, 2, 1)} = \sqrt{1^2 + 4 + 1} = \sqrt{6}$$

$$\|V\| = \sqrt{V \cdot V} = \sqrt{(1, -1, -2) \cdot (1, -1, -2)} = \sqrt{1 + 1 + 4} = \sqrt{6}$$

$$\|W\| = \sqrt{W \cdot W} = \sqrt{1 + 0 + 4} = \sqrt{5}$$

$$R.H.S \quad (1, 2, 1) \cdot (1, 0, 2) + (1, 2, -4) \cdot (1, 0, 2)$$

$$(1+6) + (1+8)$$

$$(8-8) = 0.$$

$$\|U \cdot V\| \leq \|U\| \|V\|$$

$$\|U \cdot V\| = \|(1, 2, 1) \cdot (1, 2, -4)\|$$

$$= |1^2 + 2^2 + 3(-4)| = ||1+4-12|| = |-7| = 7.$$

$$\|U\| = \sqrt{U \cdot U} = \sqrt{(1, 2, 1) \cdot (1, 2, 1)} = \sqrt{1^2 + 4 + 1} = \sqrt{6}$$

$$\|V\| = \sqrt{V \cdot V} = \sqrt{(1, -1, -2) \cdot (1, -1, -2)} = \sqrt{1 + 1 + 4} = \sqrt{6}$$

$$\|W\| = \sqrt{W \cdot W} = \sqrt{1 + 0 + 4} = \sqrt{5}$$

$$\therefore 7 \leq 7\sqrt{6} \quad \text{Ans.}$$

θ_{21}

θ_{21} is similarly to θ_{20} .

79

$$\theta_{20} \text{ (a)} = U = (1, 2, 3), V = (-4, 4, 5).$$

$$\cos \theta = \frac{U \cdot V}{\|U\| \|V\|} - \textcircled{1}$$

$$\|U\| \|V\|$$

$$U \cdot V = (1, 2, 3) \cdot (-4, 4, 5) = (-4 + 8 + 15) = 19.$$

$$\|U\| = \sqrt{1+2^2+3^2} = \sqrt{14}$$

$$\|V\| = \sqrt{(-4)^2 + (4)^2 + 5^2} = \sqrt{57}.$$

$$\cos \theta = \cos \theta = \frac{|9|}{\sqrt{14} \sqrt{57}} = 0.673.$$

$$\begin{aligned} -2c &= -5 \\ c &= 5/2 \end{aligned}$$

$$(b) \quad U = (1, 2, 3, 1) \quad V = (-3, 1, -2, 0)$$

$$\cos \theta = \frac{U \cdot V}{\|U\| \|V\|} - \textcircled{1}$$

$$U \cdot V = (1, 2, 3, 1) \cdot (-3, 1, -2, 0) = (1+2-6+0) = -3.$$

$$\|U\| = \sqrt{1+2^2+3^2+1^2} = \sqrt{14} \quad \|V\| = \sqrt{9+1+4+0} = \sqrt{14}.$$

$$\cos \theta = \frac{-3}{\sqrt{14} \sqrt{14}} = \frac{-3}{14} = -\frac{3}{14}$$

$$\boxed{\cos \theta = -\frac{3}{14}}.$$

and similarly.

For orthogonal $V \cdot W = 0$.

$$(2, c, 3) \cdot (1, -2, 1) = 0$$

$$2 - 2c + 3 = 0$$

$$-2c = -5$$

$$\boxed{c = 5/2}$$

$$W = (a, b, c), \quad X = (1, 2, 1) \quad Y = (1, -1, 1)$$

$$V \cdot W = (a, b, c) \cdot (1, 2, 1) = (a+2b+c) = 0 \quad \textcircled{1}$$

$$V \cdot X = (a, b, c) \cdot (1, -1, 1) = (a-b+c) = 0 \quad \textcircled{2}$$

$\textcircled{1} - \textcircled{2}$

$$\begin{aligned} a+2b+c &= 0 \\ a-b+c &= 0 \\ \hline 3b &= 0 \Rightarrow \boxed{b=0} \end{aligned}$$

θ_{22} S.Y.S

θ_{23} is similarly θ_{24} ($\alpha, \beta, 1$).

$$X = (2, -1, 3)$$

Unit vector in the direction of X

$$\begin{aligned} \text{Q1} &\Rightarrow a+c=0 \Rightarrow a=-c \\ \text{Q2} &\Rightarrow a+c=0 \Rightarrow a=c \\ \text{Let } c=1 &\text{ Then } a=-1, b=0 \end{aligned}$$

For possible solution put $c=1$

$$c=1, a=-1, b=0$$

$$\boxed{a=-1, b=0, c=1}$$

$$\begin{aligned} \text{Q3} &+ \text{strong inequality } \|u+v\| \leq (\|u\| + \|v\|)^2 \\ &\quad \text{if } u=(1, 2, 3, -1) \text{ and } v=(1, 0, -2, 3) \end{aligned}$$

$$u=(2, 2, 1, 2)$$

$$v=(1, 0, -2, 3)$$

$$\begin{aligned} \|u+v\| &= \sqrt{(2^2 + 2^2 + 1^2 + (-2)^2)^2} = \sqrt{4+4+1+4} = \sqrt{13} \\ \|u+v\|^2 &= (\sqrt{13})^2 = 13 \quad \text{Q1} \\ \|u\| &= \sqrt{|1+4+9|} = \sqrt{15} \end{aligned}$$

$$\|v\| = \sqrt{(1^2 + 0^2 + (-2)^2 + 3^2)^2} = \sqrt{1+4+9} = \sqrt{14}$$

$$\|u\| + \|v\| = \left(\sqrt{15} + \sqrt{14} \right)^2 = 57.98 = 58 \quad \text{Q2}$$

$$\text{From Q1 and Q2 we have } \|u+v\|^2 \leq (\|u\| + \|v\|)^2.$$

$$\text{Q1} \Rightarrow U = \frac{(2, -1, 3)}{\sqrt{14}} = \left(\frac{2}{\sqrt{14}}, -\frac{1}{\sqrt{14}}, \frac{3}{\sqrt{14}} \right)$$

$$\|M\| = \sqrt{4+1+9} = \sqrt{14}$$

Part b, c, d all same as Part (a).

Q28 is same as Q27.

$$(a) (1, 2, -3) = i+2j-3k$$

Part b, c, d also same as part (a).

$$\text{for (b) } 2i+3j-4k$$

$$2i = 2\Gamma_1 = 2 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 2 \\ 0 \\ 0 \end{bmatrix}$$

$$3j = 3\Gamma_2 = 3 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 3 \\ 0 \end{bmatrix}$$

$$-4k = -4\Gamma_3 = -4 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ -4 \end{bmatrix}$$

$$\begin{aligned} \text{3x1 matrix} \quad & \begin{bmatrix} 2 \\ 0 \\ 0 \end{bmatrix} + \begin{bmatrix} 0 \\ 3 \\ 0 \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \\ -4 \end{bmatrix} = \begin{bmatrix} 2 \\ 3 \\ -4 \end{bmatrix} \\ & \text{3x1 matrix} \quad \text{part b, c, d same} \end{aligned}$$

Ex 4.2
For Isosceles triangle two sides are equal.

$$\|R_1 - R_2\| = \sqrt{(2.5)^2 + (3.1)^2 + (4.2)^2} = \sqrt{41}$$

$$\|R_1 - R_3\| = \sqrt{(2.5)^2 + (3.1)^2 + (4.4)^2} = \sqrt{100} = 10$$

$$\|R_2 - R_3\| = \sqrt{(3.5)^2 + (1)^2 + (2.4)^2} = \sqrt{41}.$$

From above we find

$$\|R_1 - R_2\| = \|R_2 - R_3\| \text{ verified.}$$

Ex 4.2 If length of the two sides are not equal

$\cos \theta = 0$

$$\vec{U} = \vec{P}_1 \vec{P}_2 = (3-2, 1-3, 2+4) = (1, -2, 6)$$

$$\vec{V} = \vec{P}_1 \vec{P}_3 = (7-2, 0-3, 1+4) = (5, -3, 5)$$

$$\vec{W} = \vec{P}_2 \vec{P}_3 = (7-3, 0-1, 1-2) = (4, -1, -1)$$

$$\vec{U} \cdot \vec{V} = 5 + 6 + 30 = 41$$

$$\vec{V} \cdot \vec{W} = (-5 + 6 + 30) = 41$$

$$\vec{U} \cdot \vec{W} = (4 + 2 - 6) = 0$$

Jam Laser Photostate

Matrix to M.A.
Notes are available

Qasim Shahab

0300 5629915

Aarif 0345 9269401

Pk: 091-5610987

$$U = \begin{bmatrix} u \\ u_1 \\ u_{200} \end{bmatrix}$$

8% increase

$$U = \begin{bmatrix} u_1 + 0.08u_1 \\ u_2 + 0.08u_2 \\ \vdots \\ u_{200} + 0.08u_{200} \end{bmatrix} = \begin{bmatrix} (1+0.08)u_1 \\ (1+0.08)u_2 \\ \vdots \\ (1+0.08)u_{200} \end{bmatrix}$$

$$U = \begin{bmatrix} 1.08u_1 \\ 1.08u_2 \\ \vdots \\ 1.08u_{200} \end{bmatrix} = 1.08U.$$

Ex 4.2 U.V will show price of old cd players.
Speakers and cassette recorders.

$$\frac{1}{2} (t+a) (\alpha \delta \gamma).$$

$$\underline{\text{Q36}} \quad U = (1, 1, 0, 0) \quad UV = 0 \quad \text{if } V = \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix}$$

Now $UV = 0$

$$\begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix} + \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} \Rightarrow \begin{bmatrix} a+1 \\ b+1 \\ c \\ d \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\Rightarrow a=1, \quad b=1, \quad c=0, \quad d=0$$

$$V = (1, 1, 0, 0)$$

Q37 Same as Q36.

$$\underline{\text{Q38}} \quad U = (1, 0, 1) \quad \text{and } V = (a, b, c).$$

$$U \cdot V = 0$$

$$(1, 0, 1) \cdot (a, b, c) = 0$$

$$a+0+c = 0$$

$$a+c = 0 \quad \text{---} \textcircled{1}$$

For bit matrix $b=0, \quad b=1$.

If $a=0, \quad b=0$, then $a=c=0$
 If $b=1$ then $a=c=1$

Now $U+V = (x_1, y_1) + (x_2, y_2)$
 Q38 same as Q39.

Linear Transformation:- A linear transformation

L of \mathbb{R}^n into \mathbb{R}^m is a function assigning a unique vector $L(u)$ in \mathbb{R}^m to each u in \mathbb{R}^n such that

- (i) $L(u+v) = L(u) + L(v)$, for every u and v in \mathbb{R}^n
- (ii) $L(ku) = kL(u)$, for every u in \mathbb{R}^n and every scalar k .

Which of the following are linear transformation?

$$L(x, y) = (x+1, y, x+y).$$

If $\vec{U} = (x_1, y_1)$ and $\vec{V} = (x_2, y_2)$ be the two vectors in \mathbb{R}^2

$$\text{Then } L(\vec{U}) = L(x_1, y_1) = (x_1+1, y_1, x_1+y_1).$$

$$\therefore L(\vec{V}) = L(x_2, y_2) = (x_2+1, y_2, x_2+y_2)$$

$$L(\vec{U}) + L(\vec{V}) = (x_1+1, y_1, x_1+y_1) + (x_2+1, y_2, x_2+y_2)$$

$$= (x_1+x_2+2, y_1+y_2, x_1+y_1+x_2+y_2) \quad \text{---} \textcircled{1}$$

$$\text{Now } \vec{U} + \vec{V} = (x_1, y_1) + (x_2, y_2)$$

$$= (x_1+x_2, y_1+y_2).$$

$$L(\vec{U} + \vec{V}) = (x_1 + y_1 + z_1, y_1 + y_2, x_1 + x_2 + y_1 + y_2) \quad (2)$$

From eq (1) & eq (2) we have:

$$L(\vec{U} + \vec{V}) \neq L(\vec{U}) + L(\vec{V}).$$

It's not a linear transformation.

$$(b) L\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} x+y \\ y \\ x-z \end{pmatrix}$$

$$L: \mathbb{R}^3 \rightarrow \mathbb{R}^3$$

Let $\vec{U} = \begin{pmatrix} x_1 \\ y_1 \\ z_1 \end{pmatrix}$ and $\vec{V} = \begin{pmatrix} x_2 \\ y_2 \\ z_2 \end{pmatrix}$ be the two vectors in \mathbb{R}^3 .

$$\text{then } L(\vec{U}) = L\begin{pmatrix} x_1 \\ y_1 \\ z_1 \end{pmatrix} = \begin{pmatrix} x_1 + y_1 \\ y_1 \\ x_1 - z_1 \end{pmatrix}$$

$$\text{and } L(\vec{V}) = L\begin{pmatrix} x_2 \\ y_2 \\ z_2 \end{pmatrix} = \begin{pmatrix} x_2 + y_2 \\ y_2 \\ x_2 - z_2 \end{pmatrix}.$$

$$\text{Now } L(\vec{U}) + L(\vec{V}) = \begin{pmatrix} x_1 + y_1 \\ y_1 \\ x_1 - z_1 \end{pmatrix} + \begin{pmatrix} x_2 + y_2 \\ y_2 \\ x_2 - z_2 \end{pmatrix}$$

$$= \begin{pmatrix} (x_1 + y_1) + (x_2 + y_2) \\ y_1 + y_2 \\ (x_1 - z_1) + (x_2 - z_2) \end{pmatrix}$$

$$\rightarrow (1)$$

$$\text{From eq (1) & eq (2) we have: } L(\vec{U} + \vec{V}) \neq L(\vec{U}) + L(\vec{V}) \quad (A)$$

$$L(\vec{U} + \vec{V}) = L\begin{pmatrix} x_1 + x_2 \\ y_1 + y_2 \\ z_1 + z_2 \end{pmatrix} = \begin{pmatrix} (x_1 + x_2) + (y_1 + y_2) \\ y_1 + y_2 \\ (x_1 + x_2) - (z_1 + z_2) \end{pmatrix} \rightarrow (2)$$

From eq (1) & eq (2) we have:

$$L(\vec{U} + \vec{V}) \neq L(\vec{U}) + L(\vec{V}).$$

Checking 2nd Propert:

$$L(k\vec{U}) = k(L\vec{U}).$$

$$k\vec{U} = k\begin{pmatrix} x_1 \\ y_1 \\ z_1 \end{pmatrix} = \begin{pmatrix} kx_1 \\ ky_1 \\ kz_1 \end{pmatrix}.$$

$$L(k\vec{U}) = L\begin{pmatrix} kx_1 \\ ky_1 \\ kz_1 \end{pmatrix} = \begin{pmatrix} kx_1 + ky_1 \\ ky_1 \\ kx_1 - kz_1 \end{pmatrix}.$$

$$= \begin{pmatrix} x_1 + y_1 \\ y_1 \\ x_1 - z_1 \end{pmatrix}$$

$$\Rightarrow L(k\vec{U}) = k(L\vec{U}) \quad (B)$$

From eq (A) & eq (B) it's clear that it's a linear transformation from \mathbb{R}^3 to \mathbb{R}^3 .

$\times 4.3$

$\theta_2 + \theta_3$ is similarly to θ_1

$$(c) L(x, y) = (\tilde{x}+x, \tilde{y}-y)$$

sols.

$$\text{Let } \vec{U} = (x_1, y_1) \text{ & } \vec{V} = (x_2, y_2) \text{ be two vectors in } \mathbb{R}^2$$

$$\text{Then } L(\vec{U}) = L(x_1, y_1) = (\tilde{x}_1+x_1, \tilde{y}_1-y_1)$$

$$+ L(\vec{V}) = L(x_2, y_2) = (\tilde{x}_2+x_2, \tilde{y}_2-y_2)$$

$$L(\vec{U})+L(\vec{V}) = (\tilde{x}_1+x_1, \tilde{y}_1-y_1) + (\tilde{x}_2+x_2, \tilde{y}_2-y_2)$$

$$= [(x_1+\tilde{x}_1)+(x_2+\tilde{x}_2), (y_1-\tilde{y}_1)+(y_2-\tilde{y}_2)] - \text{Q.E.D}$$

$$\text{Now } \vec{U}+\vec{V} = (x_1, y_1) + (x_2, y_2)$$

$$= (x_1+x_2, y_1+y_2)$$

$$L(\vec{U}+\vec{V}) = L(x_1+x_2, y_1+y_2)$$

$$= (x_1+\tilde{x}_1+y_1, y_1-y_2)$$

$$= (x_1+y_1, y_1+y_2) + (y_1+y_2, y_1-y_2) - \text{Q.E.D}$$

From $\theta_1 + \theta_2 + \theta_3 = 0$ we have

$$L(\vec{U})+L(\vec{V}) \neq L(\vec{U}+\vec{V})$$

It's not a linear transformation.

$$(a) L \begin{pmatrix} u_1 \\ u_2 \\ u_3 \\ u_4 \end{pmatrix} = \begin{pmatrix} u_1 \\ \tilde{u}_1+u_2 \\ u_3 \\ u_4-u_3 \end{pmatrix}$$

$$L(\vec{U}) = L \begin{pmatrix} u_1 \\ u_2 \\ u_3 \\ u_4 \end{pmatrix} + \vec{V} = \begin{pmatrix} u_1 \\ u_2 \\ u_3+u_2 \\ u_4 \end{pmatrix}$$

be the two vector in \mathbb{R}^4 then

$$L(\vec{V}) = L \begin{pmatrix} v_1 \\ v_2 \\ v_3 \\ v_4 \end{pmatrix} = \begin{pmatrix} v_1 \\ v_2 \\ v_3+u_2 \\ v_4-u_3 \end{pmatrix}$$

$$L(\vec{U})+L(\vec{V}) = \begin{pmatrix} u_1 \\ u_2+u_2 \\ u_3+u_2 \\ u_4-u_3 \end{pmatrix} + \begin{pmatrix} v_1 \\ v_2+v_2 \\ v_3+u_2 \\ v_4-u_3 \end{pmatrix}$$

$$= \begin{pmatrix} u_1+v_1 \\ u_2+2v_2 \\ u_3+u_2+v_2 \\ u_4-u_3+v_4 \end{pmatrix} - \text{Q.E.D}$$

Again now

$$\vec{U}+\vec{V} = \begin{bmatrix} u_1 \\ u_2 \\ u_3 \\ u_4 \end{bmatrix} + \begin{bmatrix} v_1 \\ v_2 \\ v_3 \\ v_4 \end{bmatrix} = \begin{bmatrix} u_1+v_1 \\ u_2+v_2 \\ u_3+v_3 \\ u_4+v_4 \end{bmatrix}$$

$$L(\vec{U} + \vec{V}) = L\left(\begin{bmatrix} U_1 + V_1 \\ U_2 + V_2 \\ U_3 + V_3 \\ U_4 + V_4 \end{bmatrix}\right) = \begin{bmatrix} U_1 + V_1 \\ (U_1 + V_1)^2 + (U_2 + V_2)^2 \\ (U_1 + V_1) - (U_3 + V_3) \\ U_4 + V_4 \end{bmatrix} \rightarrow \mathbb{C}$$

From ex ① & ex ② we know

$$L(\vec{U} + \vec{V}) \neq L(\vec{U}) + L(\vec{V})$$

Is not a linear transformation.

$$(b) L\left(\begin{bmatrix} x \\ y \\ z \end{bmatrix}\right) = \begin{bmatrix} 1 & 1 & 0 \\ 0 & -1 & 2 \\ 1 & 1 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

$$= \begin{bmatrix} x+y+0z \\ 0x-y+2z \\ x+y-z \end{bmatrix} = \begin{bmatrix} x+y \\ -y+2z \\ x+y-z \end{bmatrix}$$

Let $\vec{U} = \begin{bmatrix} x_1 \\ y_1 \\ z_1 \end{bmatrix}$ and $\vec{V} = \begin{bmatrix} x_2 \\ y_2 \\ z_2 \end{bmatrix}$ be the two vector in \mathbb{R}^3 , then

$$L(\vec{U}) = L\left(\begin{bmatrix} x_1 \\ y_1 \\ z_1 \end{bmatrix}\right) = \begin{bmatrix} x_1 + y_1 \\ -y_1 + 2z_1 \\ x_1 + y_1 - z_1 \end{bmatrix}.$$

$$L(\vec{V}) = L\left(\begin{bmatrix} x_2 \\ y_2 \\ z_2 \end{bmatrix}\right) = \begin{bmatrix} x_2 + y_2 \\ -y_2 + 2z_2 \\ x_2 + y_2 - z_2 \end{bmatrix}.$$

$$\text{Now } \vec{U} + \vec{V} = \begin{bmatrix} x_1 \\ y_1 \\ z_1 \end{bmatrix} + \begin{bmatrix} x_2 \\ y_2 \\ z_2 \end{bmatrix} = \begin{bmatrix} x_1 + x_2 \\ y_1 + y_2 \\ z_1 + z_2 \end{bmatrix}.$$

$$L(\vec{U} + \vec{V}) = L\left(\begin{bmatrix} x_1 + x_2 \\ y_1 + y_2 \\ z_1 + z_2 \end{bmatrix}\right)$$

$$= \begin{bmatrix} (x_1 + x_2) + (y_1 + y_2) \\ (y_1 + y_2) + 2(z_1 + z_2) \\ (x_1 + x_2) + (y_1 + y_2) - (z_1 + z_2) \end{bmatrix} \rightarrow ②$$

From ex ① & ex ② we know.

$$L(\vec{U}) + L(\vec{V}) = L(\vec{U} + \vec{V}). \quad \text{--- (A)}$$

Checking 2nd Property:

$$L(k\vec{U}) = k(L\vec{U}) \text{ where } k \in \mathbb{R} \text{ then}$$

$$k\vec{U} = k\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} kx \\ ky \\ kz \end{bmatrix}$$

$$f(x_1 + x_2) + (y_1 + y_2) = \begin{bmatrix} x_1 + x_2 \\ y_1 + y_2 \\ z_1 + z_2 \end{bmatrix} + \begin{bmatrix} -y_1 + 2z_1 \\ -y_2 + 2z_2 \\ x_1 + y_1 - z_1 \end{bmatrix} = \begin{bmatrix} (x_1 + y_1) + (x_2 + y_2) \\ (y_1 + y_2) + 2(z_1 + z_2) \\ (x_1 + y_1 - z_1) + (-y_2 + 2z_2) \end{bmatrix} \rightarrow \mathbb{C}$$

$$L(k\vec{v}) = L \begin{bmatrix} kx_1 \\ ky_1 \\ kz_1 \end{bmatrix}$$

$$= \begin{bmatrix} kx_1 + ky_1 \\ ky_1 + kz_1 \\ kz_1 + kx_1 \end{bmatrix} = k \begin{bmatrix} x_1 + y_1 \\ y_1 + z_1 \\ z_1 + x_1 \end{bmatrix}$$

$$\Rightarrow L(k\vec{v}) = k(L\vec{v}) \longrightarrow \textcircled{1}$$


From $\textcircled{1}$ & $\textcircled{2}$ it's clear that it's a linear transformation from \mathbb{R}^3 to \mathbb{R}^3 .

(c) same as


In exercises 5-12, sketch the image of the given line under the given linear transformation L .

$$L(x,y) = (x-y) : P(2,3)$$


$$L(2,3) = (2,3).$$



$$\begin{aligned} L \begin{bmatrix} 1 & 0 \end{bmatrix} &= \begin{bmatrix} 1 & -1 \end{bmatrix} \quad ; \quad u = (1, 0) \\ L \begin{bmatrix} 0 & 1 \end{bmatrix} &= \begin{bmatrix} 0 & -1 \end{bmatrix} \\ L \begin{bmatrix} 1 & 1 \end{bmatrix} &= \begin{bmatrix} 1 & -2 \end{bmatrix} = \begin{bmatrix} 1 & 2 \end{bmatrix} \\ L \begin{bmatrix} 1 & -1 \end{bmatrix} &= \begin{bmatrix} 3 & 0 \end{bmatrix} \end{aligned}$$



$$\begin{aligned} L &= \begin{bmatrix} \cos 30^\circ & -\sin 30^\circ \\ \sin 30^\circ & \cos 30^\circ \end{bmatrix} \quad Q = 30^\circ \quad P = (-1, 3) \quad \therefore u \\ LU &= \begin{bmatrix} \cos 30^\circ & -\sin 30^\circ \\ \sin 30^\circ & \cos 30^\circ \end{bmatrix} \begin{bmatrix} -1 \\ 3 \end{bmatrix}. \\ &= \begin{bmatrix} -\cos 30^\circ - 3\sin 30^\circ \\ \sin 30^\circ + 3\cos 30^\circ \end{bmatrix} = \begin{bmatrix} -2.366, 2.098 \end{bmatrix}. \end{aligned}$$




Q_8 is similarly to Q_7 .

$$L(u) = v \quad , \quad u = (3, 2)$$

$$L(3, 2) = -(-3, 2) = (-3, -2)$$

$$\begin{bmatrix} x+2 \\ y+2 \\ x+2y+2z \end{bmatrix} = \begin{bmatrix} -1 \\ 0 \\ 0 \end{bmatrix}$$

$$v = (3, 2)$$




Q_9 is similarly to Q_9 .

$$Q_{11} \quad L\left(\begin{bmatrix} x \\ y \\ z \end{bmatrix}\right) = \begin{bmatrix} x \\ y \\ x+z \end{bmatrix} \quad u = (2, -1, 3)$$

$$L(u) = L\left(\begin{bmatrix} 2 \\ -1 \\ 3 \end{bmatrix}\right) = \begin{bmatrix} 2 \\ -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix}.$$

Q_{11} is similarly to Q_{11}



$$\therefore \vec{U} = \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix} \quad \text{This shows the image}$$

and goes to the work L .

$$\text{Ex 4.3} \quad \begin{bmatrix} u \\ L(u) \end{bmatrix} = \begin{bmatrix} x \\ y \\ x+z \end{bmatrix} \quad w = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}.$$

$$\text{Image } L(\vec{u}) = \vec{w}$$

$$L\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$$\Rightarrow x + z = 1 - (\text{i}) \quad y + z = -1 - (\text{ii}), \quad x + 2y + 2z = 0 - (\text{iii})$$

$$x = 1 - z \quad (\text{iv})$$

$$+ y = -1 - z \quad (\text{v})$$

$$2\sqrt{3} \Rightarrow (1 - z) + 2(-1 - z) + 2z = 0$$

$$1 - z - 2 - 2z + 2z = 0$$

$$-z - 1 = 0 \Rightarrow \boxed{z = -1}$$

$$\sqrt{3} \Rightarrow x = 1 - (-1) = \boxed{x = 2}$$

$$\sqrt{3} \Rightarrow y = -1 - (-1) = -1 + 1 = 0 = \boxed{y = 0}$$

$$\sqrt{3} \Rightarrow 2 + 2(0) + 2(-1) = 0$$

$$2 + 0 - 2 = 0$$

$$0 = 0.$$

Q4 is similarly to Q3.

$$(b) L(\vec{v}) = w$$

$$\begin{bmatrix} x+z \\ y+z \\ x+2y+2z \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \\ 3 \end{bmatrix}.$$

$$x+z=2 \quad (i)$$

$$y+z=-1 \quad (ii)$$

$$x+2y+2z=3 \quad (iii)$$

$$x+z=y-z \quad (iv)$$

$$y+z=-1-z \quad (v)$$

$$q \sqrt{(iii)} \Rightarrow (2-z) + 2(-1-z) + 2z = 3$$

$$2\cancel{z} + (\cancel{2} - \cancel{2}z) + 2\cancel{z} = 3$$

$$-z = 3 \Rightarrow \boxed{z = -3}$$

$$q \sqrt{(iv)} \Rightarrow x = 2 + 3 = 5 \Rightarrow \boxed{x = 5}$$

$$q \sqrt{(v)} \Rightarrow y = -1 + 3 \Rightarrow \boxed{y = 2}$$

$$\begin{bmatrix} 4x+y+3z \\ 2x-y+3z \\ 2x+2y \end{bmatrix} = \begin{bmatrix} a \\ b \\ c \end{bmatrix}.$$

$$\Rightarrow 4x+y+3z = a \quad (i)$$

$$2x-y+3z = b \quad (ii)$$

$$2x+2y = c \quad (iii)$$

From (iii) by (ii) then subst. from (i)

$$4x+y+3z = a$$

$$\frac{4x+y+3z}{2x-y+3z} = \frac{a}{b}$$

$$3y-3z = a-2b \quad (1)$$

$$\therefore \vec{U} = \begin{bmatrix} y \\ z \\ 1 \end{bmatrix} = \begin{bmatrix} 5 \\ 2 \\ 1 \end{bmatrix}$$

This shows the image so it's
the range L.

$$\text{Let } \vec{U} = \begin{bmatrix} y \\ z \\ 1 \end{bmatrix} \text{ be such that}$$

$$L\vec{U} = \vec{w}$$

$$L \begin{bmatrix} y \\ z \\ 1 \end{bmatrix} = \begin{bmatrix} a \\ b \\ c \end{bmatrix}.$$

$$\begin{bmatrix} 4 & 1 & 3 \\ 2 & -1 & 3 \\ 2 & 2 & 0 \end{bmatrix} \begin{bmatrix} y \\ z \\ 1 \end{bmatrix} = \begin{bmatrix} a \\ b \\ c \end{bmatrix}.$$

~~4x+y+3z = a
2x-y+3z = b
2x+2y = c~~

$$\text{As } \begin{bmatrix} 4 \\ -3 \end{bmatrix} = 4i - 3j \quad \text{where } L(i) = \begin{bmatrix} 2 \\ 3 \end{bmatrix} \text{ and } L(j) = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

$$\Rightarrow L \begin{bmatrix} 4 \\ -3 \end{bmatrix} = L(4i - 3j)$$

$$= 4L(i) - 3L(j)$$

$$\begin{aligned} & \text{eq ④} - \text{eq ③} \\ & 2x - y + 3z = b \\ & 2x + 2y = c \\ & -3y + 3z = b - c \end{aligned}$$

$$2\sqrt{④} + 2\sqrt{⑤}$$

$$3y - 3z = a - 2b$$

$$-3y + 3z = b - c$$

$$0 = a - 2b + b - c$$

$$a - b - c = 0$$

$$\Rightarrow c - a + b = 0 \quad \text{Ans.}$$

$\sim \sim \sim \sim$


Q16 is similarly to Q15.

$$\text{As } \begin{bmatrix} 2 \\ -3 \end{bmatrix} = 2i + j + 3k$$

$$\Rightarrow L \begin{bmatrix} 2 \\ -3 \end{bmatrix} = L(2i + j + 3k) = 2L(i) - L(j) + 3L(k)$$

$$= 2 \begin{bmatrix} 1 \\ 2 \end{bmatrix} - \begin{bmatrix} 1 \\ 2 \end{bmatrix} + 3 \begin{bmatrix} 1 \\ 3 \end{bmatrix}$$

$$= \begin{bmatrix} 2 \\ -2 \end{bmatrix} + \begin{bmatrix} 1 \\ 0 \end{bmatrix} + \begin{bmatrix} 3 \\ 9 \end{bmatrix} = \begin{bmatrix} 2-1+3 \\ -2+0+9 \end{bmatrix}$$



JAN

PHOTOSTAT

FROM MATRIC TO M.A & M.SC
NOTES ARE AVAILABLE

ADDRESS:

UNIVERSITY OF PESHAWAR,

PROPRIETOR:
QAISAR SHABAB, ARIF JAN.

Phone No: 091-5610987

Mobile No: 0300-5829015, 0345-9165402

"It's not a linear transformation."

Let $\vec{U} = (x_1, y_1)$ & $\vec{V} = (x_2, y_2)$. be the two vector
in R^2 , then

$$L(\vec{U}) = L(x_1, y_1) = (x_1 + y_1 + 1, x_1 - y_1)$$

$$L(\vec{V}) = L(x_2, y_2) = (x_2 + y_2 + 1, x_2 - y_2)$$

$$L(\vec{U}) + L(\vec{V}) = (x_1 + y_1 + 1, x_1 - y_1) + (x_2 + y_2 + 1, x_2 - y_2)$$

$$= ((x_1 + y_1) + (x_2 + y_2) + 2, (x_1 - y_1) - (x_2 - y_2)) \rightarrow \textcircled{1}$$

$$\text{Now } \vec{U} + \vec{V} = (x_1, y_1) + (x_2, y_2)$$

$$= (x_1 + x_2, y_1 + y_2)$$

$$L(\vec{U} + \vec{V}) = L(x_1 + x_2, y_1 + y_2)$$

$$= ((x_1 + x_2 + y_1 + y_2 + 1, x_1 + x_2 - y_1 - y_2)) \rightarrow \textcircled{2}$$

From ex (1) & ex (2) it's clearly clear

$$L(\vec{U} + \vec{V}) \neq L(\vec{U}) + L(\vec{V})$$

Ques If $\vec{u} = (x_1, y_1)$ and $\vec{v} = (x_2, y_2)$ be the two vec
in R^2 , then

$$\vec{u} + \vec{v} = (x_1 + y_1) + (x_2 + y_2) = (x_1 + x_2, y_1 + y_2).$$

$$L(\vec{u} + \vec{v}) = L(x_1 + x_2, y_1 + y_2).$$

$$L(\vec{u} + \vec{v}) = (\sin(x_1 + x_2), \sin(y_1 + y_2)) \quad \text{--- (1)}$$

$$L(\vec{u}) = L(x_1, y_1)$$

$$= \sin x_1 + \sin y_1$$

$$L(\vec{v}) = L(x_2, y_2)$$

$$= \sin x_2 + \sin y_2$$

$$\Rightarrow L(\vec{u}) + L(\vec{v}) = (\sin x_1 + \sin y_1) + (\sin x_2 + \sin y_2)$$

$$= (\sin(x_1 + x_2), \sin(y_1 + y_2)) \quad \text{--- (2)}$$

From eq (1) & eq (2) we have

$$L(\vec{u} + \vec{v}) = L(\vec{u}) + L(\vec{v}). \quad \text{--- (A)}$$

$$\text{Now let } k \in R \text{ then } k\vec{u} = k(x_1, y_1) = (kx_1, ky_1)$$

$$L(k\vec{u}) = L(kx_1, ky_1) = (k \sin x_1 + k \cos y_1) = k(\sin x_1 + \cos y_1)$$

$$\Rightarrow L(k\vec{u}) = k(L(\vec{u})) \quad \text{--- (B)}$$

From (A) & (B), we can say that the given function
is a linear transformation.

Theorem If $L: R^n \rightarrow R^m$ is a L.T. Then

$$L(c_1\vec{u}_1 + c_2\vec{u}_2 + \dots + c_n\vec{u}_n)$$

$$= c_1 L(\vec{u}_1) + c_2 L(\vec{u}_2) + \dots + c_n L(\vec{u}_n).$$

for any vec in $\vec{u}_1, \vec{u}_2, \dots, \vec{u}_n$ in R^n and any scalar

$$c_1, c_2, \dots, c_n$$

$$\text{For } R^3 \quad \vec{e}_1 = \vec{e}_1 = (1, 0, 0) = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}.$$

$$\vec{e}_2 = \vec{e}_2 = (0, 1, 0) = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}.$$

$$\vec{e}_3 = \vec{e}_3 = (0, 0, 1) = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}.$$

$$\text{For } R^3 \quad \vec{e} = \vec{e} = (1, 0, 0, \dots, 0) = \begin{bmatrix} 1 \\ 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}.$$

$$\vec{j} = \vec{e}_2 = (0, 1, 0, \dots, 0) = \begin{bmatrix} 0 \\ 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}.$$

$$\vec{k} = \vec{e}_3 = (0, 0, 1, \dots, 0) = \begin{bmatrix} 0 \\ 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}.$$

$$\text{For } R^n \quad \vec{e}_1 = \vec{e}_1 = (1, 0, 0, \dots, 0) = \begin{bmatrix} 1 \\ 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}.$$

$$\vec{e}_2 = (0, 1, 0, \dots, 0) = \begin{bmatrix} 0 \\ 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}.$$

$$\vec{e}_3 = (0, 0, 1, \dots, 0) = \begin{bmatrix} 0 \\ 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}.$$

$$\vdots$$

$$\vec{e}_n = (0, 0, 0, \dots, -1) = \begin{bmatrix} 0 \\ 0 \\ 0 \\ \vdots \\ -1 \end{bmatrix}.$$

$$Q_{2C} \quad L: R^2 \rightarrow R^2 \quad L\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x-y \\ xy \end{pmatrix}$$

$$\text{For } R^2 \quad e_1 = (1,0) = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad \text{and} \quad e_2 = (0,1) = \begin{pmatrix} 0 \\ 1 \end{pmatrix}.$$

$$L(e_1) = L\begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 1-0 \\ 1 \cdot 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

$$L(e_2) = L\begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 0+1 \\ 0 \cdot 1 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

Thus $A = \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}$. &

Q18 $\theta = 60^\circ$

$$L(u) = \begin{bmatrix} \cos 60 & -\sin 60 \\ \sin 60 & \cos 60 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} \frac{x}{2} + \frac{\sqrt{3}}{2}y \\ \frac{\sqrt{3}}{2}x + \frac{y}{2} \end{bmatrix}.$$

$$\text{For } R^2 \quad e_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad \text{and} \quad e_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

$$L(e_1) = L\begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} \frac{1}{2} + \frac{\sqrt{3}}{2} \cdot 0 \\ \frac{\sqrt{3}}{2} \cdot 1 + 0 \end{pmatrix} = \begin{pmatrix} \frac{1}{2} \\ \frac{\sqrt{3}}{2} \end{pmatrix}.$$

$$L(e_2) = L\begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 + \frac{\sqrt{3}}{2} \cdot 1 \\ \frac{\sqrt{3}}{2} \cdot 0 + 1 \end{pmatrix} = \begin{pmatrix} \frac{\sqrt{3}}{2} \\ 1 \end{pmatrix}.$$

$$A = \begin{pmatrix} \frac{1}{2} & \frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix}.$$

$$Q_{2D} \text{ is same as } Q_{1D}$$

$$L\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x-y \\ xy \end{pmatrix}.$$

Ex 4.3

$$\text{For } R^3 \quad e_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \quad \text{and} \quad e_3 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}.$$

$$L(e_1) = L\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 1-0 \\ 0+0 \\ 0-0 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}.$$

$$L(e_2) = L\begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 0-1 \\ 0+0 \\ 0-0 \end{pmatrix} = \begin{pmatrix} -1 \\ 0 \\ 0 \end{pmatrix}.$$

$$L(e_3) = L\begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 0+0 \\ 0-0 \\ 1-0 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}.$$

Thus $A = \begin{pmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{pmatrix}$.

Q19 $L(u) = -2u = -2 \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -2x \\ -2y \end{bmatrix}.$

$$L(e_1) = L\begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} -2 \\ 0 \end{pmatrix}$$

$$L(e_2) = L\begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ -2 \end{pmatrix}$$

$$L(e_3) = L\begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ -2 \end{pmatrix}$$

$$L(e_4) = L\begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}.$$

Thus $A = \begin{pmatrix} 2 & 0 & 0 \\ 0 & -2 & 0 \\ 0 & 0 & -2 \end{pmatrix}$.

So the message is 71 52 33 47 30 26 84
56 43 99 69 56.

(a) SEND HIM MONEY
14 5 14 4 8 9 13 13 15 14 5 25.

$$\therefore A = \begin{bmatrix} 1 & 2 & 3 \\ 1 & 1 & 2 \\ 0 & 1 & 2 \end{bmatrix}.$$

Breaking message into 4 vector
 $\begin{bmatrix} 19 \\ 5 \end{bmatrix}, \begin{bmatrix} 4 \\ 9 \end{bmatrix}, \begin{bmatrix} 13 \\ 15 \end{bmatrix}, \begin{bmatrix} 14 \\ 5 \end{bmatrix}$

$$\text{Now } L(w) = A \cdot X.$$

$$\Delta X = A \begin{bmatrix} 19 \\ 5 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \end{bmatrix} \begin{bmatrix} 19 \\ 5 \end{bmatrix} = \begin{bmatrix} 71 \\ 35 \end{bmatrix}$$

$$A \cdot \Delta X = ?$$

$$A = \begin{bmatrix} 0 & 1 & -1 \\ 2 & -2 & -1 \\ -1 & 1 & 1 \end{bmatrix}.$$

$$M_1 = \tilde{A}^{-1} L(M_1) = \begin{bmatrix} 0 & 1 & -1 \\ 2 & -2 & -1 \\ -1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 67 \\ 44 \\ 41 \end{bmatrix} = \begin{bmatrix} 3 \\ 5 \\ 12 \end{bmatrix}.$$

$$M_2 = \tilde{A}^{-1} L(M_2) = \begin{bmatrix} 0 & 1 & -1 \\ 2 & -2 & -1 \\ -1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 49 \\ 39 \\ 19 \end{bmatrix} = \begin{bmatrix} 20 \\ 9 \\ 1 \end{bmatrix}.$$

$$M_3 = \tilde{A}^{-1} L(M_3) = \begin{bmatrix} 0 & 1 & -1 \\ 2 & -2 & -1 \\ -1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 13 \\ 15 \\ 62 \end{bmatrix} = \begin{bmatrix} 14 \\ 12 \\ 25 \end{bmatrix}.$$

$$M_4 = \tilde{A}^{-1} L(M_4) = \begin{bmatrix} 0 & 1 & -1 \\ 2 & -2 & -1 \\ -1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 14 \\ 5 \\ 25 \end{bmatrix} = \begin{bmatrix} 14 \\ 15 \\ 20 \end{bmatrix}.$$

$$A \begin{bmatrix} 13 \\ 5 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \end{bmatrix} \begin{bmatrix} 13 \\ 5 \end{bmatrix} = \begin{bmatrix} 84 \\ 45 \end{bmatrix}.$$

$$A \begin{bmatrix} 14 \\ 5 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \end{bmatrix} \begin{bmatrix} 14 \\ 5 \end{bmatrix} = \begin{bmatrix} 99 \\ 56 \end{bmatrix}.$$

So the message is
3 5 18 20 1 9 14 12 25 14 15 20
C E R T A I N L Y N O T
certainly not.

$$\text{Breaking into vector in } \mathbb{R}^3$$

$$\begin{bmatrix} L^{(w)} \\ L^{(w)} \\ L^{(w)} \end{bmatrix} = \begin{bmatrix} 67 \\ 49 \\ 39 \end{bmatrix}, \begin{bmatrix} L^{(w)} \\ L^{(w)} \\ L^{(w)} \end{bmatrix} = \begin{bmatrix} 13 \\ 15 \\ 62 \end{bmatrix}, \begin{bmatrix} L^{(w)} \\ L^{(w)} \\ L^{(w)} \end{bmatrix} = \begin{bmatrix} 14 \\ 5 \\ 25 \end{bmatrix}.$$

$$L(w) = Aw \Rightarrow w = \tilde{A}^{-1} L(w)$$

Q_2 is same as Q_1 .

(Q3) Let $U = 2i - j + 3k$, $V = 2i + j$, $W = 2i - j + 2k$, $C = -3$

(i) $U \times V = -(V \times U)$.

$$U \times V = \begin{vmatrix} i & j & k \\ 1 & 2 & -3 \\ 2 & 3 & 1 \end{vmatrix}$$

$$= 11i - 7j - k \quad \text{--- Q1}$$

$$-(V \times U) = -\begin{vmatrix} i & j & k \\ 2 & 3 & 1 \\ 1 & 2 & -3 \end{vmatrix}$$

$$= -(-11i + 7j + k) = 11i - 7j - k \quad \text{--- Q2}$$

From eq Q1 eq Q2 we have.

$$U \times V = -(V \times U) \text{ proved.}$$

Part b, c, d same as part (a).

$$U \times W = \begin{vmatrix} i & j & k \\ 2 & 1 & -1 \\ 3 & 1 & 1 \end{vmatrix} = 3i - 9j + ok$$

$$= 6 + 11 + 10 = 15 \quad \text{--- Q3}$$

$$(U \times V) \cdot W = (2i + 11j + 5k) \cdot (3i + j + 2k)$$

$$= -6 + 11 + 10 = 15 \quad \text{--- Q4}$$

$$U \times W = \begin{vmatrix} i & j & k \\ 3 & 1 & -1 \\ 3 & 1 & 1 \end{vmatrix} = 3i - 9j + ok$$

$$U \cdot (V \times W) = (2i - j + 3k) \cdot (3i - 9j + ok) \\ = 6 + 9 + 0 = 15 \quad \text{--- Q5}$$

From eq Q4 & Q5 we have
 $(U \times V) \cdot W = U \cdot (V \times W)$ proved.

(Q4) $U = 2i - j + 3k$
 $V = 3i + j - k$
 $W = 3i + j + 2k$

(a) Verify equation (3) ($(U \times V) \cdot W = U \cdot (V \times W)$).

95.

Ex 5.1

Q7) This same as part (e).

Q5 is same as Q4.

Orthogonal: If $\vec{U} \cdot \vec{V}$ is orthogonal to both \vec{U} & \vec{V} i.e.

$$(\vec{U} \times \vec{V}) \cdot \vec{U} = 0 \\ (\vec{U} \times \vec{V}) \cdot \vec{V} = 0.$$

Q6 Express $a \rightarrow -(\vec{d}_1)$

$$(a) \vec{U} = 2\hat{i} + 3\hat{j} + 4\hat{k} \quad \vec{V} = -\hat{i} + \hat{j} - \hat{k}$$

$$\vec{U} \times \vec{V} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 3 & 4 \\ -1 & 1 & -1 \end{vmatrix}$$

$$= -15\hat{i} - 2\hat{j} + 9\hat{k}$$

$$\text{Now } (\vec{U} \times \vec{V}) \cdot \vec{U} = (-15\hat{i} - 2\hat{j} + 9\hat{k}) \cdot (2\hat{i} + 3\hat{j} + 4\hat{k})$$

$$= -30 - 6 + 36$$

$$= -26 + 26 = 0. \\ \therefore (\vec{U} \times \vec{V}) \cdot \vec{U} = 0.$$

$$\text{Also } (\vec{U} \times \vec{V}) \cdot \vec{V} = (-15\hat{i} - 2\hat{j} + 9\hat{k}) \cdot (-\hat{i} + \hat{j} - \hat{k})$$

$$= 15 - 6 - 9 \Rightarrow 15 - 15 = 0$$

$$(\vec{U} \times \vec{V}) \cdot \vec{V} = 0.$$

Which is orthogonal.

Q7 is same as Q6

$$\text{Area of triangle} = \frac{1}{2} |\vec{U} \times \vec{V}| \quad \text{--- (1)}$$

$$\text{Let } \vec{U} = P_1 P_2 = (-3, 1, 1, -2, 4, 3) \\ = (-4, 3, 1)$$

$$\text{And } \vec{V} = P_1 P_3 = (0, 1, 4, -2, 3, -3) \\ = (-1, 6, 0)$$

$$\vec{U} \times \vec{V} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -4 & 3 & -1 \\ -1 & 6 & 0 \end{vmatrix}$$

$$\vec{U} \times \vec{V} = -6\hat{i} - \hat{j} - 21\hat{k}$$

$$\text{Now } |\vec{U} \times \vec{V}| = \sqrt{(-6)^2 + (-1)^2 + (-21)^2} = \sqrt{478}$$

$$\therefore \sqrt{Q} \Rightarrow \text{Area of triangle} = \frac{1}{2} |\vec{U} \times \vec{V}| \\ = \frac{1}{2} \sqrt{478} \text{ (units)}^2.$$

Q10 is same as Q9.

(iii) Area of $\triangle_{gm} = |\vec{U} \times \vec{V}|$. —

$$\text{Now } \vec{U} \times \vec{V} = \begin{vmatrix} i & j & k \\ 1 & 3 & -2 \\ 3 & -1 & -1 \end{vmatrix}$$

$$= -\vec{i} - \vec{j} - 10\vec{k}$$

$$\text{Now } |\vec{U} \times \vec{V}| = \sqrt{(-5)^2 + (-5)^2 + (-10)^2} = \sqrt{150}$$

$$\therefore \text{Area of } \triangle_{gm} = \sqrt{150} \text{ (unit)}.$$

Ques $U = 2i - j$, $V = i - 2j - 2k$ and $W = 3i - j + k$.

$$\text{Volume of } \triangle_{pid} = \begin{vmatrix} U_1 & U_2 & U_3 \\ V_1 & V_2 & V_3 \\ W_1 & W_2 & W_3 \end{vmatrix}$$

$$= \begin{vmatrix} 2 & -1 & 0 \\ 1 & -2 & -2 \\ 3 & -1 & 1 \end{vmatrix}$$

$$= 2(-2+2) + 1(1+6) + 0 \\ = 2(-4) + 7 \\ = -8 + 7 = -1$$

$$\text{Vol } \triangle_{pid} = 1 \text{ (unit)}^3$$

\triangle_3 is same as \triangle_1

$$\Rightarrow \begin{vmatrix} x & y & 1 \\ -2 & -3 & 1 \\ 3 & 4 & 1 \end{vmatrix} = 0$$

$$x(-3-4) - y(-2-3) + 1(-8+9) = 0 \\ -7x + 5y + 1 = 0$$

This is the required equation of line for the given

Points:

Part (b), (c), (d) same as Part (a)

$$\text{Q}_2 \text{ is same as Q}_1.$$

$$\begin{aligned} x &= 3+2t \Rightarrow t = \frac{x-3}{2} \\ y &= -2+3t \Rightarrow t = \frac{y+2}{3} \\ z &= 4-3t \Rightarrow t = \frac{z-4}{-3} \end{aligned}$$

(a) Let $P_1(x_1, y_1) = (-2, -3)$ and $P_2(x_2, y_2) = (3, 4)$.

$$\text{Now } \begin{vmatrix} x & y & 1 \\ x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \end{vmatrix} = 0$$

$$\text{Now } \frac{x-3}{2} = \frac{y+2}{3} = \frac{z-4}{-3} \quad \text{--- (1)}$$

(a) (b), (c)

$$\sqrt{0} \Rightarrow \frac{1-3}{2} = \frac{1+2}{3} = \frac{1-4}{-3}$$

$$\frac{-2}{2} = \frac{3}{3} = \frac{-3}{-3}$$

$$-1 \neq 1 \neq 1$$

Hence all the value of t are not same, Therefore

pts are not lies on the lines.

Part (b) & (c) same as part (a)

(d) (4, -1/2, 5/2)

$$eqn Q \Rightarrow \frac{4-3}{2} = \frac{-1/2+2}{2} = \frac{5/2-4}{-3}$$

$$1/2 = 1/2 = 1/2$$

Hence all the value of t are same so pts lies on line

θ_4 is same as θ_3

$$\text{Ex (a) } P_0 = (3, 4, -2), U = (4, -5, 2).$$

$$-\infty < t < \infty$$

$$x = x_0 + at \Rightarrow x = 3 + 4t$$

$$y = y_0 + bt \Rightarrow y = 4 - 5t$$

$$z = z_0 + ct \Rightarrow z = -2 + 2t$$

part b, c, d same as part (a)

$$\text{Ex (a) } P_0 = (2, -3, 1) \text{ & } P_1 = (4, 2, 5)$$

$$\vec{U} = \vec{P}_0 \vec{P}_1 = (4-2, 2+3, 5-1) = (2, 5, 4)$$

$$\text{Now } P_0 = (2, -3, 1) \text{ & } U = (2, 5, 4)$$

$$x = x_0 + at = 2 + 2t$$

$$y = y_0 + bt \Rightarrow y = -3 + 5t$$

$$z = z_0 + ct \Rightarrow z = 1 + 4t$$

Part (b), (c) & (d) same as part (a).

$$Q(a) P_0 = \begin{pmatrix} a & b & c \\ 0 & 0 & 0 \\ 0 & -3 & 1 \end{pmatrix} + U = \begin{pmatrix} a & b & c \\ 2 & 1 & 4 \\ 2 & 5 & 4 \end{pmatrix}$$

For Symmetric form

$$\frac{x-y}{a} = \frac{y-z}{b} = \frac{z-x}{c}$$

$$\frac{x-y}{2} = \frac{y+z}{5} = \frac{z-1}{4}$$

(b), (c), (d) Part as same as Part (a)

$$-x - 2y + 4z - 3 = 0$$

Jamel Nasir
Lecturer UET Peshawar

$$\left(\begin{array}{ccc} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{array} \right)$$

$$(a) \quad 3(x-2) + 2(y+3) - 4(z-4) = 0 \quad \text{--- Q}$$

$$(0, -2, 3)$$

$$\text{eq } Q \Rightarrow 3(0-2) + 2(-8+3) - 4(3-4) = 0$$

$$0 = 0 \\ \text{So } \vec{y}, \vec{z} \text{ lie on plane.}$$

$$(b), (c), (d) \text{ Same as Part (a)}$$

$$Q(d) P_0 = (x_0, y_0, z_0) = (5, 2, 1) \\ N = (a, b, c) = (-1, -2, 1)$$

Equation of Plane is

$$a(x-x_0) + b(y-y_0) + c(z-z_0) = 0$$

$$-1(x-5) - 2(y-2) + 1(z-1) = 0$$

Part (a), (b), (c) Same as Part (d)

$$\left(\begin{array}{ccc} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{array} \right)$$

$$Q(b) P_1(2, 3, 2), P_2(-1, -2, 3), P_3(-5, -4, 2)$$

$$\text{Let } U = \overrightarrow{P_1 P_2} = (-3, -5, -1)$$

$$V = \overrightarrow{P_1 P_3} = (-7, -7, -2)$$

$$N_{\text{new}} = \vec{U} \times \vec{V} = \begin{vmatrix} x & y & z \\ -3 & -5 & -1 \\ -7 & -7 & -2 \end{vmatrix}$$

$$\Rightarrow x(10-7) - y(6-7) + z(21-35) = 0 \\ \Rightarrow x + y - 4z = 0$$

Which is the required equation of Plane.
Others part are same as.

Others part are same as.

$$\text{QII (b)} \quad \begin{array}{l} 3x - 2y - 5z + 4 = 0 \\ 2x + 3y + 4z + 8 = 0 \end{array} \quad \left\{ \begin{array}{l} C \\ Q \end{array} \right.$$

$$3x + 3y + 4z + 8 = 0$$

Xing eq (1) by "3" + eq (2) by "2"

$$\begin{array}{l} 9x - 6y - 15z + 12 = 0 \\ 4x + 6y + 8z + 16 = 0 \end{array}$$

$$13x - 7z + 28 = 0$$

$$\frac{13x + 28}{7} = z \quad \text{--- (3)}$$

$$\begin{vmatrix} -2 & 4 & 2 \\ 3 & 5 & 1 \\ 4 & 2 & -1 \end{vmatrix} = 0$$

Xing eq (1) by "2" + eq (2) by "3" + subtracting

$$-2(-5-2) - 4(-3-4) + 2(6-2) = 0$$

$$14 \neq 0$$

These points are not on the same plane.

Q13 is same as Q14.

$$\frac{-13y - 16}{22} = z \quad \text{--- (4)}$$

Comparing eq (3) + eq (4)

$$\frac{13y + 28}{7} = -\frac{13y - 16}{22} = z = t \quad (\text{say})$$

Jai Laser Photostate

Matric to M.A.
Notes are available

Qutub Shabab

Amti: 0345 9219901
Ph: 091-5610987

$$\text{Now } \begin{vmatrix} x_1 & y_1 & z_1 \\ x_2 & y_2 & z_2 \\ x_3 & y_3 & z_3 \end{vmatrix} = 0$$

$$P_1(x_1, y_1, z_1) = (-2, 4, 2)$$

$$P_2(x_2, y_2, z_2) = (3, 5, 1)$$

$$P_3(x_3, y_3, z_3) = (4, 3, -1).$$

(4)

$$\begin{aligned} \text{Q5} \\ X = 2 - 3S & \quad X = S + 2t \\ Y = 3 + 2S & \quad Y = 1 - 3t \\ Z = 4 + 2S & \quad Z = 2 + t \end{aligned}$$

$$\begin{aligned} 2 - 3S &= S + 2t \quad (1) \\ 3 + 2S &= 1 - 3t \quad (2) \\ 4 + 2S &= 2 + t \quad (3) \end{aligned}$$

$$\begin{aligned} \text{Subtracting (3) from eq(2), we get} \\ 3 + 2S &= 1 - 3t \\ \underline{4 + 2S = 2 + t} \\ -1 &= -1 - 4t \\ -4t &= 0 \\ \boxed{t = 0} \end{aligned}$$

$$\text{Also } Z = t \quad (4)$$

Equation (a), (b) & (c) are the required parametric equations of the line of intersection of the given planes.

$$(4) \quad \frac{X-2}{-2} = \frac{Y-3}{4} = \frac{Z+4}{3}$$

$$\text{Now } \frac{X-2}{-2} = \frac{Y-3}{4}$$

$\Rightarrow 4X + 2Y - 14 = 0$ (*) which is equation of plane

$$\text{And } \frac{X-2}{-2} = \frac{Z+4}{3}$$

$3X + 2Z + 2 = 0$ which is a equation of plane.

$$\begin{cases} X = 2 - 3S \\ X = S + 2t \\ X = 5 \end{cases}$$

$$\begin{aligned} \text{Now } \frac{13X + 2S}{7} &= t \\ \Rightarrow X &= \frac{7t - 2S}{13} \rightarrow (4) \end{aligned}$$

$$4 - \frac{13t - 16}{22} = t$$

$$\Rightarrow Y = \frac{22t + 16}{-13} \rightarrow (5)$$

100

$$(i) \quad x = 4t \rightarrow (a)$$

$$y = 1 + 5t \rightarrow (b)$$

$$z = 2 - t \rightarrow (c)$$

$$eq(i) \Rightarrow \frac{x}{4} = t \rightarrow (i')$$

$$eq(ii) \Rightarrow \frac{y-1}{5} = t \rightarrow (ii')$$

$$eq(iii) \Rightarrow z - 2 = t \rightarrow (iii')$$

From eq(i'), (ii') & eq(iii')

$$\frac{x}{4} = \frac{y-1}{5} = z - 2 = t$$

$$Now \quad \frac{y}{4} = \frac{y-1}{5}$$

$$\Rightarrow 5x - 4y + 4 = 0 \quad \text{--- (1) which is 1st eq. of plane.}$$

$$+ \quad y - 1 = 2 - z$$

$$y + 5z - 11 = 0 \quad \text{--- (2) which is 2nd eq. of plane.}$$

Plane.

$$\begin{array}{c} 0 \\ 0 \\ 0 \end{array}$$

Jomial Nasiv
Lekhara P.U.E.T Panhwar

$$y = 3 + 2s$$

$$y = 3 + 2(-1)$$

$$y = 3 - 2 = 1$$

$$\boxed{y = 1}$$

$$y = 1 - 3t$$

$$y = 1 - 3(0)$$

$$y = 1$$

$$\boxed{y = 1}$$

$$z = 2 + 2s$$

$$z = 2 + 0$$

$$z = 2$$

$$\boxed{z = 2}$$

$$z = 2 + t$$

$$z = 2 + 0$$

$$z = 2$$

$$\boxed{z = 2}$$

Therefore the point of intersection of the given line

$$\text{and } P(m, y, z) = P(5, 1, 2)$$

$$\sim \quad \sim \quad \sim \quad \sim \quad \sim$$

$$(a) \quad x = 2 + 2t$$

$$y = 2 + t$$

$$y = -3 - 3t \quad \text{and}$$

$$y = 4 - t$$

$$z = 2 + 2t$$

$$z = 5 - t$$

$$\text{Let } \vec{U} = (2, -3, 4)$$

$$+ \quad \vec{V} = (1, -1, -1)$$

Real Vector Spaces

Now $\vec{U} \cdot \vec{V} = (2, -3, 4) \cdot (1, -1, -1)$.

$$= 2 + 3 - 4 = 1$$

$$\vec{U} \cdot \vec{V} \neq 0$$

The given lines are not \perp .

Note $\vec{U} \cdot \vec{V} = 0$ then \perp

(d) If u and v are any elements of V then $u \oplus v$ is in V (ie V is closed under the operation \oplus).

$$(a) u \oplus v = v \oplus u \text{ for } u \text{ and } v \text{ in } V$$

$$(b) u \oplus (v \oplus w) = (u \oplus v) \oplus w \text{ for } u, v \text{ and } w \text{ in } V.$$

(c) There is an element 0 in V such that

$$u \oplus 0 = 0 \oplus u = u \text{ for all } u \text{ in } V$$

(d) For each u in V there is an element $-u$ in V such that $u \oplus -u = 0$.

- (e) If u is any element of V and c is any real number then $c u$ is in V (ie V is closed under the operation \odot).
- (f) $c(u \oplus v) = c u \oplus c v$, for all real numbers c and all u and v in V .

Shop #53 coffee shop bazar Islamia college Peshawar university



Notes are available
Metric to M.A.

Notes are available
Metric to M.A.

Shop #53 coffee shop bazar Islamia college Peshawar university

Qasir Shahab

0300 56229015

Arif: 0345 9288401
Ph: 091-5610987

Jam Laser Photostate

Arif: 0345 9288401
Ph: 091-5610987

Arif: 0345 9288401
Ph: 091-5610987

Jam Laser Photostate

Arif: 0345 9288401
Ph: 091-5610987

(g) $c \odot (d \oplus u) = (cd) \oplus u$ for all real numbers c and
and all u in V .

(h) $1 \odot u = u$ for all u in V .

The elements of V are called vectors. The real numbers
are called scalars. The operation \oplus is called vector
addition. The operation \odot is called scalar.

Example 2 Consider the set V of all ordered triple of
real numbers of the form (x, y, z) and define the
operations \oplus and \odot by

$$(x, y, z) \oplus (x', y', z') = (x+x', y+y', z)$$

$$c \odot (x, y, z) = (cx, cy, cz)$$

Check the properties to show that it is a vector
space or not.

$$\text{Let } U = (x, y, z) \quad V = (x', y', z')$$

$$(a) U \oplus V = V \oplus U$$

$$\Rightarrow U \oplus V = (x, y, z) \oplus (x', y', z') \\ = (x+x', y+y', z) \rightarrow (i)$$

$$+ V \oplus U = (x', y', z) \oplus (x, y, z) \\ = (x'+x, y'+y, z) \rightarrow (ii)$$

From (i) & (ii) we have

$$U \oplus V = V \oplus U$$

$$(b) U \oplus (V \oplus W) = (U \oplus V) \oplus W. \quad \text{if } W = (a, b, c)$$

$$(U \oplus (V \oplus W)) = U \oplus [(x', y', 0) \oplus (a, b, c)]$$

$$= U \oplus [x+a, y+b, a+c]$$

$$= (x, y, 0) \oplus (x+a, y+b+c)$$

$$= (x+a, y+j+b, c) \quad \text{--- (i)}$$

$$+ (U \oplus V) \oplus W = \{(x, y, 0) \oplus (x', y', 0)\} \oplus (a, b, c)$$

$$= (x+x', y+y', 0) \oplus (a, b, c)$$

$$= (x+x'+a, y+j+b, c) \quad \text{--- (ii)}$$

From eq (i) & eq (ii) we have

$$U \oplus (V \oplus W) = (U \oplus V) \oplus W.$$

$$(c) U \oplus 0 = 0 \oplus U = U.$$

$$U \oplus 0 = (x, y, 0) \oplus 0 = (x+0, y+0, 0+0) = (x, y, 0) = U$$

$$+ 0 \oplus U = (0, 0, 0) \oplus (x, y, 0) = (0+x, 0+y, 0+0) = (x, y, 0) = U$$

$$\text{Hence } U \oplus 0 = 0 \oplus U = U.$$

$$(d) U \oplus -U = 0.$$

$$(x, y, 0) \oplus -(x, y, 0) = (x, y, 0) \oplus (-x, -y, 0)$$

$$(x-x, y-y, 0) = (0, 0, 0) = 0$$

$$\text{Hence } U \oplus -U = 0.$$

Now checking the property for scalar multiplication:

(e) $c \odot (U \oplus V) = c \odot ((x, y, 0) \oplus (x', y', 0))$

$$= c \odot (x+a, y+b, 0)$$

$$= (c(x), c(y), 0) \quad \text{--- (i)}$$

$$+ c \odot U \oplus c \odot V = c \odot (x, y, 0) \oplus c \odot (x', y', 0)$$

$$= (cx, cy, 0) \oplus (cx', cy', 0)$$

$$= (cx+cx', cy+cy', 0) \quad \text{--- (ii)}$$

From eq (i) & eq (ii) we have

$$c \odot (U \oplus V) = c \odot U \oplus c \odot V$$

$$(f) (c+d) \odot U = c \odot U + d \odot U$$

$$(c+d) \odot U = (c+d) \odot (x, y, 0)$$

$$= (cx+dy, cy+dy, 0)$$

$$= (cx+dx, cy+dy, 0) \quad \text{--- (i)}$$

$$+ (c+d) \odot U = c \odot U + d \odot U$$

vis the set of all ordered pairs of real numbers (x, y) where $x > 0$ & $y > 0$

$$(x, y) \oplus (x', y') = (x+x', y+y')$$

$$+ (x, y) = (cx, cy)$$

(g) $\text{co}(\text{dou}) = \text{cdou}$.

$$\begin{aligned}\text{co(dou)} &= \text{co}(\text{do}(x, y, 0)) \\ &= \text{co}(\text{d}(x, dy, 0)) \\ &= (\text{cd}x, \text{cd}y, 0) \quad \text{--- (ii)}\end{aligned}$$

$$\begin{aligned}\text{d(cdou)} &= (\text{cd})\text{o}(\text{co}, y, 0) \\ &= (cdx, cdy, 0) \quad \text{--- (iii)}\end{aligned}$$

From eq (ii) & eq (iii) we have

$$\text{co(dou)} = \text{cdou}$$

$$\text{d. } 1 \odot u = u$$

$$1 \odot (x, y, 0) = (1 \cdot x, 1 \cdot y, 1 \cdot 0)$$

$$= (x, y, 0)$$

$$\text{b) } u \oplus (v \oplus w) = (u \oplus v) \oplus w \text{ let } w = (a, b)$$

Here it satisfies all the properties therefore V is a vector space.

$$= (cx+cdx, cy+cdy, 0) \quad \text{--- (ii)}$$

From eq (ii) & eq (iii) we have

$$(\text{cd})\text{o}u = \text{co}u \oplus \text{dou}$$

Checking the Properties of vector addition

$$\text{u} \oplus \text{v} = (x, y) + \text{v} = (x', y')$$

$$\text{a) } \text{u} \oplus \text{v} = \text{v} \oplus \text{u}$$

$$\begin{aligned}\text{u} \oplus \text{v} &= (x, y) \oplus (x', y') \\ &= (x+x', y+y') \quad \text{--- (i)}\end{aligned}$$

$$\begin{aligned}\text{u} \oplus \text{v} &= (x', y') \oplus (x, y) \quad \text{--- (ii)} \\ &= (x'+x, y'+y) \quad \text{--- (iii)}\end{aligned}$$

From eq (i) & eq (iii) we have

$$\text{u} \oplus \text{v} = \text{v} \oplus \text{u}$$

$$\text{b) } u \oplus (v \oplus w) = (u \oplus v) \oplus w \text{ let } w = (a, b)$$

$$U \oplus (V \otimes W) = U \oplus ((W', j') \otimes (a, b))$$

$$\begin{aligned} &= (W, j) \oplus (W' + a, j' + b) \\ &= (W + W', j + j' + b) - c(i) \end{aligned}$$

$$+ (U \oplus V) \otimes W = ((W, j) \oplus (W', j')) \otimes (a, b)$$

$$\begin{aligned} &= ((W + W', j + j') \oplus (a, b)) \\ &= ((W + W', j + j' + b) - c(i)) \end{aligned}$$

From $\mathfrak{A}_V^{(i)} + \mathfrak{A}_V^{(ii)}$ we have

$$U \oplus (V \otimes W) = U \oplus V + W$$

$$(c) U \oplus 0 = 0 \oplus U = U$$

$$U \oplus 0 = (W, j) \oplus (0, 0) = (W + 0, j + 0) = (W, j) = U$$

$$0 \oplus U = (0, 0) \oplus (W, j) = (0 + W, 0 + j) = (W, j) = U$$

$$\text{Hence } U \oplus 0 = 0 \oplus U = U.$$

$$(d) U \oplus -U = 0$$

$$(W, j) \oplus -(W, j) = (W, j) \oplus (-W, -j)$$

$$(W - W, j - j) = (0, 0) = 0$$

$$\text{Hence } U \oplus -U = 0. \text{ closed under } \oplus$$

New checking the properties for scalar multiplication
 (e) $c \odot (U \otimes V) = c \odot U \otimes c \odot V$

$$c \odot (U \otimes V) = (\odot((W, j)) \odot (W', j'))$$

$$\begin{aligned} &= (\odot(W + W', j + j')) \\ &= c(W + W', c(j + j')) - c \\ &= c(W + W', c(j + j')) - c \end{aligned}$$

$$+ c \odot U \odot c \odot V = c \odot (W', j') \odot (cW, cj)$$

$$= (cW, cj) \odot (cW', cj')$$

$$= (c(W + W'), c(j + j'))$$

$$= (c(W + W'), c(j + j')) - \odot$$

From $\mathfrak{A}_V^{(i)} + \mathfrak{A}_V^{(ii)}$ we have

$$c \odot (U \otimes V) = c \odot U \oplus c \odot V$$

But if $c < 0$ then $(W, j) + (W', j') < 0$ so if

not closed under \odot .

Q2
 V is the set of all ordered triplets of real numbers
 in the form (a, b, c) .

$$(a, b, c) \oplus (a', b', c') = (a+a', b+b', c+c')$$

$$= (a, b+c', b+c', c+c')$$

$$(a, b, c) \odot (a', b', c') = (a, b+a', c+a')$$

$$+ c(a, b, c) = (a, b, c+a).$$

Sol:
 $U = (0, y, z) + V = (0, y', z')$

Checking the properties for vector addition (+)

$$\textcircled{1} \quad U \oplus V = V \oplus U$$

$$\begin{aligned} &= U \oplus V = (0, y, z) \oplus (0, y', z') \\ &= (0+0, y+y', z+z') \\ &= (0, y+y', z+z'). \quad \text{--- (i)} \end{aligned}$$

From eq (i) & eq (iii) we have.

$$U \oplus (V \oplus W) = (U \oplus V) \oplus W$$

$$\begin{aligned} &+ V \oplus W = (0, y', z') \oplus (0, y, z) \\ &= (0+0, y+y', z+z') \\ &= (0, y+y', z+z'). \quad \text{--- (ii)} \end{aligned}$$

$$\text{(c) } U \oplus 0 = 0 \oplus U = U$$

$$\begin{aligned} &U \oplus 0 = (0, y, z) \oplus (0, 0, 0) = (0+0, y+0, z+0) = (0, y, z) = U \\ &0 \oplus U = (0, 0, 0) \oplus (0, y, z) = (0+0, 0+y, 0+z) = (0, y, z) = U \end{aligned}$$

From eq (i) & eq (iii) we have.

$$U \oplus V = V \oplus U.$$

$$\text{Hence } U \oplus 0 = 0 \oplus U = U.$$

$$\begin{aligned} \text{(b) } U \oplus (V \oplus W) &= (U \oplus V) \oplus W + \text{if } W = (a, b, c) \\ &\Rightarrow U \oplus (V \oplus W) = U \oplus [(0, y, z) \oplus (a, b, c)] \\ &= U \oplus (a(0, y, z) + (a, b, c)) \\ &= (0, y, z) \oplus (a, y+b, z+c) \\ &= (a, y+b, z+c) \quad \text{--- (iii)} \end{aligned}$$

$$(c_1\delta) \odot u = c_1 \delta \odot (c_1 y, z)$$

$$= (0, c_1 y, c_1 z) \oplus (0, d_1 y, d_1 z)$$

$$\begin{aligned} &= (0, c_1 y + d_1 y, c_1 z + d_1 z) - 0 \\ &= (0, c_1 y, c_1 z) - 0 \\ &= (0, c_1 y, c_1 z) \end{aligned}$$

$$Hence \quad u \oplus -u = 0$$

closed under \oplus .

Now checking the properties for scalar multiplication

$$\textcircled{2} \quad co(u+v) = (co u) \oplus (co v)$$

$$co(cu+cv) = co((0, y, z) \oplus (0, y', z'))$$

$$= co[0+0, y+y', z+z']$$

$$= co(0, y+y' + z+z')$$

$$= (0, c(cy+y') + c(cz+z')) \quad \text{--- (i)}$$

$$(co(u) \oplus co(v)) = [co((0, y, z)] \oplus [co((0, y', z'))]$$

$$= (0, cy, cz) \oplus (0, cy', cz')$$

$$= (0+0, cy+y', cz+c z')$$

$$= (0, c(cy+y') + c(cz+z')) \quad \text{--- (ii)}$$

$$From eq (i) \oplus eq (ii) we have,$$

$$co(u+v) = (co u) \oplus (co v)$$

$$Hence \quad u \oplus -u = 0$$

Ex 6.1

110

$$(d) 1 \odot U = U$$

$$1 \odot (x, y, z) = (x, y, z) = U$$


How it satisfies all the properties Therefore V is a vector space.

Q3
 V is the set of all polynomials of the form $at^2 + bt + c$ where a, b and c are real numbers with $b=a+1$

$$\begin{aligned} & (a_1t^2 + bt + c_1) \oplus (a_2t^2 + bt + c_2) \\ &= (a_1+a_2)t^2 + (b_1+b_2)t + (c_1+c_2) \end{aligned}$$

$$\text{of } \gamma \odot (at^2 + bt + c) = (\gamma a)t^2 + (\gamma b)t + \gamma c$$

Not closed under \odot b/c we get γ instead of 1.



Hint

$$\text{Let } U = at^2 + bt + c_1 \text{ and } V = at^2 + bt + c_2.$$

$$U = \begin{bmatrix} a_1 & b_1 \\ c_1 & d_1 \end{bmatrix} \quad V = \begin{bmatrix} a_2 & b_2 \\ c_2 & d_2 \end{bmatrix}$$

$$\Rightarrow U \oplus V = (a_1t^2 + bt + c_1) \oplus (a_2t^2 + bt + c_2)$$

$$U \oplus V = (a_1+a_2)t^2 + (b_1+b_2)t + (c_1+c_2)$$

$$\text{Put } b_1 = a_1 + 1, \quad b_2 = a_2 + 1$$

$$\begin{aligned} U \oplus V &= (a_1+a_2)t^2 + (a_1+1+a_2+1)t + (c_1+c_2) \\ &= (a_1+a_2)t^2 + (a_1+a_2+2)t + (c_1+c_2) \end{aligned}$$

Not closed under \odot b/c we get γ instead of 1.
 As given $(at^2 + bt + c) + b = a+1$.

Now showing the properties of scalar multiplication

$$\begin{aligned} \gamma \odot (at^2 + bt + c) &= \gamma at^2 + \gamma bt + \gamma c \\ &= \gamma a_1 t^2 + \gamma (a_1+1)t + \gamma c_1 \\ &= \gamma a_1 t^2 + (\gamma a_1 + \gamma) t + \gamma c_1 \end{aligned}$$

①

Q11, Q12 + Q13 same as Q2.

Q14 same as Q1

Subspaces: Let V be a vector space. Then W will be subspace of V if

- (a) $U \oplus V$ is in W
- (b) $k \in R$, kU is in W .

Q11 Which of the following subsets of R^3 are subspaces of R^3 ? The set of all vectors having the form

(a) $\{a, b, 2\}$

$$\text{Let } \vec{U} = (a, b, 2) \text{ and } \vec{V} = (a', b', 2) \text{ are two vectors}$$

in W then

$$\vec{U} \oplus \vec{V} = (a, b, 2) \oplus (a', b', 2)$$

$$= (a+a', b+b', 4) \notin W$$

which is not in W . Hence W is not a subspace.

(b)

If $\vec{U} = (a, b, c)$ & $\vec{V} = (a', b', c')$ are two vectors in W then $a+b=c$, $a'+b'=c'$

$$P_1 \vec{U} \oplus \vec{V} = (a, b, c) \oplus (a', b', c')$$

$$= (a+a', b+b', c+c') \in W$$

Q12

Q13

Q14

Q15

Q16

Q17

Q18

Q19

Q20

Q21

Q22

Q23

Q24

Q25

Q26

Q27

Q28

Q29

Q30

Q31

Q32

Q33

Q34

Q35

Q36

Q37

Q38

Q39

Q40

Q41

Q42

Q43

Q44

Q45

Q46

Q47

Q48

Q49

Q50

Q51

Q52

Q53

Q54

Q55

Q56

Q57

Q58

Q59

Q60

Q61

Q62

Q63

Q64

Q65

Q66

Q67

Q68

Q69

Q70

Q71

Q72

Q73

Q74

Q75

Q76

Q77

Q78

Q79

Q80

Q81

Q82

Q83

Q84

Q85

Q86

Q87

Q88

Q89

Q90

Q91

Q92

Q93

Q94

Q95

Q96

Q97

Q98

Q99

Q100

Q101

Q102

Q103

Q104

Q105

Q106

Q107

Q108

Q109

Q110

Q111

Q112

Q113

Q114

Q115

Q116

Q117

Q118

Q119

Q120

Q121

Q122

Q123

Q124

Q125

Q126

Q127

Q128

Q129

Q130

Q131

Q132

Q133

Q134

Q135

Q136

Q137

Q138

Q139

Q140

Q141

Q142

Q143

Q144

Q145

Q146

Q147

Q148

Q149

Q150

Q151

Q152

Q153

Q154

Q155

Q156

Q157

Q158

Q159

Q160

Q161

Q162

Q163

Q164

Q165

Q166

Q167

Q168

Q169

Q170

Q171

Q172

Q173

Q174

Q175

Q176

Q177

Q178

Q179

Q180

Q181

Q182

Q183

Q184

Q185

Q186

Q187

Q188

Q189

Q190

Q191

Q192

Q193

Q194

Q195

Q196

Q197

Q198

Q199

Q200

Q201

Q202

Q203

Q204

Q205

Q206

Q207

Q208

Q209

Q210

Q211

Q212

Q213

Q214

Q215

Q216

Q217

Q218

Q219

Q220

Q221

Q222

Q223

Q224

Q225

Q226

Q227

Q228

Q229

Q230

Q231

Q232

Q233

Q234

Q235

Q236

Q237

Q238

Q239

Q240

Q241

Q242

Q243

Q244

Q245

Q246

Q247

Q248

Q249

Q250

Q251

Q252

Q253

Q254

Q255

Q256

Q257

Q258

Q259

Q260

Q261

Q262

Q263

Q264

Q265

Q266

Q267

Q268

Q269

Q270

Q271

Q272

Q273

Q274

Q275

Q276

Q277

Q278

Q279

Q280

Q281

Ex. 2

(a) Let $\vec{U} = (a, b, c)$ & $\vec{V} = (a', b', c')$ in W . Then $a=c=0$, $a'=c'=0$.

$$\vec{U} \oplus \vec{V} = (a+a'+b+b', a+b'+a'+b') \\ = (a+a', b+b', (a+a')+(b+b')) \in W.$$

$$(b) U \oplus V = (a, b, c) \oplus (a', b', c') \\ = (a+a', b+b', c+c')$$

$$= (0, b+b', +0) \in W$$

$$(b) k \circ U = k \circ (a, b, c) \\ = k \circ (a, b, c)$$

$$= ((k \circ a), k \circ b, k \circ c) \in W$$

Hence W is the subspace of R^3 .

$$(b) k \circ U = k \circ (a, b, c)$$

$$= (k \circ a, k \circ b, k \circ c)$$

$$= (k \circ 0, k \circ b, k \circ 0)$$

$$= (0, k \circ b, 0) \in W$$

Hence W is the subspace of R^3 .

$$(b) (a, b, c) \text{ where } a=-c$$

If $\vec{U} = (a, b, c)$ & $\vec{V} = (a', b', c')$ are two vectors in W then $a=-c$, $a'=-c$

$$(a) U \oplus V = (a, b, c) \oplus (a', b', c') \\ = (a+a', b+b', c+c') \\ = (-c+c', b+b', c+c')$$

$$\vec{U} \oplus \vec{V} = (-c + c') + (b + b') + (c + c') \in W$$

$$(b) k\vec{U} = k(a, b, c)$$

$$= kac, kab, kac$$

$$= kac, kab, kac$$

$$= (-kc, kb, kc) \notin W$$

$$k < 0 \& k > 0$$

(c) If $\vec{U} = (a, b, c) \notin W$ and $\vec{V} = (a', b', c') \in W$

$$\text{then } b = 2a + 1, b' = 2a' + 1$$

so W is not a subspace of \mathbb{R}^4 .

$$\vec{U} \oplus \vec{V} = (a, b, c) \oplus (a', b', c')$$

if $a_1 \neq a'_1$, then $a_1 + a'_1 \neq 0$ is similarly for b .

$$\text{if } a_1 = a'_1 + 1 \text{ and } a_0 = 0$$

$$= (a + a'_1, b + b'_1, c + c')$$

$$= (a + a'_1, (2a + 1)(2a'_1 + 1), c + c')$$

$$= (a + a'_1, 2(a + a'_1) + 2, c + c') \notin W$$

So W is not a subspace of \mathbb{R}^4 .

$$(d) k\vec{U} = k(a, b, c, d) \oplus \vec{V} = (a'_1, b'_1, c'_1, d'_1) \in W$$

vector in \mathbb{R}^4 then $a - b = 2 \neq a'_1 - b'_1 = 2$
 $a = 2 + b \neq a'_1 = 2 + b'$

$$(e) \vec{U} \oplus \vec{V} = (a, b, c, d) \oplus (a'_1, b'_1, c'_1, d'_1)$$

$$= (a + a'_1, b + b'_1, c + c'_1, d + d'_1)$$

$$= (b + 2 + b'_1, b + b'_1, c + c'_1, d + d'_1)$$

$$= (b + b'_1 + 4, b + b'_1, c + c'_1, d + d'_1) \notin W$$

$$(f) k\vec{U} = k(a_1t^2 + a_2t + a_0) \oplus \vec{V} = a'_1t^2 + a'_2t + a'_0 \in W$$

$$= (a_1 + a'_1)t^2 + (a_2 + a'_2)t + (a_0 + a'_0) \in W$$

$$k\vec{U} = k(a_1t^2 + a_2t + a_0)$$

$$= (ka_1t^2 + ka_2t + ka_0) \in W \text{ subspace.}$$

$$(b) \quad a_2t^2 + a_1t + a_0 \quad \text{where } a_1 = 2a_0$$

$$\text{Let } \vec{U} = a_2t^2 + a_1t + a_0 \quad \text{and} \quad \vec{V} = a_2't^2 + a_1't + a_0'$$

$$(a) \quad \vec{U} \oplus \vec{V} = (a_2t^2 + a_1t + a_0) \oplus (a_2't^2 + a_1't + a_0')$$

$$= (a_2 + a_2')t^2 + (a_1 + a_1')t + (a_0 + a_0')$$

$$= (a_2 + a_2')t^2 + (a_1 + a_1')t + (a_0 + a_0')$$

$$(b) \quad k \otimes U = k(a_2t^2 + a_1t + a_0)$$

$$= k a_2 t^2 + k a_1 t + k a_0$$

$$= k a_2 t^2 + k a_1 t + k a_0 \in W \quad \text{Subspace.}$$

$$(c) \quad a_2t^2 + a_1t + a_0, \quad \text{where } a_2 + a_1 + a_0 = 2. \Rightarrow a_{2,2} = 2 - a_1 - a_0$$

$$\text{Let } \vec{U} = a_2t^2 + a_1t + a_0 + \vec{V} = a_2't^2 + a_1't + a_0'$$

$$\vec{U} \oplus \vec{V} = (a_2t^2 + a_1t + a_0) \oplus (a_2't^2 + a_1't + a_0')$$

$$= (a_2 + a_2')t^2 + (a_1 + a_1')t + (a_0 + a_0')$$

$$= (2 - a_1 - a_0 + 2 - a_1' - a_0')t^2 + (a_1 + a_1')t + (a_0 + a_0') \notin W$$

not a subspace.

\vec{Q}_9 is similarly to \vec{Q}_8

$$\begin{aligned} & \text{Let } w_1 = a_1u + b_1v \\ & \text{and } w_2 = a_2u + b_2v \end{aligned}$$

$$\begin{aligned} & (a) \quad w_1 \otimes w_2 = (a_1u + b_1v) \otimes (a_2u + b_2v) \\ & = a_1u + a_2u + b_1v + b_2v \end{aligned}$$

$$= (a_{11} + a_{12})u + (b_1 + b_2)v \in W$$

$$(b) \quad k \otimes w_1 = k \otimes (a_1u + b_1v)$$

$$= (ka_1)u + (kb_1)v \in W$$

So W is a Subspace of \mathbb{R}^3 .

\vec{Q}_{15} is as same as \vec{Q}_{14} .

$$\text{Let } \vec{U} = \begin{bmatrix} a & b & c \\ d & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \quad \text{and} \quad \vec{V} = \begin{bmatrix} a_1 & b_1 & c_1 \\ d_1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad b = a + k \\ b_1 = a_1 + k,$$

$$\vec{U} \oplus \vec{V} = \begin{bmatrix} a & b & c \\ d & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \oplus \begin{bmatrix} a_1 & b_1 & c_1 \\ d_1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

$$= \begin{bmatrix} a+a_1 & b+b_1 & c+c_1 \\ d+d_1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

$$U \oplus V = \begin{bmatrix} a+a_1 & (a+c)_1(a_1+c_1) & c+c_1 \\ d+d_1 & 0 & 0 \end{bmatrix} \in W$$

$$U \ominus V = \begin{bmatrix} a & b & c \\ d & e & f \end{bmatrix} \oplus \begin{bmatrix} a & b & c \\ d & e & f \end{bmatrix}$$

$$\text{and } \vec{U} = \begin{bmatrix} a & b & c \\ d & e & f \end{bmatrix} + \vec{V} = \begin{bmatrix} a & b & c \\ d & e & f \end{bmatrix} + \begin{bmatrix} a & b & c \\ d & e & f \end{bmatrix} = \begin{bmatrix} 2a & 2b & 2c \\ 2d & 2e & 2f \end{bmatrix}$$

$$(b) k \otimes U = k \otimes \begin{bmatrix} a & b & c \\ d & e & f \end{bmatrix}$$

$$= k \otimes \begin{bmatrix} a & a+c & c \\ d & 0 & 0 \end{bmatrix}$$

$$= \begin{bmatrix} ka & k(a+b) & kc \\ kd & 0 & 0 \end{bmatrix} \in W$$

Subspace

$$(b) \vec{U} = \begin{bmatrix} a & b & c \\ d & e & f \end{bmatrix} + \vec{V} = \begin{bmatrix} a & b & c \\ d & e & f \end{bmatrix} + C \neq 0.$$

$$(a) \vec{U} \oplus \vec{V} = \begin{bmatrix} a & b & c \\ d & e & f \end{bmatrix} \oplus \begin{bmatrix} a_1 & b_1 & c_1 \\ d_1 & e_1 & f_1 \end{bmatrix}$$

$$= \begin{bmatrix} a+a_1 & b+b_1 & c+c_1 \\ d+d_1 & e+e_1 & f+f_1 \end{bmatrix} \in W$$

$$(b) k \otimes U = k \otimes \begin{bmatrix} a & b & c \\ d & e & f \end{bmatrix} = \begin{bmatrix} ka & kb & kc \\ kd & ke & kf \end{bmatrix}$$

$$= \begin{bmatrix} -2ac & kb & kc \\ kd & ke & kf \end{bmatrix} \in W$$

which is subspace

(17) is similarly Q16

Q16 is subspace

$$= \begin{bmatrix} -2ac & kb & kc \\ kd & ke & kf \end{bmatrix} \in W$$

$$(b) k \otimes U = k \otimes \begin{bmatrix} a & b & c \\ d & e & f \end{bmatrix}$$

$$= \begin{bmatrix} ka & kb & kc \\ kd & ke & kf \end{bmatrix} \in W$$

$$= \begin{bmatrix} -2ac & kb & kc \\ kd & ke & kf \end{bmatrix} \in W$$

Ques (b) $\{C_1, C_2\}$ is same as part (a).

116

Definition - If $S = \{V_1, V_2, \dots, V_k\}$ is a set of vector in a vector space V . Then the set of all vectors in V that are linear combinations of the vectors in S is denoted by $\text{Span } S$ or $\text{span}\{V_1, V_2, \dots, V_k\}$.

Spans or $\text{span}\{V_1, V_2, \dots, V_k\}$

Ans

$$V_1 = (1, 0, 0, 1), V_2 = (1, -1, 0, 0), V_3 = (0, 1, 2, 1)$$

(a) $V = (-1, 4, 2, 2)$.

If $C_1V_1 + C_2V_2 + C_3V_3 = V$ then $C_1, V_2, V_3 \in V$

$$C_1(1, 0, 0, 1) + C_2(1, -1, 0, 0) + C_3(0, 1, 2, 1) = (-1, 4, 2, 2)$$

$$(C_1, 0, 0, C_1) + (C_2, -C_2, 0, 0) + (0, C_3, 2C_3, C_3) = (-1, 4, 2, 2)$$

$$C_1 + C_2 = -1 \quad \text{(i)}$$

$$-C_2 + C_3 = 4 \quad \text{(ii)}$$

$$2C_3 = 2 \quad \text{(iii)}$$

$$C_1 + C_3 = 2 \quad \text{(iv)}$$

$$\stackrel{\text{eqn (iii)}}{\Rightarrow} 2C_3 = 2 \Rightarrow C_3 = 1$$

$$\stackrel{\text{eqn (iv)}}{\Rightarrow} C_1 + C_3 = 2 \Rightarrow C_1 = 1$$

$$\stackrel{\text{eqn (i)}}{\Rightarrow} C_1 + (-C_2 + 1) = -1 \Rightarrow C_2 = 4 - 1 = 3 \Rightarrow C_2 = 3$$

$$\stackrel{\text{eqn (ii)}}{\Rightarrow} C_1 + 1 = 2 \Rightarrow C_1 = 2 - 1 = 1 \Rightarrow C_1 = 1$$

$$\sqrt{0} \Rightarrow C_1 + (-3) = -1 \Rightarrow C_1 = -1 + 3 = 2 \Rightarrow C_1 = 2$$

So solution is not possible. So V is not span of (V_1, V_2, V_3) .

$$A_1 = \begin{bmatrix} 1 & -1 \\ 0 & 3 \end{bmatrix}, A_2 = \begin{bmatrix} 1 & 1 \\ 0 & 2 \end{bmatrix}, A_3 = \begin{bmatrix} 2 & 2 \\ -1 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 5 & 1 \\ -1 & 9 \end{bmatrix} = C_1 \begin{bmatrix} 1 & -1 \\ 0 & 3 \end{bmatrix} + C_2 \begin{bmatrix} 1 & 1 \\ 0 & 2 \end{bmatrix} + C_3 \begin{bmatrix} 2 & 2 \\ -1 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} C_1 - C_1 \\ 0 & 3C_1 \end{bmatrix} + \begin{bmatrix} C_2 & C_2 \\ 0 & 2C_2 \end{bmatrix} + \begin{bmatrix} 2C_3 & 2C_3 \\ -C_3 & C_3 \end{bmatrix}$$

$$\begin{bmatrix} 5 & 1 \\ -1 & 9 \end{bmatrix} = \begin{bmatrix} C_1 + C_2 + 2C_3 & -C_1 + C_2 + 2C_3 \\ 3C_1 + 2C_2 + C_3 \end{bmatrix}$$

$$C_1 + C_2 + 2C_3 = 5 \quad \text{--- (1)}$$

$$C_1 + C_2 + 2C_3 = 1 \quad \text{--- (2)}$$

$$-C_3 = -1 \quad \text{--- (3)}$$

$$3C_1 + 2C_2 + C_3 = 9 \quad \text{--- (4)}$$

$$\stackrel{\text{eqn (2)}}{\Rightarrow} -C_3 = -1 \Rightarrow C_3 = 1$$

$$\stackrel{\text{eqn (3)}}{\Rightarrow} C_1 + C_2 + 2C_3 = 1 \Rightarrow C_1 + C_2 = 3 - 2C_3$$

$$\stackrel{\text{eqn (4)}}{\Rightarrow} C_1 + C_2 + 2C_3 = 9 \Rightarrow C_1 + C_2 = 3 - 2C_3$$

$$\Rightarrow C_1 + C_2 = 3 - 2C_3 \Rightarrow C_1 + C_2 = 3 - 2(-1) \Rightarrow C_1 + C_2 = 5 \quad \text{--- (5)}$$

$$\Rightarrow C_1 + C_2 + 2C_3 = 1 \Rightarrow -C_1 + C_2 = -1 \quad \text{--- (6)}$$

Prob $c_3=1$ in $\text{eq } \textcircled{4}$

$$3c_1+2c_2+1=9$$

$$3c_1+2c_2=8 \quad \textcircled{B}$$

$\textcircled{5} + \textcircled{1}$

$$\begin{array}{l} c_1+c_2=3 \\ -c_1+c_2=-1 \end{array}$$

$$2c_2=2$$

$$\boxed{c_2=1}$$

X'ing eq $\textcircled{5}$ by $\textcircled{3}$ & add with \textcircled{B}

$$\begin{array}{l} -3c_1+3c_2=-3 \\ 3c_1+2c_2=8 \end{array}$$

$$\boxed{c_2=5}$$

$$\boxed{c_2=1}$$

$$3\sqrt{\theta} \Rightarrow 3c_1+2c_1=8$$

$$3c_1=8-2=6$$

$$\boxed{c_1=2}$$

$$c_1=2$$

$$\text{So } \begin{bmatrix} 2 \\ -9 \end{bmatrix} \text{ belongs to space if } c_3=1,$$

Prob (b), (c) & (d) Same as Prob (a).

$$w_1+w_1 = \begin{bmatrix} 0 \\ 0 \end{bmatrix} = w_4$$

$$\begin{aligned} w_3+w_4 &= \begin{bmatrix} 0 \\ 0 \end{bmatrix} = w_2 \\ w_2+w_3 &= \begin{bmatrix} 0 \\ 0 \end{bmatrix} = w_1 \\ w_2+w_4 &= \begin{bmatrix} 0 \\ 0 \end{bmatrix} = w_3 \end{aligned}$$

$$W_1 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}, W_2 = \begin{bmatrix} 0 \\ 0 \end{bmatrix}, W_3 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, W_4 = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$W = \left\{ \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right\}$$

$$W = \left\{ \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right\}$$

bit more

Ex 6.9

117

$$W = \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right\}$$

$$\text{Let } w_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, w_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}, w_3 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, w_4 = \begin{bmatrix} 0 \\ 0 \end{bmatrix}.$$

$$w_1+w_2 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} = w_3$$

$$w_1+w_3 = \begin{bmatrix} 1 \\ 1 \end{bmatrix} = w_4$$

$$w_1+w_4 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} = w_1$$

$$w_2+w_3 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} = w_1$$

$$w_2+w_4 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} = w_2$$

$$w_3+w_4 = \begin{bmatrix} 1 \\ 1 \end{bmatrix} = w_3$$

It's Subspace.

Q5) Let $V = \mathbb{R}^4$. Determine if W , the set of all vectors in V with second entry zero is a subspace of V .

$$W = \left\{ \begin{bmatrix} w_1 & w_2 & w_3 & w_4 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \mid \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} w_1 \\ w_2 \\ w_3 \\ w_4 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} \right\}$$

$$w_1 w_2 = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix} = w_2$$

$$w_1 + w_3 = \begin{bmatrix} 0 \\ 1 \\ 0 \\ 1 \end{bmatrix} = w_3$$

$$w_1 + w_4 = \begin{bmatrix} 0 \\ 1 \\ 1 \\ 0 \end{bmatrix} = w_4$$

$$w_1 + w_5 = \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix} = w_5$$

$$w_1 + w_6 = w_6$$

$$w_1 + w_7 = w_7$$

$$w_1 + w_8 = w_8$$

$$w_1 + w_1 = w_1$$

$$w_2 + w_2 = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} = w_1$$

$$w_3 + w_3 = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} = w_3$$

Similarly other also hold.

Not vector space as $w_1 w_8 = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} \notin W$

$$W = \left\{ \begin{bmatrix} w_1 \\ w_2 \\ w_3 \\ w_4 \end{bmatrix} \mid \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} w_1 \\ w_2 \\ w_3 \\ w_4 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} \right\}$$

Ues if

$$U = c_1 V_1 + c_2 V_2 + c_3 V_3$$

$$\begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} = c_1 \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + c_2 \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix} + c_3 \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} c_1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + \begin{bmatrix} c_2 \\ 1 \\ 0 \\ 0 \end{bmatrix} + \begin{bmatrix} c_3 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} c_1 + c_2 + c_3 \\ 0 \\ c_2 + c_3 \\ c_3 \end{bmatrix}$$

$$\boxed{\text{S.P.} \Rightarrow c_2 + 1 = 0} \quad \text{by 1 mark}$$

$$\boxed{c_2 = -1} \quad \text{on } \boxed{c_2 = 1}$$

$$c_1 + 1 + 1 = 1$$

$$c_1 = 1 - 2 = -1 = 1$$

$$\boxed{c_1 = 1}$$

$$\begin{aligned} c_1 + c_2 + c_3 &= 1 & \text{--- (1)} \\ c_2 + c_3 &= 0 & \text{--- (2)} \\ c_3 &= 1 - c_1 - c_2 & \text{--- (3)} \end{aligned}$$

$$\boxed{c_3 = 1}$$

Which of the following vectors spans \mathbb{R}^2

τ belongs to whom $c_1 = c_2 = c_3 = 1$

Q 33 is same as Q 32.

$\sim \text{O} \sim \text{O}$

Javed Nasir

Lecturer UET Peshawar

Spring Semester & Fall 2013
Electrical Engg.

$$\text{Now } V = C_1 V_1 + C_2 V_2$$

$$(a, b) = C_1 (1, 2) + C_2 (-1, 1)$$

$$(a, b) = (C_1, 2C_1) + (-C_2, C_2)$$

$$(a, b) = (C_1 - C_2, 2C_1 + C_2)$$

$$C_1 - C_2 = a$$

$$2C_1 + C_2 = b$$


C(i) & C(ii)

$$\begin{cases} C_1 - C_2 = a \\ 2C_1 + C_2 = b \end{cases}$$

$$\frac{3C_1 = a+b}{3C_1 = a+b} \Rightarrow C_1 = \frac{a+b}{3}$$

$$C_1 = \frac{a+b}{3} - C_2 = a$$

$$C_2 = \frac{a+b}{3} - a = \frac{a+b-3a}{3} = -\frac{2a+b}{3}$$



Jan Laser Photostate
Matriic to M.A.
Notes are available

Shop #58 coffee shop bazar Islamia college Peshawar university

Qaisr Shahab

Arif: 0345 9289401

Ph: 091-5610987

Qaisr Shahab

Arif: 0345 9289401

Ph: 091-5610987

$$\boxed{C_2 = -\frac{2a+b}{3}}$$

So solution exist & do span \mathbb{R}^2

~~Spanning set~~

$$(b) (a, 0), (1, 1), (-2, -2)$$

$$\text{Let } V = (a, b), \quad V_1 = (0, 0), \quad V_2 = (1, 1), \quad V_3 = (-2, -2)$$

$$\text{Now } V = c_1 V_1 + c_2 V_2 + c_3 V_3$$

$$(a, b) = c_1(0, 0) + c_2(1, 1) + c_3(-2, -2)$$

$$(a, b) = (c_1, 0) + (c_2, c_2) + (-2c_3, -2c_3)$$

$$(a, b) = (0 + c_2 - 2c_3, 0 + c_2 - 2c_3)$$

$$0 + c_2 - 2c_3 = a \quad \text{--- (1)}$$

$$0 + c_2 - 2c_3 = b \quad \text{--- (2)}$$

$$(1) - (2)$$

$$\cancel{0 + c_2 - 2c_3 = a}$$

$$\cancel{0 + c_2 - 2c_3 = b}$$

$$\underline{\underline{0 \neq a - b}}$$

$$c_1 = -\left(\frac{b-a}{2}\right) = \boxed{\frac{c-b}{2} = c_1}$$

Solution doesn't exist so V_1, V_2, V_3 don't span \mathbb{R}^2

$$\text{Part C same part b & (d) same (c)}$$

These different values of c_1 so solution does not exist Nofa span.

Ex 6.3

Which of the following vector span \mathbb{R}^4

- (a)
- (1, 0, 0, 1)
 - (0, 1, 0, 0)
 - (1, 1, 1, 1)
 - (1, 1, 1, 0)

JAN

PHOTOSTAT

FROM MATRIC TO M.A & M.SC
NOTES ARE AVAILABLE

ADDRESS:

JAN PHOTOSTAT, COFFEE SHOP MARKET,
UNIVERSITY OF PESHAWAR

PROPRIETOR:
QAISAR SHABAB, ARIF JAN

Phone No: 091-5610987

Mobile No: 0300-5829015, 0345-9165402
Arif Jan: 0345-9289401

$$\text{Let } V = (a, b, c, d), V_1 = (b, 0, 0, 1), V_2 = (0, b, 0, 0) \quad V_3 = (1, b, 1, 1)$$

$$V_4 = (1, b, 1, 0)$$

$$\text{Now } V = C_1V_1 + C_2V_2 + C_3V_3 + C_4V_4$$

$$(a, b, c, d) = C_1(1, 0, 0, 1) + C_2(0, b, 0, 0) + C_3(1, b, 1, 1) + C_4(1, b, 1, 0)$$

$$(a, b, c, d) = (C_1, 0, 0, C_1) + (0, C_2, 0, 0) + (C_3, C_3, C_3, C_3) + (C_4, C_4, C_4, C_4)$$

$$(a, b, c, d) = (C_1 + C_3 + C_4, C_2 + C_3 + C_4, C_3 + C_4, C_1 + C_3)$$

$$C_1 + C_3 + C_4 = a \quad (1)$$

$$C_2 + C_3 + C_4 = b \quad (2)$$

$$C_3 + C_4 = c \quad (3)$$

$$C_1 + C_3 = d \quad (4)$$

$$P.S.C.H.S.D. in $\mathbb{R}^4$$$

$$a \neq 0 \Rightarrow C_4 \neq d \Rightarrow a = 0 \Rightarrow \boxed{C_4 = 0 - d}$$

$$a \neq 0 \Rightarrow C_3 + (a - d) = C \Rightarrow \boxed{C_3 = C - a + d}$$

$$a \neq 0 \Rightarrow C_1 + (C - a + d) = d \Rightarrow C_1 = d - (C - a + d)$$

$$a \neq 0 \Rightarrow C_1 + (C - a + d) = d \Rightarrow C_1 = d - C + a - d = -C + a$$

$$\text{Q} \Rightarrow C_1 = a - c$$

$$\text{Q} \Rightarrow a - C_2 + C_3 = c$$

$$-C_2 + C_3 = c - a \quad \text{--- (4)}$$

$$\text{Q} + \text{4}$$

$$\frac{C_3 + C_2 = b}{C_3 - C_2 = c - a}$$

$$2C_3 = c - a + b$$

$$C_3 = \frac{c-a+b}{2}$$

$$\begin{aligned} & \text{Q} \\ & \text{Q} \Rightarrow C_2 + (C_{-a+d}) + (a-d) = b \\ & C_2 + C_{-d+d} + a - d = b \\ & C_2 = b - c \end{aligned}$$

$$\text{Hence solution exist so spans } R^4$$

Other parts are same as part a

$$(a)f^2t^2, t^2t, t+1$$

$$\text{Pf } P_1(a, b, c), \quad P_1 = (t^2 + 1), \quad P_2 = (t^2 + t), \quad P_3 = (t + 1)$$

$$P = C_1P_1 + C_2P_2 + C_3P_3$$

$$(a, b, c) = C_1(t^2 + 1) + C_2(t^2 + t) + C_3(t + 1)$$

$$= (C_1t^2 + C_1) + (C_2t^2 + C_2t) + (C_3t + C_3)$$

= ~~similarly~~

$$(a, b, c) = (C_1 + C_2)t^2 + (C_2 + C_3)t + (C_1 + C_3)$$

$$C_1 + C_2 = a - 0$$

$$C_3 + C_2 = b - Q$$

$$C_1 + C_3 = c - Q$$

$$C_1 = \frac{a-b+c}{2}$$

$$\begin{aligned} & \text{Q} \Rightarrow C_1 = a - C_2 \\ & = a - \left(\frac{b+a-c}{2} \right) = \frac{2a-b-c}{2} \end{aligned}$$

$$C_1 = a - C$$

So the solution exist hence a poly no: $\text{Span } P_2$

Q_5 is same as Q_4 .

Q6
Augmented form

$$\left[\begin{array}{cccc|c} 1 & 0 & 1 & 0 & 0 \\ -2 & 3 & 1 & 0 & 0 \\ -1 & 2 & 1 & 0 & 0 \\ -1 & 1 & 0 & 0 & 0 \end{array} \right]$$

$$\left[\begin{array}{cccc|c} 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & -2 & 1 & 0 \\ 0 & 0 & -1 & 1 & 0 \\ 1 & 1 & 0 & 0 & 0 \end{array} \right] R_2-R_1$$

$$\left[\begin{array}{cccc|c} 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 \end{array} \right] R_3-2R_1$$

$$\left[\begin{array}{cccc|c} 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 \end{array} \right] R_3-R_1$$

$$\left[\begin{array}{cccc|c} 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 \end{array} \right] R_4-R_1$$

$c_1 + c_3 = 0$

$c_1 = -c_3$

$c_1 = -\gamma$

So Span exist has infinite many solns.

$Q7$ is same $\underline{\underline{Q6}}$

$$X_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \quad X_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \quad X_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix}, \quad X_4 = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}$$

$X = c_1 X_1 + c_2 X_2 + c_3 X_3$

$$\begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} \pm c_1 \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + c_2 \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix} + c_3 \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix}.$$

$$\begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} c_1 + c_2 + c_3 \\ 2c_1 + 0 + 0 \\ 0 - c_2 + 2c_3 \\ c_1 + c_2 + 0 \end{bmatrix}$$

$$2(c_2 + 2c_3 + c_4) = 0$$

$2(c_2 + 2c_3) = 0$

$2(c_2 + 2c_3 + 2c_4) = 0$

$$\begin{aligned} C_1 + C_2 + C_3 &= 0 \quad (1) \\ 2C_1 + 6C_3 &= 0 \quad (2) \\ -C_2 + 2C_3 &= 0 \quad (3) \\ C_1 + C_2 &= 0 \quad (4) \end{aligned}$$

$$\begin{aligned} C_1 + C_2 + C_3 &= 0 \quad (1) \\ 2C_1 + 6C_3 &= 0 \quad (2) \\ -C_2 + 2C_3 &= 0 \quad (3) \\ C_1 + C_2 &= 0 \quad (4) \end{aligned}$$

Put $C_1 + C_2$ value in eq(1)

$$eq(4) \Rightarrow C_1 + C_2 = 0 \Rightarrow \boxed{C_3 = 0}$$

$$eq(3) \Rightarrow -C_2 + 2C_3 = 0 \Rightarrow \boxed{C_2 = 0}$$

$$eq(4) \Rightarrow C_1 + 0 = 0 \Rightarrow \boxed{C_1 = 0}$$

Hence solution exist so $\{x_1, x_2, x_3\}$ is L.I.

θ_3 is same as θ_4 .

$$\begin{array}{c} X \\ \hline \end{array}$$

$$V_1 = (1, 2, -1) \quad V_2 = (3, 2, 5)$$

$$Now C_1 V_1 + C_2 V_2 = 0$$

$$C_1(1, 2, -1) + C_2(3, 2, 5) = 0$$

$$C_1 + 3C_2 = 0 \quad (1)$$

$$2C_1 + 2C_2 = 0 \quad (2)$$

$$-C_1 + 5C_2 = 0 \quad (3)$$

$$\begin{aligned} (1) - (3) \\ C_1 + 3C_2 = 0 \\ -C_1 + 5C_2 = 0 \\ \hline 8C_2 = 0 \\ C_2 = 0 \end{aligned}$$

$$2\sqrt{0} \Rightarrow C_1 + 3C_2 = 0 \Rightarrow \boxed{C_1 = 0}$$

Hence linearly independent.

(b), (c), (d) is similarly.

θ_{11} is similarly to θ_{10}

$$\begin{array}{c} X \\ \hline \end{array}$$

$$\theta_2 = t^2 + 1, \quad \theta_3 = t^{-2}, \quad \theta_4 = t + 3.$$

$$Now C_1 \theta_1 + C_2 \theta_2 + C_3 \theta_3 = 0$$

$$C_1(t^2 + 1) + C_2(t^{-2}) + C_3(t + 3) = 0$$

$$(C_1 t^2 + C_1) + (C_2 t^{-2}) + (C_3 t + C_3) = 0$$

$$(C_1) t^2 + (C_2 + C_3) t + (C_1 - C_2 + C_3) = 0$$

$$C_1 = 0 \quad \boxed{1} \Rightarrow \boxed{C_1 = 0}$$

$$C_2 + C_3 = 0 \quad \boxed{2}$$

$$C_1 - C_2 + C_3 = 0 \quad \boxed{3}$$

$P_1 + C_2 = 0$ in eq ②

$$2P_2 \Rightarrow 0 - C_2 + C_3 = 0$$

$$-C_2 + C_3 = 0 \quad \text{---} \quad ③$$

② + ④

$$\begin{array}{l} C_1 + C_3 = 0 \\ -C_2 + C_3 = 0 \end{array}$$

$$2C_3 = 0$$

$$2P_2 \Rightarrow C_2 + 0 = 0$$

$$\boxed{C_2 = 0}$$

$C_1 = 0, C_2 = 0, C_3 = 0$ so linearly independent.

Part b, c, & d is similarly

$$\left(\begin{array}{rrrr} 1 & 2 & 3 & 2 \\ 1 & 3 & 1 & 0 \\ -1 & 2 & 1 & 0 \\ 2 & 1 & 1 & 0 \end{array} \right)$$

$$C_1 + 2C_2 + 3C_3 + 2C_4 = 0 \quad ①$$

$$C_1 + 3C_2 + C_3 + 2C_4 = 0 \quad ②$$

$$C_1 + C_2 + 2C_3 + C_4 = 0 \quad ③$$

$$C_1 + 2C_2 + C_3 + C_4 = 0 \quad ④$$

Now $C_1V_1 + C_2V_2 + C_3V_3 + C_4V_4 = 0$

$$\left\{ \left[\begin{array}{r} 1 \\ 1 \\ 1 \\ 1 \end{array} \right], \left[\begin{array}{r} 2 \\ 3 \\ 1 \\ 0 \end{array} \right], \left[\begin{array}{r} 3 \\ 1 \\ 2 \\ 0 \end{array} \right], \left[\begin{array}{r} 1 \\ 2 \\ 1 \\ 0 \end{array} \right] \right\}$$

Exercise 10

Consider the vector space $M_{2,2}$. Follow the direction of

Ex 6.3

Daur Shahab 0300 5829015 Jam Laser Photostate Matrix to M.A. Notes are available 	Aarif 0345 9289401 Ph: 091-5610987 Daur Shahab 0300 5829015 Jam Laser Photostate Matrix to M.A. Notes are available 
--	---

$$\left(\begin{array}{rrrr} 1 & 2 & 3 & 2 \\ 0 & 1 & -2 & 0 \\ 0 & -1 & -1 & 1 \\ 0 & 0 & -2 & 1 \end{array} \right) \xrightarrow{\begin{array}{l} R_2 - R_1 \\ R_3 - R_1 \\ R_4 - R_1 \end{array}} \left(\begin{array}{rrrr} 1 & 2 & 3 & 2 \\ 0 & 1 & -2 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -2 & 1 \end{array} \right) \xrightarrow{\begin{array}{l} R_1 + R_2 \\ R_3 + R_2 \end{array}}$$

③

126

(a) $\{ \cos t, \sin t, e^t \}$

$$V_1 = \cos t, V_2 = \sin t, V_3 = e^t$$

$$\text{Now } C_1 V_1 + C_2 V_2 + C_3 V_3 = 0$$

$$C_1 = C_2 = C_3 = 0 \quad \text{Linearly Independent.}$$

$$\begin{aligned} & \left[\begin{array}{cccc|c} 1 & 2 & 3 & 2 & 1 & 0 \\ 0 & 1 & -2 & 0 & 1 & 0 \\ 0 & 0 & 1 & \sqrt{3} & 0 & 0 \\ 0 & 0 & -2 & -1 & 0 & 0 \end{array} \right] \sim \\ & \left[\begin{array}{cccc|c} 1 & 2 & 3 & 2 & 1 & 0 \\ 0 & 1 & -2 & 0 & 1 & 0 \\ 0 & 0 & 1 & \sqrt{3} & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 \end{array} \right] \sim R_3, R_4 \\ & -\sqrt{3}C_4 = \boxed{C_4=0} \end{aligned}$$

$$C_3 + \boxed{C_3=0} = 0$$

$$C_3 + \boxed{C_3=0} = 0$$

$$C_3 + \boxed{C_3=0} = 0$$

$$C_2 + \boxed{C_2=0} = 0$$

$$C_2 + \boxed{C_2=0} = 0$$

$$\boxed{C_2=0}$$

$$C_1 + 2C_2 + 3C_3 + 2C_4 = 0$$

$$C_1 + 2(\boxed{C_2=0}) + 3(C_3) + 2(C_4) = 0$$

$$\boxed{C_1=0}$$

$$C_1 = C_2 = C_3 = C_4 = 0$$

So it's linearly independent.

(d) $\{\cos t, \sin t, e^{2t}\}$

$$V_1 = \cos t, V_2 = \sin t, V_3 = e^{2t} = e^{2t} - e^{2t}$$

$$\text{Now } C_1 V_1 + C_2 V_2 + C_3 V_3 = 0$$

$$C_1 \cos t + C_2 \sin t + C_3 (e^{2t} - e^{2t}) = 0$$

$$(C_1 + C_3) \cos t + (C_2 - C_3) e^{2t} = 0$$

$$C_1 + C_3 = 0 \quad \text{---} \quad 0$$

$$C_2 - C_3 = 0 \quad \text{---} \quad 0$$

Let $\boxed{c_3 = \gamma}$

$$\text{eq } Q \Rightarrow c_2 - \gamma = 0 \Rightarrow \boxed{c_2 = +\gamma}$$

$$\text{eq } P \Rightarrow c_1 + \gamma = 0 \Rightarrow \boxed{c_1 = -\gamma}.$$

Which is linearly dependent on non-trivial.

or

$$\begin{array}{l} \text{Let} \\ V_1 = (1, 0, -1) \end{array}$$

$$V_2 = (2, 1, 2)$$

$$V_3 = (1, 1, \epsilon)$$

$$\text{Now } C_1 V_1 + C_2 V_2 + C_3 V_3 = 0$$

$$C_1(-1, 0, -1) + C_2(2, 1, 2) + C_3(1, 1, \epsilon) = 0$$

$$-C_1 + 2C_2 + C_3 = 0 \quad \textcircled{1}$$

$$+2C_2 + C_3 = 0 \quad \textcircled{2}$$

$$-C_1 + 2C_2 + C_3 = 0 \quad \textcircled{3}$$

Given L. Dependent so the solution is non-trivial

then $\det = 0$

$$\begin{vmatrix} -1 & 2 & 1 \\ 0 & 1 & 1 \\ -1 & 2 & \epsilon \end{vmatrix} = 0$$

Q6

$$\text{Let } V_1 = t + 3$$

$$V_2 = 2t + \lambda^2 + 2$$

$$\text{Now } C_1 V_1 + C_2 V_2 = 0.$$

$$C_1(t+3) + C_2(2t + \lambda^2 + 2) = 0$$

$$C_1 t + 3C_1 + 2C_2 t + C_2 \lambda^2 + 2C_2 = 0$$

$$C_1 t + 2C_2 t = 0 \quad \textcircled{1}$$

$$3C_1 + C_2(\lambda^2 + 2) = 0 \quad \textcircled{2}$$

Det = 0 if non-trivial solution

$$\lambda^2 + 2 = 0 \Rightarrow \lambda = \pm 2i$$

$$\begin{vmatrix} 1 & 2 \\ 3 & \lambda^2 + 2 \end{vmatrix} = 0 \quad \lambda^2 + 2 = 0$$

$$\begin{vmatrix} -1 & 1 & 1 \\ 2 & 2 & -2 \\ 0 & 1 & -1 \end{vmatrix} \begin{vmatrix} 0 & 1 \\ -1 & \epsilon \end{vmatrix} = 0$$

$$-1(C-2) - 2(0+\epsilon) + 1(0+\epsilon) = 0$$

$$-C + 2 - 2\epsilon + \epsilon = 0$$

$$-C + 2 - 2\epsilon + \epsilon = 0$$

$$-C + 2 = 0$$

$$-C + 2 = 0$$

$$\Rightarrow \boxed{C=2}$$

$$\boxed{Q17} \quad V_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad V_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad V_3 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \quad V = \begin{bmatrix} a \\ b \end{bmatrix}$$

$$C_1 V_1 + C_2 V_2 + C_3 V_3 = V$$

$$C_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + C_2 \begin{bmatrix} 0 \\ 1 \end{bmatrix} + C_3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} a \\ b \end{bmatrix}$$

$$\begin{bmatrix} a \\ 0 \end{bmatrix} + \begin{bmatrix} 0 \\ C_2 \end{bmatrix} + \begin{bmatrix} C_3 \\ C_3 \end{bmatrix} = \begin{bmatrix} a \\ b \end{bmatrix}$$

$$\begin{bmatrix} a \\ 0 \end{bmatrix} + \begin{bmatrix} 0 \\ C_2 \end{bmatrix} + \begin{bmatrix} 0 \\ C_3 \end{bmatrix} = \begin{bmatrix} a \\ b \end{bmatrix}$$

$$\begin{array}{l} C_1 + C_3 = a \\ 0 + C_2 = b \\ 0 + C_2 + C_3 = b \end{array}$$

$$\textcircled{1} - \textcircled{2}$$

$$C_1 + C_3 = a$$

$$\begin{array}{l} C_1 + C_3 = a \\ 0 + C_2 = b \\ 0 + C_2 + C_3 = b \end{array}$$

$$\textcircled{1} - \textcircled{2}$$

$$C_1 + C_3 = a$$

$$0 + C_2 + C_3 = b$$

$$\frac{C_1 = a - c}{C_1 = a_v - c} \Rightarrow \boxed{C_1 = a_v - c} \Rightarrow \boxed{C_1 = a + c}$$

$$a + c + a_v + c_3 = a$$

$$(1) \quad a + c_3 = a + c + b = 0(a) + c_3 = a + c + b \quad \text{For B.M}$$

$$c_3 = a + c + b - \boxed{a} = \boxed{c + b + a_v = c_3}$$

$$\begin{aligned} C_3 &= a + c + b \\ C_3 &= C_1 + b \\ C_3 - C_1 &\neq b \end{aligned}$$

$$C_3 + C_1 = b \quad (\text{In B.M})$$

$$C + b + a_v + a + c = b$$

$$C(1+1) + (1+1)a + b = b$$

$$C(a) + (a)a + b = b$$

$$0 + 0 + b = b$$

$$b = b \text{ verified.}$$

So 2nd soln B³

Ques same w Q17.

BQ is same.

Q2D is same w B3

Q2L is similarly

Q2A is

\sim

JAN

PHOTOSTAT

FROM MATRIC TO M.A & M.SC
NOTES ARE AVAILABLE

ADDRESS:

JAN PHOTOSTAT, COFFEE SHOP MARKET,
UNIVERSITY OF PESHAWAR

PROPRIETOR:
QAISAR SHABAB, ARIF JAN

Phone No: 091-5610987

Mobile No: 0300-5829015, 0345-9165401

Q = { $\vec{v}_1, \vec{v}_2, \vec{v}_3, \vec{v}_4, \vec{v}_5$ } w.r.t.
 $\vec{v}_1 = (1, 2, 3)$,
 $\vec{v}_2 = (2, 1, 4)$,
 $\vec{v}_3 = (-1, -1, 2)$,
 $\vec{v}_4 = (0, 1, 2)$, & $\vec{v}_5 = (1, 1, 1)$

Note that V is the Null Space of Matrix A whose rows are the given vectors

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 1 & 4 \\ -1 & -1 & 2 \\ 0 & 1 & 2 \\ 1 & 1 & 1 \end{bmatrix}$$

$$\sim \left[\begin{array}{ccc|cc} 1 & 2 & 3 & -2R_1+R_2 & R_1+R_3 \\ 0 & 0 & 0 & -1 & -R_1+R_5 \\ 0 & 1 & 2 & 1 & -R_1+R_5 \end{array} \right]$$

$$\left[\begin{array}{ccc|cc} 1 & 2 & 3 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 2 & 0 & 0 \end{array} \right]$$

\mathbf{Q}_2 is similar to \mathbf{Q}_1

130

$$\begin{bmatrix} ? & ? & ? \\ 0 & 0 & 0 \\ 1 & -1 & 1 \end{bmatrix} \begin{array}{l} -2R_2 + R_1 \\ 2R_4 + R_3 \\ R_4, R_5 \end{array}$$

$$\begin{bmatrix} ? & ? & ? \\ 0 & 0 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{array}{l} -R_3 + R_4 \\ R_4, R_5 \end{array}$$

$$\begin{bmatrix} ? & ? & ? \\ 0 & 0 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{array}{l} -7 \\ 5 \\ 4 \end{array}$$

$$\begin{bmatrix} ? & ? & ? \\ 0 & 0 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{array}{l} -R_3 + R_4 \\ R_4, R_5 \end{array}$$

$$\begin{bmatrix} ? & ? & ? \\ 0 & 0 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{array}{l} -SR_3 + R_2 \\ 7R_3 + R_1 \\ 3R_3 + R_4 \end{array}$$

$$R_3/4$$

which is reduced now echelon form.

Note here $w_1 = (1, 0, 0)$
 $w_2 = (0, 1, 0)$
 $w_3 = (0, 0, 1)$ from a basis of V .

$$\begin{bmatrix} ? & ? & ? \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{array}{l} -2R_2 + R_1 \\ 4R_2 + R_3 \\ 3R_2 + R_4 \\ 7R_2 + R_5 \end{array}$$

$$\begin{bmatrix} ? & ? & ? \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{array}{l} -2R_2 + R_1 \\ 4R_2 + R_3 \\ 3R_2 + R_4 \\ 7R_2 + R_5 \end{array}$$

$$\begin{bmatrix} ? & ? & ? \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{array}{l} -2R_1 + R_2 \\ 3R_1 + R_3 \\ -3R_1 + R_4 \\ 5R_1 + R_5 \end{array}$$

$$\text{Now } A = \begin{bmatrix} 1 & 2 & 1 \\ 3 & 2 & 1 \\ 3 & 2 & 1 \\ 3 & 2 & 1 \end{bmatrix}$$

$$V_1 = \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix} \quad V_2 = \begin{bmatrix} 3 \\ 2 \\ 2 \end{bmatrix} \quad V_3 = \begin{bmatrix} 3 \\ 2 \\ 2 \end{bmatrix} \quad V_4 = \begin{bmatrix} 3 \\ 2 \\ 2 \end{bmatrix} \quad V_5 = \begin{bmatrix} 3 \\ 2 \\ 2 \end{bmatrix}$$

$$\begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}$$

Scanned by CamScanner

Linearly independent solution $\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \right\}$

IIQ

$$A = \begin{bmatrix} -1 & 2 & -1 \\ 2 & -1 & 3 \\ 3 & 2 & -1 \\ 0 & 0 & 4 \\ -1 & -1 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 2 \\ 0 & 0 & 0 & 0 & 3 \\ 0 & 0 & 0 & 0 & 4 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\begin{array}{l} R_1 - 2R_1 \\ R_4 - 3R_1 \\ R_5 - 5R_1 \end{array}$$

$$\begin{bmatrix} 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 2 \\ 0 & 0 & 0 & 0 & 3 \\ 0 & 0 & 0 & 0 & 4 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix} - (R_2 + R_3)$$

$$\begin{bmatrix} 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 \\ -25 & -10 & -2 & -2 & 0 \end{bmatrix}$$

$R_{3/5}$

$$\begin{bmatrix} 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 \\ -25 & -10 & -2 & -2 & 0 \end{bmatrix}$$

$$\begin{array}{l} R_1 - 2R_2 \\ R_3 - 2R_2 \\ R_4 - 4R_2 \\ R_5 + 10R_1 \end{array}$$

$$\begin{bmatrix} 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\begin{array}{l} R_5 - 4R_4 \\ R_3 - 6R_4 \\ R_2 - 2R_4 \\ R_1 + R_4 \end{array}$$

$$\begin{bmatrix} 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\begin{array}{l} V_5 R_4 \\ R_4 \end{array}$$

$$\begin{bmatrix} 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\begin{array}{l} R_4 + 6R_2 \\ R_5 + 5R_3 \\ R_2 + 2R_3 \\ R_1 - 5R_3 \end{array}$$

$$\therefore w_1 = (1, 0, 0, 0)$$

$$w_2 = (0, 1, 0, 0)$$

$$w_3 = (0, 0, 1, 0)$$

$$w_4 = (0, 0, 0, 1) \quad \text{form a basis for } V$$

Jan Laser Photostate

Qais Shabab

0300 5829015

Aasif 0345 02289401
Ph: 091-3610987

Matrix to M.A.
Notes are available



Shop#58 coffee shop bazar Islamia college Peshawar University

Ex 6

whose augmented matrix is

$$\left[\begin{array}{ccccc} 1 & -1 & -3 & -2 & 1 \\ 2 & 9 & 8 & 3 & 0 \\ -1 & -1 & 3 & 2 & 0 \end{array} \right] = [A^T : 0]$$

In the coefficient matrix is A^T . Transforming the augmented matrix $[A^T : 0]$ is to reduced row echelon form we obtain

$$\left[\begin{array}{ccccc} 1 & -1 & -3 & -2 & 0 \\ 0 & 7 & 14 & 7 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right] \xrightarrow{R_2 \leftrightarrow R_3}$$

$$\left[\begin{array}{ccccc} 1 & 2 & -1 & 0 & 0 \\ 0 & 7 & 0 & 0 & 0 \\ 0 & 14 & 0 & 0 & 0 \end{array} \right] \xrightarrow{R_1 + R_2}$$

$$\left[\begin{array}{ccccc} 1 & 2 & -1 & 0 & 0 \\ 0 & 7 & 0 & 0 & 0 \\ 0 & 14 & 0 & 0 & 0 \end{array} \right] \xrightarrow{R_2 \leftrightarrow R_3}$$

$$\left[\begin{array}{ccccc} 1 & 2 & -1 & 0 & 0 \\ 0 & 7 & 0 & 0 & 0 \\ 0 & 14 & 0 & 0 & 0 \end{array} \right] \xrightarrow{R_1 + R_2}$$

$$\text{Hence } w_1 = (1, 0, -1)$$

$w_2 = (0, 1, 0)$ is basis for A .

Ans.

(b) consisting of vector that are new vectors of A

Q5 (a) consisting of vector that are new vectors of A

$$A = \left[\begin{array}{ccc} 1 & 2 & -1 \\ 1 & 9 & -1 \\ -3 & 8 & 3 \\ -2 & 3 & 2 \end{array} \right]$$

$$\left[\begin{array}{ccccc} 1 & 2 & -1 & 0 & 0 \\ 0 & 7 & 0 & 0 & 0 \\ 0 & 14 & 0 & 0 & 0 \end{array} \right] \xrightarrow{R_1 + R_2}$$

$$\left[\begin{array}{ccccc} 1 & 2 & -1 & 0 & 0 \\ 0 & 7 & 0 & 0 & 0 \\ 0 & 14 & 0 & 0 & 0 \end{array} \right] \xrightarrow{R_2 \leftrightarrow R_3}$$

$$\left[\begin{array}{ccccc} 1 & 2 & -1 & 0 & 0 \\ 0 & 7 & 0 & 0 & 0 \\ 0 & 14 & 0 & 0 & 0 \end{array} \right] \xrightarrow{R_1 + R_2}$$

$$\text{Hence } w_1 = (1, 0, -1)$$

$w_2 = (0, 1, 0)$ is basis for A .

Ans.

(b) consisting of vector that are new vectors of A

Since the leading 1's in column 1, 2, we conclude that the 1st two rows of A form a basis for the new space A that is $\{(1, 2, -1), (1, 9, -1)\}$.

\mathcal{Q}_6 is similar to \mathcal{Q}_5 .

$$\begin{bmatrix} 1 & -2 & 7 & 0 \\ 1 & -1 & 4 & 0 \\ 3 & -2 & -3 & 5 \\ 2 & -1 & 3 & 5 \end{bmatrix} \xrightarrow{\text{R}_1 + R_2} \begin{bmatrix} 2 & -1 & 0 & 0 \\ 0 & 0 & 0 & 5 \\ 0 & 0 & 0 & 0 \\ 0 & -3 & 5 & 5 \end{bmatrix} \xrightarrow{\text{S.R}_4 + R_1} \begin{bmatrix} 2 & -1 & 0 & 0 \\ 0 & 0 & 0 & 5 \\ 0 & 0 & 0 & 0 \\ 0 & -3 & 5 & 5 \end{bmatrix}$$

Ex 6.6
133

(a)

$$\begin{bmatrix} 1 & -1 & 3 & 2 \\ -2 & -1 & 2 & -1 \\ 0 & 4 & -3 & 1 \\ 0 & 5 & 3 & -1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -1 & 3 & 2 \\ 0 & 4 & -3 & 1 \\ 0 & 5 & 3 & -1 \\ 0 & 5 & 3 & -1 \end{bmatrix} \xrightarrow{2R_1 + R_2} \begin{bmatrix} 1 & -1 & 3 & 2 \\ 0 & 4 & -3 & 1 \\ 0 & 5 & 3 & -1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -1 & 3 & 2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \xrightarrow{-R_1 + R_2} \begin{bmatrix} 1 & -1 & 3 & 2 \\ 0 & 4 & -3 & 1 \\ 0 & 5 & 3 & -1 \\ 0 & 5 & 3 & -1 \end{bmatrix} \xrightarrow{2R_1 + R_2} \begin{bmatrix} 1 & -1 & 3 & 2 \\ 0 & 4 & -3 & 1 \\ 0 & 5 & 3 & -1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

i) what augmented matrix is
 $\begin{bmatrix} 1 & -2 & 7 & 0 & 0 \\ -1 & 4 & 0 & 0 & 0 \\ 3 & -2 & 1 & 0 & 0 \\ 2 & -1 & 3 & 0 & 0 \end{bmatrix} = \begin{bmatrix} A & | & b \end{bmatrix}$
 i.e. the coefficient matrix is \bar{A} . Transforming the augmented matrix $\begin{bmatrix} \bar{A} & | & b \end{bmatrix}$ to reduced row echelon form we obtain

$$\begin{bmatrix} 1 & -2 & 7 & 0 & 0 \\ -1 & 4 & 0 & 0 & 0 \\ 3 & -2 & 1 & 0 & 0 \\ 2 & -1 & 3 & 0 & 0 \end{bmatrix} \xrightarrow{-R_1 + R_2} \begin{bmatrix} 1 & -2 & 7 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 3 & -2 & 1 & 0 & 0 \\ 2 & -1 & 3 & 0 & 0 \end{bmatrix} \xrightarrow{3R_2 + R_3} \begin{bmatrix} 1 & -2 & 7 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 2 & -1 & 3 & 0 & 0 \end{bmatrix} \xrightarrow{2R_4 + R_1} \begin{bmatrix} 1 & -2 & 7 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \xrightarrow{-8R_2 + R_1} \begin{bmatrix} 1 & -2 & 7 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$\begin{bmatrix} 1 & -2 & 7 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$ is the basis for the column space A.

$$\lambda = \begin{bmatrix} 1 & -2 & 5 \\ 2 & 3 & 2 \\ 0 & 7 & 8 \end{bmatrix}$$

134

(a) basis for null space of A

$$\begin{array}{c} \left[\begin{array}{ccc|c} 1 & -2 & 5 & 0 \\ 0 & -7 & 8 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] R_{2,2R_1} \\ \hline \left[\begin{array}{ccc|c} 1 & -2 & 5 & 0 \\ 0 & 1 & -8/7 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \end{array}$$

$$\begin{array}{c} \left[\begin{array}{cccc|c} 1 & 0 & 1 & 0 & 10 \\ 0 & 1 & -3 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \end{array} \right] R_3R_4 \\ \hline \left[\begin{array}{cccc|c} 1 & 0 & 1 & 0 & 10 \\ 0 & 1 & -3 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \end{array} \right] -5R_4 + R_2 \\ \hline \left[\begin{array}{cccc|c} 1 & 0 & 1 & 0 & 10 \\ 0 & 1 & -3 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \end{array} \right] \end{array}$$

Since the leading 1's are in columns 1, 2, 3 & 4 we conclude that the first two rows of A form a basis for the Null Space of A that is

$$\begin{array}{c} \left[\begin{array}{cc|c} 1 & -2 & 5 \\ 0 & -7 & 8 \end{array} \right] \frac{1}{7}R_2 \\ \hline \left[\begin{array}{cc|c} 1 & -2 & 5 \\ 0 & 1 & -8/7 \end{array} \right] \end{array}$$

$$\begin{array}{c} \left[\begin{array}{ccc|c} 1 & 0 & 19/7 & 0 \\ 0 & 1 & -8/7 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] R_{1,2R_1} \\ \hline \left[\begin{array}{ccc|c} 1 & 0 & 19/7 & 0 \\ 0 & 1 & -8/7 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] R_{3,7R_2} \\ \hline \left[\begin{array}{ccc|c} 1 & 0 & 19/7 & 0 \\ 0 & 1 & -8/7 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \end{array}$$

basis for null space of A

$$A = \left\{ \left[1, 0, 19/7 \right], \left[0, 1, -8/7 \right] \right\}$$

basis for column space of A

$$\tilde{A} = \begin{bmatrix} 1 & 2 & 0 \\ -1 & 3 & 7 \\ 5 & 2 & 8 \end{bmatrix}$$

Augmented matrix is

$$\begin{array}{c} \left[\begin{array}{ccc|c} 1 & 2 & 0 & 1 \\ 2 & 3 & 2 & 0 \\ 0 & 7 & 8 & 0 \end{array} \right] [A : b] \\ \hline \left[\begin{array}{ccc|c} 1 & 2 & 0 & 1 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{array} \right] \end{array}$$

Javed Nasir
Lecturer U.E.T Peshawar.

(From Sp
Toko.)

$$\begin{array}{r} \overline{0\ 0\ 0} \\ - 0\ 0 \\ \hline 0\ 0\ 0 \end{array}$$

As Ramsey is dead in Gloucester so now he is dead

A
T
D
P

$$\frac{\partial \theta}{\partial \phi} = \theta$$

to have good spots of

$$\begin{array}{r} \overline{54} \\ - 23 \\ \hline 31 \end{array}$$

$$\left[\begin{array}{ccccc} 0 & 0 & -1 & 2 & 0 \\ -8 & 7 & 1 & 0 & 0 \\ 8 & -7 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 \\ \end{array} \right] R_2 + 2R_1 \\ R_3 - 5R_1$$

$$\begin{array}{r} \text{?} \\ \hline 00 - \\ -8 \\ \hline 12 \\ -1 \\ \hline 10 \\ -8 \\ \hline 20 \\ -10 \\ \hline 10 \\ -10 \\ \hline 0 \end{array}$$

$$\begin{array}{r} \boxed{00-00} \\ \hline 00-00 \\ \hline R_3 + 8R_2 \end{array}$$

$\{[1.02], [0 \ 1 \ -1]\}$ form Rain

$$\left[\begin{array}{ccccc} 0 & 0 & 0 & 0 & -1 \\ -14 & -22 & & & \\ -18 & -12 & & & \\ -12 & -12 & & & \\ \hline 0 & 0 & 0 & 0 & \end{array} \right] \xrightarrow{\begin{array}{l} -3R_1 + R_2 \\ -2R_1 + R_4 \\ -R_1 + R_5 \end{array}} \left[\begin{array}{ccccc} 0 & 0 & 0 & 0 & -1 \\ -2R_1 + R_2 & & & & \\ -2R_1 + R_4 & & & & \\ -R_1 + R_5 & & & & \end{array} \right]$$

$$\begin{array}{r} \overline{-pWN} \\ -p\bar{q} - w \\ \overline{qr1\infty +} \\ \overline{00000} \end{array}$$

augmented marker

$$\frac{P}{n} = \frac{\omega}{N}$$

T
S
E
R
E
R
E

$$\begin{array}{r} 0 \ 0 \\ 0 - 0 \\ \hline 0 \ 1 \ 0 \end{array}$$

$$\left[\begin{array}{ccccc} 1 & 3 & 7 & 10 \\ 0 & 1 & 6R_5 & 10 \\ 0 & -4 & -22 & 10 \\ 0 & -8 & -12 & 10 \\ 0 & -2 & -2 & 10 \end{array} \right] - R_5 R_2$$

$$\left[\begin{array}{ccccc} 1 & 0 & 14R_5 & 10 \\ 0 & 1 & 6R_5 & 10 \\ 0 & 0 & -8 & 0 \\ 0 & -8 & -12 & 0 \\ 0 & -2 & -2 & 0 \end{array} \right] - 3R_2 + R_1$$

$$\left[\begin{array}{ccccc} 1 & 0 & 14R_5 & 10 \\ 0 & 1 & 6R_5 & 10 \\ 0 & 0 & -8 & 0 \\ 0 & -8 & -12 & 0 \\ 0 & -2 & -2 & 0 \end{array} \right] - 7R_5 + R_3$$

$$\left[\begin{array}{ccccc} 1 & 0 & 14R_5 & 10 \\ 0 & 1 & 6R_5 & 10 \\ 0 & 0 & -8 & 0 \\ 0 & -8 & -12 & 0 \\ 0 & -2 & -2 & 0 \end{array} \right] - 4R_5 + R_4$$

$$\left[\begin{array}{ccccc} 1 & 0 & 14R_5 & 10 \\ 0 & 1 & 6R_5 & 10 \\ 0 & 0 & -8 & 0 \\ 0 & -8 & -12 & 0 \\ 0 & -2 & -2 & 0 \end{array} \right] - 2R_2 + R_5$$

$$\left[\begin{array}{ccccc} 1 & 0 & 14R_5 & 10 \\ 0 & 1 & 6R_5 & 10 \\ 0 & 0 & -8 & 0 \\ 0 & -6 & -22 & 10 \\ 0 & -2 & -2 & 0 \end{array} \right] - R_5 R_2$$

$$\left[\begin{array}{ccccc} 1 & 0 & 0 & 0 & 10 \\ 0 & 0 & 0 & 0 & 10 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 10 \\ 1 & 0 & 0 & 1 & 0 \end{array} \right] - 4R_5 + R_4$$

$$\left[\begin{array}{ccccc} 1 & 0 & 0 & 0 & 10 \\ 0 & 0 & 0 & 0 & 10 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 10 \\ 1 & 0 & 0 & 1 & 0 \end{array} \right] - 2R_5 + R_2$$

$$\left[\begin{array}{ccccc} 1 & 0 & 0 & 0 & 10 \\ 0 & 0 & 0 & 0 & 10 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 10 \\ 1 & 0 & 0 & 1 & 0 \end{array} \right] - 6R_5 + R_4$$

$$\left[\begin{array}{ccccc} 1 & 0 & 0 & 0 & 10 \\ 0 & 0 & 0 & 0 & 10 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 10 \\ 1 & 0 & 0 & 1 & 0 \end{array} \right] - 8R_5 + R_2$$

Since the Non-zero row is 3 so rank = 3.

(b) where augmented matrix is $[A|0]$

$$\left[\begin{array}{ccccc} 1 & 2 & 3 & 2 & 1 & 0 \\ 0 & -5 & -11 & -8 & -2 & 0 \\ 0 & -6 & -22 & -12 & -2 & 0 \end{array} \right] = [A|0]$$

$$\left[\begin{array}{ccccc} 1 & 2 & 3 & 2 & 1 & 0 \\ 0 & -5 & -11 & -8 & -2 & 0 \\ 0 & -6 & -22 & -12 & -2 & 0 \end{array} \right] - 3R_1 + R_2$$

$$\left[\begin{array}{ccccc} 1 & 2 & 3 & 2 & 1 & 0 \\ 0 & 1 & 14R_5 & 8R_5 & 2R_5 & 0 \\ 0 & -6 & -22 & -12 & -2 & 0 \end{array} \right] - R_1$$

$$\left[\begin{array}{ccccc} 1 & 2 & 3 & 2 & 1 & 0 \\ 0 & 1 & 14R_5 & 8R_5 & 2R_5 & 0 \\ 0 & -6 & -22 & -12 & -2 & 0 \end{array} \right] - 2R_2 + R_1$$

Ex 6.6

136

$$A = \begin{bmatrix} -1 & 2 & 3 \\ -1 & 2 & -1 \\ 3 & -1 & 2 \end{bmatrix}$$

137

$$\sim \left[\begin{array}{ccc|c} 1 & 0 & -7/5 & -4/5 & 1/5 & 1 \\ 0 & 1 & 11/5 & 8/5 & 1/5 & 0 \\ 0 & 0 & -4/5 & -12/5 & 2/5 & 0 \end{array} \right] \quad 6R_2 + R_1$$

$$\sim \left[\begin{array}{ccc|c} 1 & 0 & -7/5 & -4/5 & 1/5 & 1 \\ 0 & 1 & 11/5 & 8/5 & 4/5 & 0 \\ 0 & 0 & 1 & 12/5 & -2/5 & 0 \end{array} \right] - 8R_3 R_1$$

$$\sim \left[\begin{array}{ccc|c} 1 & 0 & 0 & -62/75 & 31/225 & 1/5 \\ 0 & 1 & 0 & 76/75 & 68/225 & 0 \\ 0 & 0 & 1 & 12/45 & -2/45 & 0 \end{array} \right] - \frac{11}{5} R_3 + R_2$$

$$\sim \left[\begin{array}{ccc|c} 1 & 0 & 3 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & -2 & 0 \end{array} \right] R_1 - 2R_2$$

So non-zero rows = 3. Rank = 3.

Row rank = column rank = 3.

θ_{12} is similar to θ_{11}

So rank = 3.
For nullity $AX=0$

$$\left[\begin{array}{cc|c} 1 & 0 & 3 \\ 0 & 1 & -1 \\ 0 & 0 & -2 \end{array} \right]$$

Jamal Nasir
Lecture U.E.T Peshawar

$$\begin{aligned} -2x_3 &= 0 \Rightarrow x_3 = 0 \\ x_1 + x_3 &= 0 \Rightarrow x_1 = 0 \\ x_1 + 3x_3 &= 0 \Rightarrow x_1 = 0 \end{aligned}$$

$\theta_{13}, \theta_{14}, \theta_{16}, \theta_{17}$ is similarly to θ_{15}

138 Lineally independent. Inols when $\det \neq 0$

Q25 If $\text{Rank } A = n = 3$

If $\text{Rank } A = n = 3$
then there is unique solution.

$\omega \rho \rho$
 $\nu_0 -$
 $\omega - 0$

$$2(0-2)-1(6-3)+0(4-0)$$

1
2

$-7 \neq 0$ so linearly independent

$$\begin{array}{r} 2 \\ \hline 00 \\ 100 \\ 00 \\ \hline 111 \\ \hline R_3 - 3R_1 \end{array}$$

$$\begin{array}{r} \\ \hline 0 & 0 & -2 & -2 \\ 0 & -1 & -7 & 0 \\ \hline 7 & & & \end{array} \quad R_{4,8}$$

$$\left. \begin{array}{r} 0 \\ 0 \\ - \\ 0 \\ - \\ 0 \\ - \\ 0 \end{array} \right] \begin{array}{l} 2 \\ 2 \\ 2 \\ 2 \\ 2 \\ 2 \\ 2 \\ 2 \end{array} R_3 + 8R_2$$

Non-jealousy 2

for York = 2

No one is older.

θ_{27} is same as θ_{26}

$$\begin{array}{r}
 00 \\
 0 - 2 \\
 00 \\
 \hline
 282
 \end{array}$$

D
11
-O-
O-P
w6w

Non-trivial solution occurs when rank $< n$.

$$\begin{pmatrix} -1 & 2 \\ 0 & -3 \\ 0 & 0 \end{pmatrix} 2P_2 + P_3$$

RanuA < 3 so non-trivial

Ex 6.6

Solution exist when

Rank of A = Rank of $[A:b]$

$$\left[\begin{array}{ccccc} 1 & -2 & -3 & 4 & 1 \\ 2 & -1 & -5 & 6 & 1 \\ 2 & -3 & -1 & -2 & 1 \end{array} \right] \sim \left[\begin{array}{ccccc} 1 & -2 & -3 & 4 & 1 \\ 0 & 1 & 2 & -4 & -1 \\ 0 & 1 & 2 & -4 & -1 \end{array} \right]$$

$$\left[\begin{array}{ccccc} 1 & -2 & -3 & 4 & 1 \\ 0 & 1 & 2 & -4 & -1 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right] \sim \left[\begin{array}{ccccc} 1 & -2 & -3 & 4 & 1 \\ 0 & 1 & 2 & -4 & -1 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right]$$

Rank of A = 3 & Rank of $[A:b] = 2$ Rank of A \neq Rank of $[A:b]$ \Rightarrow So it's has no solution \emptyset 34, \emptyset 36 is similarly \Rightarrow So it has no solution \emptyset 30 & \emptyset 32 is same on \emptyset 31

Solution exist when

Rank A = Rank of $[A:b]$

$$\left[\begin{array}{ccccc} 1 & 2 & 5 & -2 & 1 \\ 2 & 3 & -2 & 4 & 0 \\ 5 & 1 & 0 & 2 & 0 \end{array} \right] \sim \left[\begin{array}{ccccc} 1 & 2 & 5 & -2 & 1 \\ 0 & -1 & -12 & 8 & 0 \\ 0 & -9 & -25 & 12 & 0 \end{array} \right] \sim \left[\begin{array}{ccccc} 1 & 2 & 5 & -2 & 1 \\ 0 & 1 & 12 & -8 & 0 \\ 0 & -9 & -25 & 12 & 0 \end{array} \right] \sim \left[\begin{array}{ccccc} 1 & 2 & 5 & -2 & 1 \\ 0 & 1 & 12 & -8 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right]$$

$$\left[\begin{array}{ccccc} 1 & 2 & 5 & -2 & 1 \\ 0 & 1 & 12 & -8 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right] \sim \left[\begin{array}{ccccc} 1 & 2 & 5 & -2 & 1 \\ 0 & 1 & 12 & -8 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right] \sim \left[\begin{array}{ccccc} 1 & 2 & 5 & -2 & 1 \\ 0 & 1 & 12 & -8 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right] \sim \left[\begin{array}{ccccc} 1 & 2 & 5 & -2 & 1 \\ 0 & 1 & 12 & -8 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right]$$

Rank A = Rank of $[A:b]$

bit matrix

$$\text{Q3} \quad \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 0 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 0 \end{bmatrix} R_{S+R_1}$$

$$\sim \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{bmatrix} R_{S+R_2}$$

$$\text{Non-zero rows} = 2$$

$$\text{rank} = 2.$$

Q4

$$\begin{bmatrix} 1 & 0 & 0 \\ -1 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} R_{4+R_1}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} R_{3+R_2}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} R_{4+R_3}$$

$$\text{rank} = 3 \Rightarrow \text{rank} = 3$$

$$Q_{24} \text{ and } Q_{40} \text{ similarly.}$$

For eigenvalues

$$\det(\lambda I_n - A) = 0$$

$$\det(\lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} - \begin{bmatrix} 3 & -1 \\ -2 & 2 \end{bmatrix}) = 0$$

$$\det(\begin{bmatrix} \lambda & 0 \\ 0 & \lambda \end{bmatrix} - \begin{bmatrix} 3 & -1 \\ -2 & 2 \end{bmatrix}) = 0$$

$$\det \begin{pmatrix} \lambda-3 & 1 \\ 2 & \lambda-2 \end{pmatrix} = 0$$

$$(\lambda-3)(\lambda-2) - 2 = 0$$

$$\lambda^2 - 3\lambda + 6 - 2 = 0$$

$$\lambda^2 - 5\lambda + 4 = 0$$

$$\boxed{\lambda = 1} \quad \boxed{\lambda = 4}$$

For eigenvectors

$$(\lambda I_n - A)x = 0$$

$$\text{For } \lambda = 1$$

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} - \begin{bmatrix} 3 & -1 \\ -2 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = 0$$

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} - \begin{bmatrix} 3 & -1 \\ -2 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}.$$

$$\begin{bmatrix} -2 & 1 \\ 2 & -1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}.$$

$$-2x_1 + x_2 = 0$$

$$2x_1 - x_2 = 0$$

$$x_1 = x_2$$

$$x_2 = \gamma (\text{any real no.})$$

$$x = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

For $\lambda = 4$

$$4 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} - \begin{bmatrix} 3 & -1 \\ -2 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}.$$

$$\begin{bmatrix} 4 & 0 \\ 0 & 4 \end{bmatrix} - \begin{bmatrix} 3 & -1 \\ -2 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}.$$

$$\begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}.$$

$$x_1 + x_2 = 0$$

$$2x_1 + 2x_2 = 0$$

$$x_1 = -x_2$$

$$x_2 = \gamma (\text{any real no.})$$

$$x = \begin{bmatrix} -\gamma \\ \gamma \end{bmatrix}$$

$$\text{Q.E.D. } A = \begin{bmatrix} 2 & 2 & 3 \\ 1 & 2 & -1 \\ 2 & -2 & 1 \end{bmatrix}$$

For eigenvalues
 $\det(\lambda I_n - A) = 0$

$$\det \left(\lambda \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} - \begin{bmatrix} 2 & 2 & 3 \\ 1 & 2 & -1 \\ 2 & -2 & 1 \end{bmatrix} \right) = 0$$

$$\det \left(\begin{bmatrix} \lambda & 0 & 0 \\ 0 & \lambda & 0 \\ 0 & 0 & \lambda \end{bmatrix} - \begin{bmatrix} 2 & 2 & 3 \\ 1 & 2 & -1 \\ 2 & -2 & 1 \end{bmatrix} \right) = 0$$

$$\det \left(\begin{bmatrix} \lambda - 2 & -2 & -3 \\ -1 & \lambda - 2 & -1 \\ -2 & 2 & \lambda - 1 \end{bmatrix} \right) = 0$$

$$\lambda - 2 \left[(\lambda - 1)(\lambda - 2) + 2 \right] + 2 \left[(-\lambda - 1) - 2 \right] - 3 \left[-2 + 2(\lambda - 2) \right] = 0$$

$$\Rightarrow (\lambda - 2)(\lambda^2 - 3\lambda + 4) + (-2\lambda - 2) + (-6\lambda + 18) = 0$$

$$\lambda^3 - 5\lambda^2 + 2\lambda + 8 = 0$$

$$\lambda = -1$$

$$\begin{array}{c|ccc} 8 & 1 & -1 & 1 \\ & & -5 & 2 \\ \hline & 1 & -1 & 6 \\ & & -6 & 8 \\ \hline & & 0 & 0 \end{array}$$

$$(\lambda + 1)(\lambda^2 - 6\lambda + 8) = 0$$

$$\lambda + 1 = 0$$

$$\lambda - 1$$

$$\lambda^2 - 4\lambda + 8 = 0$$

$$\lambda^2 - 4\lambda - 2\lambda + 8 = 0$$

$$(\lambda - 4)(\lambda - 4) = 0$$

$$\lambda - 2 = 0, \lambda - 4 = 0$$

$$\lambda = 2, \lambda = 4.$$

$$\text{Now } \lambda = -1, \lambda = 2, \lambda = 4.$$

For eigenvector

$$(\lambda I_n - A)X = 0$$

$$\lambda = -1$$

$$= \begin{pmatrix} -1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix} - \begin{pmatrix} 2 & 2 & 3 \\ 1 & 2 & 1 \\ 2 & 2 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

$$\lambda = 2, \lambda = 4 \quad S.V.S$$



$$\lambda = -1$$

$$= \begin{pmatrix} -1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix} - \begin{pmatrix} 2 & 2 & 3 \\ 1 & 2 & 1 \\ 2 & 2 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}.$$

Jill Laser Photostate

Matric to M.A.
Notes are available



Jill Laser Photostate

Matric to M.A.
Notes are available



$$\begin{pmatrix} -3 & -2 & -3 \\ -1 & -3 & -1 \\ -2 & -2 & -2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

Qur'Shabab
0300 5829015
Arif: 0345 9228401
Ph: 091-5510987
Qur'Shabab
0300 5829015
Arif: 0345 9228401
Ph: 091-5510987
Matric to M.A.
Notes are available

Shop #58 coffee shop bazar Islamia college Peshawar university
Shop #58 coffee shop bazar Islamia college Peshawar university

$$(A/h) = \begin{bmatrix} -3 & -2 & -3 & 1 \\ -1 & -3 & -1 & 1 \\ -2 & -2 & -2 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

$$\underline{Q3} \quad \text{Def } A = \begin{bmatrix} 1 & 2 & 1 \\ 0 & 1 & 2 \\ -1 & 3 & 2 \end{bmatrix}$$

The characteristic Poly: is

$$P(\lambda) = \text{Def } (\lambda I_3 - A)$$

$$= \text{Def} \left(\begin{bmatrix} \lambda & 0 & 0 \\ 0 & \lambda & 0 \\ 0 & 0 & \lambda \end{bmatrix} - \begin{bmatrix} 1 & 2 & 1 \\ 0 & 1 & 2 \\ -1 & 3 & 2 \end{bmatrix} \right)$$

$$= \text{Def} \left(\begin{bmatrix} \lambda-1 & -2 & -1 \\ 0 & \lambda-1 & -2 \\ 1 & -3 & \lambda-3 \end{bmatrix} \right)$$

$$= (\lambda-1)[(\lambda-1)(\lambda-2)-6] + 2(0+2) - 1(0+1-\lambda)$$

$$= (\lambda-1)(\lambda^2 - 3\lambda + 2) + 4 + \lambda - 1$$

$P(\lambda) = \lambda^3 - 4\lambda^2 + 7$ which is required char poly.

$\theta_4, \theta_5, \theta_6, \theta_7$ is similarly for Q3

$$A = \begin{bmatrix} 0 & 1 & 2 \\ 0 & 0 & 3 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\begin{array}{c} \text{Ex. 8.1} \\ \text{143} \end{array}$$

The characteristic Poly. of matrix A is given by.

$$P(\lambda) = \text{Def } (\lambda I_3 - A)$$

$$= \text{Def} \left(\begin{bmatrix} \lambda & -1 & -2 \\ 0 & \lambda & -3 \\ 0 & 0 & \lambda \end{bmatrix} \right)$$

$$= \lambda(\lambda^2 + 0) + 1(0 + 0) - 2(0 + 0)$$

$$P(A) = \lambda^3$$

$$\text{Now } P(\lambda) = \lambda^3 = 0$$

$\lambda = 0$ eigen value.

For Eigenvalue

$$(\lambda I_3 - A)X = 0$$

$$\left(\lambda \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} - \begin{bmatrix} 0 & -2 & 0 \\ 0 & 0 & 3 \\ 0 & 0 & 0 \end{bmatrix} \right) \begin{bmatrix} X_1 \\ X_2 \\ X_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Jai Laser Photostate

Color Photos

copy services

Aarti Books publisher
Ph. 0391-560097

Omar Shahab
0300 540015

Aarti Books publisher
Ph. 0391-560097

Jai Laser Photostate

Matrix to M.A.

Notes are available

Shreeji coffee shop bazar Islamia college Peshawar university

Shreeji coffee shop bazar Islamia college Peshawar university



$$\begin{pmatrix} 0 & -1 & 0 & -1 \\ 0 & 0 & 0 & 3 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

$$\begin{pmatrix} 0 & -1 & -2 & 0 \\ 0 & 0 & -3 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

$$(A/x) \begin{pmatrix} 0 & -1 & -2 & 0 \\ 0 & 0 & -3 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

~~3x3~~ $| x_3 = 0$

$$-x_2 - 2x_3 = 0$$

$$-x_2 = 2x_3$$

$$-x_2 = 2(0)$$

$$\boxed{-x_2 = 0}$$

$$| x_1 = x$$

$$X = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} x \\ x \\ x \\ x \end{bmatrix}$$

$\theta_9, \theta_{10}, \theta_{11}, \theta_{12}, \theta_{13}, \theta_{14}, \theta_{15}$

Similarly:

