

0.0.1 Transfer Matrix method

Given the Hamiltonian above we can write the corresponding partition function in the following symmetric form:

$$Z_N(k, h) = \sum_{S_1=\pm 1} \sum_{S_2=\pm 1} \cdots \sum_{S_N=\pm 1} \left[e^{kS_1S_2 + \frac{h}{2}(S_1+S_2)} \right] \left[e^{kS_2S_3 + \frac{h}{2}(S_2+S_3)} \right] \cdots \left[e^{kS_NS_1 + \frac{h}{2}(S_N+S_1)} \right] \quad (1)$$

Suppose you have a sort of $\sum_j M_{ij} P_{jk}$, what we have done is doing something like that. In the previous form Z_N can be written as a product of matrices

$$\begin{aligned} Z_N(h, k) &= \sum_{S_1=\pm 1} \cdots \sum_{S_N=\pm 1} \prod_{i=1}^N \exp \left[kS_iS_{i+1} + \frac{h}{2}(S_i + S_{i+1}) \right] \\ &= \sum_{S_1=\pm 1} \cdots \sum_{S_N=\pm 1} \langle S_1 | \mathbb{T} | S_2 \rangle \langle S_2 | \mathbb{T} | S_3 \rangle \cdots \langle S_N | \mathbb{T} | S_1 \rangle \end{aligned} \quad (2)$$

where \mathbb{T} is a 2×2 matrix defined as

$$\langle S | \mathbb{T} | S' \rangle = \exp \left[kSS' + \frac{h}{2}(S + S') \right] \quad (3)$$

Note that the labels of the matrix corresponds to the values of S_i . Hence its dimension depends on the number of possible values a spin S_i can assume. It can also depend on how many spins are involved in the interacting terms that are present in the hamiltonian ($k_{LL} \sum S_iS_{i+1}S_{i+2}S_{i+3}$). For Ising $S_i = \pm 1$ and nearest neighbour interaction implies that we have 2 values and that \mathbb{T} is a 2×2 matrix whose components are

$$\langle +1 | \mathbb{T} | +1 \rangle = \exp[k + h] \quad (4a)$$

$$\langle +1 | \mathbb{T} | -1 \rangle = \langle -1 | \mathbb{T} | +1 \rangle = \exp[k - h] \quad (4b)$$

$$\langle -1 | \mathbb{T} | -1 \rangle = \exp[-k] \quad (4c)$$

The explicit representation is

$$\mathbb{T} = \begin{pmatrix} e^{k+h} & e^{-k} \\ e^{-k} & e^{k-h} \end{pmatrix} \quad (5)$$

Let us now introduce some useful notations and relations using the bra-ket formalism: subequations

$$|S_i^{(+)}\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix}_i \quad |S_i^{(-)}\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix}_i \quad (6)$$

$$\langle S_i^{(+)} | = (1^*, 0)_i \quad \langle S_i^{(-)} | = (0, 1^*)_i \quad (7)$$

The identity relation is:

$$\sum_{S_i=\pm 1} |S_i\rangle \langle S_i| = |S_i^{(+)}\rangle \langle S_i^{(+)}| + |S_i^{(-)}\rangle \langle S_i^{(-)}| = \mathbb{1} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad (8)$$

By using the last property we can write

$$\begin{aligned} Z_N(K, h) &= \sum_{S_1=\pm 1} \cdots \sum_{S_N=\pm 1} \langle S_1 | \mathbb{T} | S_2 \rangle \langle S_2 | \mathbb{T} | S_3 \rangle \cdots \langle S_N | \mathbb{T} | S_1 \rangle \cdots \\ &= \sum_{S_1=\pm 1} \langle S_1 | \mathbb{T}^N | S_1 \rangle = \text{Tr}[\mathbb{T}^N] \end{aligned} \quad (9)$$

this is exactly the trace of the matrix. Being \mathbb{T} symmetric, we can diagonalize it by an unitary transformation

$$\mathbb{T}_D = \mathbb{P}^{-1} \mathbb{T} \mathbb{P} \quad (10)$$

with $\mathbb{P} \mathbb{P}^{-1} = \mathbb{1}$.

$$\begin{aligned} \text{Tr}[\mathbb{T}^N] &= \text{Tr} \left[\underbrace{\mathbb{T} \mathbb{T} \mathbb{T} \dots \mathbb{T}}_N \right] = \text{Tr} [\mathbb{P} \mathbb{P}^{-1} \mathbb{T} \mathbb{P} \mathbb{P}^{-1} \mathbb{T} \mathbb{P} \dots \mathbb{P}^{-1} \mathbb{T} \mathbb{P} \mathbb{P}^{-1}] \\ &= \text{Tr} [\mathbb{P} \mathbb{T}_D^N \mathbb{P}^{-1}] \underset{\substack{\text{cyclic property} \\ \text{of the trace}}}{=} \text{Tr} [\mathbb{T}_D^N \mathbb{P}^{-1} \mathbb{P}] \\ &= \text{Tr} [\mathbb{T}_D^N] \end{aligned} \quad (11)$$

where

$$\mathbb{T}_D = \begin{pmatrix} \lambda_+ & 0 \\ 0 & \lambda_- \end{pmatrix} \quad \mathbb{T}_D^N = \begin{pmatrix} \lambda_+^N & 0 \\ 0 & \lambda_-^N \end{pmatrix} \quad (12)$$

with λ_{\pm} are the eigenvalues with $\lambda_+ > \lambda_-$.

Remark. \mathbb{P} is the similitude matrix whose columns are given by the eigenvectors of λ_{\pm} .

We finally have:

$$Z_N(K, h) = \text{Tr} [\mathbb{T}_D^N] = \lambda_+^N + \lambda_-^N \quad (13)$$

Remark. As mentioned previously the dimension of the transfer matrix \mathbb{T} and hence the number of eigenvalues $\{\lambda\}$ depend both on the possible values of S_i and on the number of sites involved in terms of the Hamiltonian (range of interaction).

Example 1. For example consider the Ising ($S_i = \pm 1$) with nearest neighbour and next nearest neighbour interactions. The hamiltonian is:

$$\mathcal{H} = k_1 \sum_i S_i S_{i+1} + k_2 \sum_i S_i S_{i+1} S_{i+2} S_{i+3} \quad (14)$$

Because of the second term now there are $2^4 = 16$ possible configurations that can be described by using a 4×4 transfer matrix that we can write formally as

0.1 Lesson

We have $S_i = +1, 0, -1$, therefore it can assume three different values. This is a *deluted* ising model.

Let us suppose there are $(n+2)$ possible values:

$$\left\langle S_i^{(3)} \right| = (0, 0, 1^*, 0, \dots) \quad (15)$$

$$\left| S_i^{(3)} \right\rangle = \begin{pmatrix} 0 \\ 0 \\ 1 \\ \vdots \\ 0 \end{pmatrix} \quad (16)$$

$$\sum_{S_i} |S_i\rangle \langle S_i| = \mathbb{1}, \quad \mathbb{1} \in (n+2) \times (n+2) \quad (17)$$

$$S_i = \sum_{S_i} |S_i\rangle S_i \langle S_i| \quad (18)$$

Now $\{\lambda_+, \lambda_-, \lambda_1, \dots, \lambda_n\}$, with $\lambda_+ > \lambda_- \geq \lambda_1 \geq \dots \geq \lambda_n$.

$$Z_N(\{k\}, h) = \lambda_+^N + \lambda_-^N + \sum_{i=1}^n \lambda_i^N \quad (19)$$

$$\mathbb{T} = \mathbb{P}\mathbb{T}_D \mathbb{P}^{-1} = \sum_i |t_i\rangle \lambda_i \langle t_i| \quad (20)$$

Now we are interested in the limit of the bulk free energy:

$$F_N() = -k_B T \log Z_N() \quad (21)$$

In general, looking at the thermodynamic limit:

$$f_b(\{k\}, h) = \lim_{N \rightarrow \infty} \frac{1}{N} F_N = \lim_{N \rightarrow \infty} \frac{1}{N} (-k_B T) \log \left[\lambda_+^N + \lambda_-^N + \sum_{i=1}^n \lambda_i^N \right] \quad (22)$$

$$\rightarrow \lim_{N \rightarrow \infty} \frac{-k_B T}{N} \log \left[\lambda_+^N \left(1 + \frac{\lambda_-^N}{\lambda_+^N} + \sum_i \left(\frac{\lambda_i}{\lambda_+} \right)^N \right) \right] = -k_B T \log \lambda_+ \quad (23)$$

So we have obtained

$$f_b = -k_B T \log \lambda_+ \quad (24)$$

This is simply because λ_+ is the largest.

Theorem 0.1.1 (Perron-Frobenius). *Let A be a $m \times m$ matrix. If A is finite ($m < \infty$) and $A_{ij} > 0, \forall i, j$, ($A_{ij} = A_{ij}(\vec{x})$) therefore λ_+ has the following properties:*

1. $\lambda_+ \in \mathbb{R}^+$
2. $\lambda_+ \neq$ from $\{\lambda_i\}_{i=1, \dots, m-1}$
3. λ_+ is a analytic function of its arguments

Try to change $A_{ij} > 0$ or the hypothesis that A is finite and see what is obtained.

0.2 Correlation function

Now we calculate the two points correlation function. We want the fluctuation respect to the average:

$$\Gamma_R = \langle S_1 S_R \rangle - \langle S_1 \rangle \langle S_R \rangle \quad (25)$$

we expect from physics that

$$\Gamma_R \underset{R \rightarrow \infty}{\sim} \exp[-R/\xi] \quad (26)$$

$$\xi^{-1} = \lim_{R \rightarrow \infty} \left[-\frac{1}{R} \log [\langle S_1 S_R \rangle - \langle S_1 \rangle \langle S_R \rangle] \right] \quad (27)$$

$$\langle S_1 S_R \rangle_N = \frac{1}{Z_N} \sum_{\{S\}} S_1 S_R \exp[-\beta \mathcal{H}] \quad (28)$$

$$= \frac{1}{Z_N} \sum_{\{S\}} S_1 \langle S_1 | \mathbb{T} | S_2 \rangle \dots \langle S_{R-1} | \mathbb{T} | S_R \rangle S_R \langle S_R | \mathbb{T} | S_{R+1} \rangle \dots \langle S_N | \mathbb{T} | S_1 \rangle \quad (29)$$

$$= \frac{1}{Z_N} \sum_{S_1, S_R} S_1 \langle S_1 | \mathbb{T}^{R-1} | S_R \rangle S_R \langle S_R | \mathbb{T}^{N-R+1} | S_1 \rangle \quad (30)$$

$$\mathbb{T}^{R-1} = \sum_{i=1}^{n+2} |t_i\rangle \lambda_i^{R-1} \langle t_i| \quad (31)$$

$$\mathbb{T}^{N-R+1} = \sum_{i=1}^{n+2} |t_i\rangle \lambda_i^{N-R+1} \langle t_i| \quad (32)$$

$$\langle S_1 | \mathbb{T}^{R-1} | S_R \rangle = \sum_{i=1}^{n+2} \langle S_i | t_i \rangle \lambda_i^{R-1} \langle t_i | S_R \rangle \quad (33)$$

$$\sum_{\{S\}} S_1 S_R e^{-\beta \mathcal{H}} = \sum_{S_1 S_R} S_1 \sum_{i=1}^{n+2} \langle S_1 | t_i \rangle \lambda_i^{R-1} \langle t_i | S_R \rangle S_R \sum_{j=1}^{n+2} \langle S_R | t_j \rangle \lambda_j^{N-R+1} \langle t_j | S_1 \rangle \quad (34)$$

Define:

$$\mathbb{S}_1 = \sum_{S_1} |S_1\rangle S_1 \langle S_1| \quad (35)$$

$$\mathbb{S}_R = \sum_{S_R} |S_R\rangle S_R \langle S_R| \quad (36)$$

$$\rightarrow = \sum_{ij} \langle t_j | \mathbb{S}_1 | t_i \rangle \lambda_i^{R-1} \langle t_i | \mathbb{S}_R | t_j \rangle \lambda_j^{N-R+1} \quad (37)$$