Surveillance Coverage, Treatment Precedence, and Workload Constraints in Machine Learning Early Warning for Sepsis: An Operational Evaluation

Alireza Shayan[1], **Somayeh Afrasiabi**[1,*]

[1]Department of Computer Science and Engineering & IT,

School of Electrical and Computer Engineering, Shiraz University, Shiraz, Iran

Ali404@hafez.shirazu.ac.ir, afrasiabi@shirazu.ac.ir

[*]Corresponding Author: Somayeh Afrasiabi (afrasiabi@shirazu.ac.ir)

Abstract

Objectives: To evaluate how surveillance coverage, anchor selection, and alert constraints govern machine learning early warning for sepsis in intensive care unit (ICU) surveillance.

Materials and Methods: We retrospectively evaluated eight predictive architectures across 8,254 adult first ICU stays from MIMIC-III MetaVision (2008–2012; 97,700 hourly windows; prevalence 0.694%) under a prespecified second-day surveillance protocol (hours 19–30). CareVue provided development data, and historical prescriptions provided sensitivity bounds. We formalized an operational cascade ($S \rightarrow V \rightarrow O \rightarrow D$), evaluated an outcome-stratified hybrid risk set ($\mathcal{R}_0^{\text{hybrid}}$ and $\mathcal{R}_2^{\text{hybrid}}$) censoring cases at infection suspicion and controls at earliest prescription, and estimated burden-matched frontiers under 12-hour alert suppression.

Results: Among 408 consensus Sepsis-3 stays, 257 (63.0%) developed onset after hour 36 ($V = 0$). Opportunity depended strongly on the selected anchor: 150 of 151 covered stays (99.3%) preceded the infection-suspicion anchor (T_{susp}), but under point-assigned earliest hospital prescription start dates, 0 covered stays remained eligible (0 definite, 0

point-assigned), precluding discrimination estimation. On the outcome-stratified hybrid set, discrimination remained stable (BiMamba SSM: AUROC 0.6433 on $\mathcal{R}_0^{\text{hybrid}}$ vs. 0.6326 canonical). Under an alert budget of 10 episodes/100 patient-days, PPV was 0.98%–1.72% (NNE: 58.1–101.8), with timely yield 0.98%–1.72%.

Discussion: Evaluation feasibility and apparent performance depended heavily on clinical anchor selection. Stable discrimination on the hybrid set reflects outcome-specific censoring rather than uniform prospective policy; under earliest-prescription restriction, point-assigned discrimination was incomputable due to zero retained positive cases.

Conclusion: Sepsis early-warning evaluations must report protocol coverage, avoid outcome-dependent evaluation rules, and benchmark against explicit alert budgets.

Keywords: Sepsis Early Warning, Surveillance Truncation, Anchor Selection, Alert Budget Frontier, State-Space Models.

1 Introduction

1.1 Background and Significance

Sepsis remains a leading cause of critical illness, inpatient mortality, and healthcare expenditure globally [1,2]. Clinical treatment urgency depends critically on presentation: the Surviving Sepsis Campaign international guidelines recommend immediate antimicrobial therapy, ideally within one hour, for possible septic shock or a high likelihood of sepsis, while recommending rapid diagnostic assessment for possible sepsis without shock [3, 4]. To support prompt intervention, the biomedical informatics community has heavily invested in machine learning (ML) models for early warning. Scores of retrospective investigations have reported high discriminative capability, frequently reporting area under the receiver operating characteristic curve (AUROC) values between 0.80 and 0.88 ahead of consensus onset [5]. However, published ICU prediction studies differ substantially in outcome implementation, temporal alignment, and evaluation design; consequently, reported AUROCs should not be interpreted as directly comparable measures of deployment readiness [5, 6]. Prior MIMIC-III analyses have shown that alternative operational onset definitions can change measured performance as much as or more than changing model

class [22]. These findings motivate separating reference-event construction from surveillance coverage and alert-policy evaluation. Time-sensitive sepsis benchmarks have also incorporated utility functions that reward and penalize alerts according to their timing [7]. Such approaches reinforce the need to report temporal evaluation rules, although benchmark utility is distinct from measured clinical benefit.

Clinical evaluations have reported heterogeneous performance and outcome associations [8,9], while clinician-vignette studies have identified discrepancies between operational onset labels and recommended treatment timing [11]. External evaluation of the original Epic Sepsis Model demonstrated that reported performance depended heavily on the evaluation level and timing rules: hospitalization-level and prediction-level workload estimates were not interchangeable, emphasizing the need to report the exact unit of evaluation [8]. Prospective evaluations of early-warning systems such as TREWS have found associations between timely alert confirmation and improved outcomes among selected patients identified prior to antibiotic initiation [9, 10]. However, because provider response was observational rather than randomized, these findings support potential clinical utility without establishing universal causal efficacy [9].

Furthermore, ICU clinicians already encounter substantial alarm exposure from bedside physiological monitors [14]. In other clinical decision-support settings, repeated alerts and work complexity have been associated with lower alert acceptance, supporting explicit reporting of notification burden [15, 16].

This disconnect between retrospective benchmarks and prospective clinical utility is commonly attributed to discriminative collapse: the conjecture that ML models fail because they cannot anticipate sepsis prior to clinician recognition or empiric treatment [12]. Pretreatment evaluation has previously been proposed to distinguish genuine advance warning from recognition of care processes already underway [12]. However, existing evaluations frequently conflate three distinct operational mechanisms:

1. **Surveillance Protocol Coverage (V):** Protocol surveillance windows (e.g., fixed observation periods) may miss episodes occurring outside the evaluated interval.

2. **Treatment Precedence (O):** Decision windows may occur after clinicians have already recognized infection and initiated antimicrobial therapy.

3. **Clinical Alert-Budget Constraints (D):** Even when models maintain pre-treatment discrimination, bedside alert-rate constraints bound real-world sensitivity and positive predictive value (PPV).

To distinguish these evaluation components, we conducted an operational evaluation across 8 representative predictive architectures using retrospectively extracted electronic health record (EHR) data from the Beth Israel Deaconess Medical Center (BIDMC) MIMIC-III database (v1.4; CareVue development and MetaVision evaluation) [19]. Our contributions are fourfold:

1. **Outcome-Stratified Hybrid Discrimination Analysis:** We evaluate discrimination on outcome-stratified hybrid sets ($\mathcal{R}_0^{\text{hybrid}}$ and $\mathcal{R}_2^{\text{hybrid}}$), demonstrating that while mathematical discrimination changes minimally, differences from canonical discrimination reflect both evaluated observations and outcome-specific selection rather than uniform prospective eligibility.

2. **Operational Alert-Budget Frontier:** We estimate retrospective burden-matched operating characteristics under a 12-hour alert suppression rule across prespecified workload scenarios (5, 10, and 20 emitted alert episodes per 100 patient-days).

3. **Three-Stage Population Decomposition and Evaluability:** We formalize the cascade $S \to V \to O \to D$, showing that under the second-day protocol, 63.0% of sepsis stays develop onset after surveillance, while evaluability under pre-exposure constraints depends entirely on the selected anchor (99.3% under T_{susp} vs. zero positive cases under point-assigned $T_{\text{Rx,start}}$).

4. **Multi-Clock Timestamp Uncertainty Analysis:** We bound prescription-proxy opportunity using calendar-day intervals in historical MIMIC-III prescriptions, conditional on assigned Sepsis-3 timestamps and the observed surveillance schedule, and contrast interval bounds with declared point-assigned prescription proxies.

2 Materials and Methods

2.1 Study Design and Cohort Definition

We conducted a retrospective evaluation with sequential prediction replay using the MIMIC-III critical care repository (v1.4) from BIDMC [19], evaluating transportability across clinical information systems from CareVue (2001–2008 development) to MetaVision (2008–2012 prospective-replay benchmark), with broad historical prescriptions providing a sensitivity analysis. Analyses used versioned extraction queries with dataset identifiers and checksums recorded in the release manifest. The study cohort was defined as adult patients ($\geq$ 18 years) admitted to an intensive care unit with a total ICU length of stay LOS $\geq$ 30h. To avoid within-patient clustering across multiple admissions, only the chronological first ICU stay per patient was retained. Complete cohort derivation schemas, exclusion criteria, train/validation/test partitions, feature dictionaries, missing-data rules, model architectures, and executable Sepsis-3 definitions are detailed in the Supplementary Methods Package.

Surveillance was evaluated under a prespecified second-day ICU surveillance protocol: predictions were generated at 12 scheduled hourly decision bins $[t-1, t]$ for $t \in \{19, 20, \ldots, 30\}$ hours post-admission. The test partition comprised 8,254 unique patient ICU stays (97,700 evaluated windows, 4,070.8 patient-days of surveillance volume). Stays were censored at Sepsis-3 onset (N = 110 stays, 623 windows) or upon observation-availability cessation prior to hour 30 under the prespecified feature-generation schedule (N = 180 control stays, 725 windows; all cohort stays had total LOS $\geq$ 30h). Reporting was organized in accordance with TRIPOD+AI and STROBE recommendations [20, 21].

2.2 Outcome Definition and Operational Clocks

Sepsis was defined according to a study-specific operational adaptation of consensus Sepsis-3 criteria [1, 2]: an acute increase in Sequential Organ Failure Assessment (SOFA) score $\geq$ 2 points over baseline ($\text{SOFA}_{\text{base}}$, evaluated over the 24-hour lookback $[T_{\text{susp}} - 24\text{h}, T_{\text{susp}}]$) within a clinical evaluation window $[T_{\text{susp}} - 24\text{h}, T_{\text{susp}} + 12\text{h}]$ relative to suspected infection (defined as blood culture specimen acquisition paired with systemic antimicrobial administration; detailed in

Supplementary Section S1.5).

To resolve temporal precision and event ambiguity without conflating orders, intended start dates, and physical administrations, we track six operational milestones (with ICU admission as the reference origin $t = 0$, followed by five operational event clocks; Figure 1):

- $T_{\text{admit},i}$: ICU admission timestamp ($t = 0$, reference origin).
- $T_{\text{Rx,start},i}$: Earliest documented systemic antimicrobial prescription start date (`prescriptions.startdate` calendar date coarsened to midnight 00:00:00).
- $T_{\text{abx,pair},i}$: Systemic antimicrobial order date participating in the qualifying infection suspicion pair.
- $T_{\text{culture,pair},i}$: Blood culture specimen collection timestamp (`microbiologyevents.charttime`, minute-level precision).
- $T_{\text{susp},i}$: Operational infection-suspicion anchor, defined per Seymour et al. [2] as $\min(T_{\text{abx,pair}}, T_{\text{culture,pair}})$.
- $T_{\text{S3},i}$: Retrospective consensus Sepsis-3 onset timestamp.

When blood culture precedes antimicrobial prescription ($T_{\text{culture,pair}} < T_{\text{abx,pair}}$), $T_{\text{susp}} = T_{\text{culture,pair}} \neq T_{\text{abx,pair}}$. Let $h = 1$ hour and define the dimensionless lookback coordinate $x_i = (T_{\text{susp},i} - 24h - T_{\text{admit},i})/h$. The search window evaluates discrete hourly bins starting at baseline_start_h$_i = \lfloor x_i \rfloor$ and ending at window_end_h$_i = \lfloor (T_{\text{susp},i} + 12h - T_{\text{admit},i})/h \rfloor$. Because the upper boundary bin is floored, evaluated hours never exceed $T_{\text{susp}} + 12h$, strictly guaranteeing $T_{\text{susp},i} - T_{\text{S3},i} \geq -12.0h$ (empirical minimum: $-11.2h$). For episodes where acute ΔSOFA ≥ 2 is first observed at the earliest search bin ($N = 274/408$, 67.2%), onset is assigned as $T_{\text{S3},i} = T_{\text{admit},i} + h\lfloor x_i \rfloor$, yielding the operational lead gap:

$$G_i = T_{\text{susp},i} - T_{\text{S3},i} = 24h + h \ \text{frac}(x_i) \tag{1}$$

Because $0 \leq \text{frac}(x_i) < 1$ (dimensionless), first-bin-assigned episodes satisfy $24h \leq G_i < 25h$, with an analytical supremum of $25h$ (the reported $24.99h$ is the observed empirical maximum). In 274 of 408 episodes, the assigned onset coincided with the first evaluated search bin. Consequently,




Figure 1: Operational Clocks, Outcome-Stratified Decision Windows, and Population Opportunity Attrition. Panel A illustrates the clinical timeline for an ICU-incident sepsis patient, depicting the search-window lead $G_i = 24h + h\ \mathrm{frac}(x_i) < 25h$ for first-bin-assigned episodes ($N = 274/408$, 67.2%; median +24.2h; empirical max 24.99h), the pre-suspicion decision set $\mathcal{A}_i(\delta) = \{t \in \mathcal{T}_i : 0 < T_{\mathrm{S3},i} - t \leq 6\mathrm{h}, t + \delta \leq T_{\mathrm{susp},i}\}$, and the post-anchor period censored in $\mathcal{R}_\delta^{\mathrm{hybrid}}$. Panel B displays population opportunity attrition comparing Broad Historical MIMIC-III (earliest prescription proxy $T_{\mathrm{Rx,start}}$) and MIMIC-III MetaVision (suspicion anchor T_{susp}), highlighting eligibility under declared midnight assignment. *Alt-text:* Two-panel diagram displaying timeline clocks and mathematical risk-set definitions in Panel A, and horizontal bar charts comparing surveillance coverage and treatment-precedence opportunity in Panel B.

onset timing and derived warning intervals were sensitive to the search-window boundary and bin-assignment convention and should not be interpreted as directly observed biological onset.

2.3 Operational Cascade and Outcome-Stratified Hybrid Risk Sets

Let C denote the evaluated test cohort of unique ICU stays ($N = 8,254$), and let $\mathcal{T}_i \subseteq \{19, 20, \ldots, 30\}$ denote the set of hourly decision bins evaluated for stay i. For patients who never develop consensus Sepsis-3 during hospitalization, we set $T_{\mathrm{S3},i} = \infty$.

The full evaluated early-warning risk set is defined as:

$$\mathcal{R} = \left\{(i,t) : i \in C,\ t \in \mathcal{T}_i,\ t < T_{\mathrm{S3},i}\right\} \tag{2}$$

The binary early-warning prediction target at anticipation horizon $H = 6\mathrm{h}$ is:

$$Y_i(t;H) = \mathbf{1}\left\{t < T_{\mathrm{S3},i} \leq t + H\right\} \tag{3}$$

In the canonical unconstrained risk set ($\mathcal{R}_{\mathrm{canonical}} = \mathcal{R}$), this definition yields exactly 97,700

evaluated hourly windows, comprising 678 horizon-positive observations ($Y = 1$) and 97,022 negative observations ($Y = 0$, prevalence $\pi = 0.694\%$).

To evaluate restricted decision windows, we define the outcome-stratified hybrid risk sets relative to reference clock C_i:

$$\mathcal{R}_{\delta}^{\text{hybrid}} = \left\{(i, t) \in \mathcal{R} : t + \delta \leq C_i\right\} \tag{4}$$

where $C_i = T_{\text{susp},i} = \min(T_{\text{culture,pair},i}, T_{\text{abx,pair},i})$ for incident sepsis stays ($S_i = 1$; determined by $T_{\text{abx,pair}}$ in 399 stays and blood culture charttime in 9 stays), and $C_i = T_{\text{Rx,start},i}$ for non-sepsis control stays ($S_i = 0$) if systemic antimicrobials were ordered during hospitalization ($C_i = \infty$ otherwise). Exactly 2,287 control stays had $T_{\text{Rx,start}} \leq 19$h, censoring all their windows. We evaluated $\mathcal{R}_0^{\text{hybrid}}$ ($\delta = 0$h; 70,192 windows across 5,967 stays; prevalence 0.960‰) and $\mathcal{R}_2^{\text{hybrid}}$ ($\delta = 2$h; 70,129 windows across 5,967 stays; prevalence 0.945‰).

Importantly, sepsis cases and controls were restricted using different clinical-process anchors. Differences from canonical discrimination therefore reflect both changes in the evaluated observations and the outcome-specific selection rule; they are not estimates of performance under a uniform prospective eligibility policy. Under a uniform earliest-prescription policy ($C_i = T_{\text{Rx,start},i}$ across all stays), exactly 0 of 151 covered sepsis stays have point-assigned pre-prescription decision bins ($O_{\text{Rx}}^{\text{point}} = 0, O_{\text{Rx}}^{\text{def}} = 0$). Consequently, the number of horizon-positive observations is zero, rendering discrimination metrics (AUROC and AUPRC) not estimable under point-assigned midnight prescription start dates.

2.4 Alert-Budget Operating Frontiers and Refractory Policy

To quantify simulated emitted-alert burden, we evaluated fixed alert budgets $q \in \{5.0, 10.0, 20.0\}$ emitted alert episodes per 100 surveillance patient-days. Under a 12-hour suppression rule, each stay emits at most one alert episode within the evaluated 12-hour interval ($N_{\text{episodes},i} \leq 1$).

We estimated retrospective burden-matched operating characteristics by selecting score cutoffs from the held-out score distribution. These operating points describe empirical performance at matched emitted-alert rates; they do not represent thresholds prospectively fixed for online

deployment. Patient selection used the maximum score over the evaluated interval, whereas detection timing was assessed at the first eligible threshold crossing under the complete tie-selection rule:

$$t_{\text{first},i} = \min\{t \in \mathcal{T}_i : \text{Elig}_i(t; \tau, \mathcal{B}) = 1\} \tag{5}$$

where $\mathcal{B}$ denotes the deterministic prioritization rule:

$$\text{Elig}_i(t; \tau, \mathcal{B}) = \mathbf{1}\{i \in \mathcal{S}_{\text{alerted}}(K) \ \wedge \ s_i(t) \geq \tau\} \tag{6}$$

with $\mathcal{S}_{\text{alerted}}(K)$ defined as the top K patient replicas under the lexicographical sort:

$$\left(-\max_t s_i(t),\ \text{ICUSTAY_ID},\ \text{copy_idx}\right)$$

2.5 Predictive Models and Statistical Inference

We evaluated 8 models spanning three major architectural paradigms:

1. **State-Space Models:** Bidirectional Mamba (BiMamba SSM) utilizing linear-time selective state-space layers [17, 18].

2. **Deep Sequence Transformers:** Mask-Aware Clinical Transformer (incorporating explicit missingness masks) and Baseline Unmasked Transformer.

3. **Tabular and Ensemble Baselines:** Random Forest, XGBoost, LightGBM, Logistic Regression (L2), and a Multi-Model Stacking Ensemble.

Models consumed a causal 20-hour observation lookback $[t - 20\text{h}, t)$ across 21 physiological time series. All models were trained on fixed pre-surveillance development partitions with hyperparameters and feature normalization frozen prior to this operational evaluation. Complete architecture details, hyperparameter grids, and the stacking protocol are detailed in the Supplementary Methods Package.

Uncertainty was quantified using patient-clustered paired bootstrapping ($B = 1,000$ resamples). For Procedure B (Empirical Burden-Matched Bootstrap), exposure $E_b = \frac{1}{24} \sum_{i \in b} h_i$ and alert target $K_b = \text{round}(q \cdot E_b/100)$ were dynamically recomputed within each resample.

NNE confidence intervals were derived by exact reciprocal transformation of unrounded PPV percentiles: $\text{NNE}_{\text{low}} = 100/\text{PPV}_{\text{upper}}$ and $\text{NNE}_{\text{high}} = 100/\text{PPV}_{\text{low}}$ (with $+\infty$ when $\text{PPV}_{\text{low}} = 0$).

3 Results

3.1 Cohort Progression and Denominator Accounting

Table 1 traces all 8,254 ICU admissions through surveillance eligibility, onset timing, lookback horizons, and pre-treatment actionability. Among 408 consensus Sepsis-3 stays, 257 (63.0%) had clinical onset after hour 36 (median onset: 49.0h post-admission). Under the specified observation schedule and 6-hour horizon, these 257 stays had no evaluated horizon-positive decision time ($V = 0$). The remaining 151 stays (37.0%) had active protocol surveillance coverage ($V = 1$). Of the 151 covered stays, 150 had at least one horizon-positive decision time at or before recorded treatment initiation (99.3% within-covered opportunity; 36.8% of total incident sepsis), providing pre-proxy opportunity under midnight assignment ($\mathcal{R}_0$).

3.2 Outcome-Stratified Hybrid Discrimination Analysis

Table 2 presents discrimination across Canonical Sepsis-3, Hybrid Set ($\mathcal{R}_0^{\text{hybrid}}$), and Hybrid Set with 2h Buffer ($\mathcal{R}_2^{\text{hybrid}}$). Model discrimination remained stable or slightly higher in hybrid risk sets (BiMamba SSM: AUROC 0.6326 canonical vs. 0.6433 on $\mathcal{R}_0^{\text{hybrid}}$, paired ΔAUROC = +0.0107, 95% CI: +0.0060 to +0.0175; Stacking Ensemble: 0.6284 vs. 0.6351, paired ΔAUROC = +0.0067). The increase in AUPRC (BiMamba: 1.209% to 1.656%) reflects higher baseline positive prevalence in the restricted set (0.694% vs. 0.960%). These results are directionally consistent with the antibiotic-order sensitivity findings of Dutta et al. (2026) [13]. However, because cases and controls were restricted using different anchors (T_{susp} vs. $T_{\text{Rx,start}}$), this comparison reflects outcome-specific selection rather than a uniform clinical eligibility policy.

3.3 Observed Burden–Detection Trade-Off Under This Protocol

Table 3 and Figure 2 demonstrate the observed trade-off under the prespecified alert-budget scenarios: bedside detection yield is constrained by alert-rate limits. Alerts were selected across the full canonical surveillance stream (4,070.8 patient-days; 97,700 windows across all 8,254 test stays); C_i was used only retrospectively to adjudicate advance-warning timing ($t_{\text{first},i} + \delta \leq T_{\text{susp},i}$ for sepsis cases), rather than excluding observations prior to alert selection.

At an emitted workload of 10 alert episodes per 100 patient-days (407 emitted episodes across 4,070.8 patient-days):

- Across all 8 models, episode-level PPV ranged from 0.98% to 1.72%, corresponding to 58.1 to 101.8 simulated evaluations per detected patient (NNE).
- BiMamba SSM and LightGBM each uniquely detected **7 patients** (PPV = 1.72%, 95% CI: 0.49%–2.95%; NNE = 58.1, 95% CI: 33.9–203.5).
- Stacking Ensemble, XGBoost, and Random Forest each detected **6 patients** (PPV = 1.47%, NNE = 67.8).
- All qualifying emitted alerts occurred at least two hours before the operational infection-suspicion anchor ($t_{\text{first},i} + 2\text{h} \leq T_{\text{susp},i}$).
- Model rankings varied across budgets: at 20 episodes/100 pt-days, XGBoost detected 15 patients (NNE = 54.3) and Mask-Aware Transformer detected 14 patients (NNE = 58.1), highlighting workload-dependent operating behavior.

3.4 Two-Axis Subgroup and Timestamp Uncertainty Matrix

Table 4 presents the cross-tabulation of Sepsis Onset Timing against Treatment Proxy Timing. Because MIMIC-III prescription start dates (`prescriptions.startdate`) lack time of day, order initiation is evaluated both under declared point assignment (midnight a_i) and under calendar-day intervals $[a_i, b_i) = [\text{midnight}, \text{next midnight})$.

Opportunity depended strongly on the selected clinical-process anchor. Under the primary benchmark evaluating decision opportunity relative to the infection-suspicion anchor (T_{susp}), 150




Figure 2: Operational Alert-Budget Performance Frontier. Panel A displays episode-level positive predictive value (PPV %) across prespecified alert budgets. Panel B displays the Number Needed to Evaluate (NNE) per detected patient under a 12-hour alert suppression rule. Lines illustrate trends across evaluated budget tiers; complete bootstrap confidence intervals and boundary-degenerate notes are reported in Table 3.

of 151 covered patients (99.3% within-covered; 36.8% of total incident sepsis) had decision bins preceding T_{susp} under declared midnight point assignment a_i. Because $a_i \leq T_{\text{susp},i}$ across $[a_i, b_i)$, these 150 stays are definitely eligible throughout the calendar-day interval ($O_i^{\text{def}} = 150$), with only 1 indeterminate stay ($O_i^{\text{indet}} = 1$), bounding opportunity between 36.8% and 37.0% ($[150, 151]/408$; $[99.3\%, 100.0\%]$ within-covered). All indicators strictly satisfy $O_i^{\text{def}} \leq O_i^{\text{point}} \leq O_i^{\text{poss}}$ and $O_i^{\text{poss}} = O_i^{\text{def}} + O_i^{\text{indet}}$.

In contrast, opportunity was substantially lower and uncertain relative to the earliest hospital prescription-start proxy ($T_{\text{Rx,start}}$): exactly 0 covered stays had definite opportunity under midnight point-assignment ($O_i^{\text{def}} = 0, O_i^{\text{point}} = 0$), with 47 indeterminate stays ($O_i^{\text{indet}} = 47$), establishing bounds $[0.0\%, 11.5\%]$ ($[0, 47]/408$; $[0.0\%, 31.1\%]$ within-covered). Because no horizon-positive cases remained before point-assigned earliest prescriptions, this task was not evaluable for discrimination under midnight point assignment. In broad historical MIMIC-III ($N = 3,581$), intervals generate 1 definitely eligible stay, 110 indeterminate stays, and bounds $[0.03\%, 3.10\%]$. The 150/151 result under T_{susp} does not establish that patients were unexposed to antimicrobials or that models predicted prior to first administration. Equally, zero definite opportunity under $T_{\text{Rx,start}}$ reflects date coarsening, not verified bedside exposure.




Figure 3: Cumulative Population Coverage and Actionability Decomposition. Stepwise attrition from 408 incident sepsis episodes through protocol coverage, pre-treatment opportunity, and alert-budget detection.

3.5 Cumulative Population Coverage Decomposition

Figure 3 illustrates the cumulative population coverage decomposition:

$$\frac{151}{408} \times \frac{150}{151} \times \frac{7}{150} = \frac{7}{408} = 1.72\%$$

1. **Surveillance Truncation** ($151/408 = 37.0\%$)**:** 63.0% of sepsis patients develop onset after hour 36, outside the prespecified second-day surveillance protocol.

2. **Pretreatment Actionability** ($150/151 = 99.3\%$)**:** Of the 151 covered stays, 150 had at least one horizon-positive decision time at or before recorded treatment initiation.

3. **Workload Detection Yield** ($7/150 = 4.67\%$)**:** Under an alert budget of 10 episodes/100 patient-days, BiMamba SSM detects 7 patients (1.72% whole-population yield; NNE = 58.1).

4 Discussion

This investigation characterizes the operational trade-offs governing machine learning early warning for sepsis in ICU surveillance:

1. **Outcome-Dependent Anchor Selection Governs Evaluability:** In directional consistency with Dutta et al. [13], model discrimination remained stable on the outcome-stratified hybrid set ($\mathcal{R}_0^{\text{hybrid}}$ and $\mathcal{R}_2^{\text{hybrid}}$). However, this comparison reflects an outcome-dependent selection rule (suspicion anchor for cases, earliest prescription for controls) rather than a uniform prospective policy. Under a uniform point-assigned earliest-prescription rule, zero horizon-positive cases remained, rendering discrimination not estimable.

2. **Detection Yield Is Constrained by Workload and Prevalence:** AUROC values near 0.64 are modest. At 10 emitted episodes per 100 evaluation-bin patient-days, PPV ranged from 0.98% to 1.72%, corresponding to 58.1–101.8 simulated evaluations per detected patient (NNE); observed sensitivity was below 2%.

3. **Surveillance Protocol Coverage Explains Unreached Cases:** Under the evaluated second-day protocol, 63.0% of incident cases develop onset after the monitoring window terminates, preventing multi-hour warning.

4.1 Distinguishing Care-Process Forecasting from Sepsis Anticipation

A critical conceptual error in prior literature is **endpoint substitution**. When benchmarks predict "antibiotic initiation within 6 hours across all patients", they evaluate care-process prediction (predicting clinician ordering behavior, including prophylactic prescribing in non-septic patients). This task is fundamentally distinct from pre-treatment sepsis detection. Pretreatment sepsis evaluation should retain its specified sepsis outcome while restricting eligible decision times; replacing that outcome with antibiotic receipt defines a different prediction task.

Clinician-vignette studies address when clinicians judge treatment appropriate [11]; our analysis addresses prediction performance relative to operational EHR proxies. These are complementary questions, and proxy precedence does not establish clinical appropriateness. Furthermore, while changes under prescription-proxy restriction were small in this cohort, differences in outcome construction, population selection, timestamp precision, and clinical-action definition preclude direct replication of Dutta et al.'s order-based sensitivity analysis [13].

Repeated and low-information notifications have been associated with reduced clinician

acceptance in other decision-support settings [15]. While we quantify simulated emitted-alert exposure and evaluation yield, our analysis did not measure cognitive overload, alert acceptance, or alarm fatigue directly [15, 16]. Clinical impact also depends on whether alerts are reviewed and acted upon [10]. Our evaluation estimates opportunities and simulated review burden; it does not estimate provider adoption or changes in treatment timing.

4.2 Limitations

The analysis estimates operating characteristics in a selected long-stay ICU cohort rather than clinical effectiveness. Burden-matched cutoffs were derived retrospectively from complete evaluated score trajectories, and identifier-based selection was required for tied boundary scores. Accordingly, the frontier does not represent a prospectively locked policy or guarantee a specified alert rate in online deployment.

Temporal precision also constrains interpretation: MIMIC-III prescription dates cannot be treated as directly observed administration times [19, 22]. Our order-proxy analyses therefore require explicit uncertainty bounds and cannot establish hour-resolved clinical treatment delay. Observed treatment precedence does not establish that infection was unrecognized or that earlier antimicrobial administration would have been appropriate. The study did not measure clinician response, review duration, treatment changes, alarm fatigue, or patient outcomes; NNE therefore represents a simulated evaluation count rather than measured clinical workload. Alarm-reduction interventions may suppress both nonactionable and clinically important signals; our short evaluation interval cannot establish the safety or optimal duration of a refractory policy during sustained surveillance [16]. Finally, our cohorts originate from a single medical center (BIDMC) across EHR system transitions (CareVue to MetaVision); multi-center validation across diverse health systems remains essential.

5 Conclusion

In this restricted, long-stay ICU cohort, observed detection yield depended on surveillance coverage, baseline discrimination, and emitted-alert policies. Evaluation feasibility was highly

anchor-dependent: horizon-positive observations were retained before the selected suspicion anchor, but none remained before the earliest prescription-start proxy under midnight assignment, precluding discrimination estimation for that task. Machine learning early-warning evaluations must transparently identify outcome-dependent evaluation rules, report anchor-dependent evaluability, and benchmark under explicit workload constraints.

Ethics Approval and Consent to Participate

This study was conducted using de-identified data from the MIMIC-III (v1.4) database, collected at Beth Israel Deaconess Medical Center (BIDMC). Data collection was approved by the Institutional Review Boards of BIDMC and the Massachusetts Institute of Technology (MIT). The data were de-identified in accordance with Health Insurance Portability and Accountability Act (HIPAA) Safe Harbor standards; informed consent was waived by the reviewing IRBs. Access was granted under PhysioNet Credentialed Data Use Agreement.

Competing Interests

The authors declare that they have no competing financial or non-financial interests.

Funding

This research received no external grant or commercial funding.

Authors' Contributions

AS conceived the study, developed the analytical framework, implemented the pipelines, performed the experiments, and drafted the manuscript. SA provided academic supervision, contributed to methodological design, critically revised the manuscript, and approved the final draft.

AI Assistance Disclosure

During the preparation of this manuscript, the authors utilized large language model and AI assistance (including Perplexity AI and Anthropic/Google AI interfaces) for adversarial peer-review pre-submission auditing, statistical consistency cross-checking, text refinement, and LaTeX typesetting. The authors independently designed and executed all empirical modeling pipelines, statistical data extraction algorithms, and bootstrap evaluations. The authors reviewed, validated, and edited all text and artifacts, and take full responsibility for the integrity, methodology, and conclusions of this work.

Code and Data Availability

Prediction model architectures and frozen weight tensors are archived under analytical data freeze commit `b0932ca` (release manifest `1.0.0-freeze`); evaluation pipeline and bootstrap execution code are archived under release commit `1292103`; manuscript reporting and multi-clock packaging are archived under commit `4038fa2`. All preprocessing pipelines and configurations are available under MIT License at `https://github.com/alireza/Sepsis-Onset-Timing-Paradox`. De-identified patient data are available from PhysioNet under the MIMIC-III Data Use Agreement.

References

[1] Singer M, Deutschman CS, Seymour CW, et al. The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA*. 2016;315(8):801–810. DOI: 10.1001/jama.2016.0287.

[2] Seymour CW, Liu VX, Iwashyna TJ, et al. Assessment of clinical criteria for sepsis: for the Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA*. 2016;315(8):762–774. DOI: 10.1001/jama.2016.0288.

[3] Evans L, Rhodes A, Alhazzani W, et al. Surviving sepsis campaign: international guidelines for management of sepsis and septic shock 2021. *Intensive Care Med*. 2021;47(11):1181–1247. DOI: 10.1007/s00134-021-06506-y.

[4] Kumar A, Roberts D, Wood KE, et al. Duration of hypotension before initiation of effective antimicrobial therapy is the critical determinant of survival in human septic shock. *Crit Care Med*. 2006;34(6):1589–1596. DOI: 10.1097/01.CCM.0000217961.75225.E9.

[5] Moor M, Rieck B, Horn M, Jutzeler CR, Borgwardt K. Early prediction of sepsis in the ICU using machine learning: a systematic review. *Front Med*. 2021;8:607952. DOI: 10.3389/fmed.2021.607952.

[6] Wang Z, Wang W, Sun C, et al. A methodological systematic review of validation and performance of sepsis real-time prediction models. *npj Digit Med*. 2025;8:190. DOI: 10.1038/s41746-025-01587-1.

[7] Reyna MA, Josef CS, Jeter R, et al. Early prediction of sepsis from clinical data: the PhysioNet/Computing in Cardiology Challenge 2019. *Crit Care Med*. 2020;48(2):210–217. DOI: 10.1097/CCM.0000000000004145.

[8] Wong A, Otles E, Donnelly JP, et al. External validation of a widely implemented proprietary sepsis prediction model in hospitalized patients. *JAMA Intern Med*. 2021;181(8):1065–1070. DOI: 10.1001/jamainternmed.2021.2626.

[9] Adams R, Henry KE, Sridharan A, et al. Prospective, multi-site study of patient outcomes after implementation of the TREWS machine learning-based early warning system for sepsis. *Nat Med*. 2022;28(7):1455–1460. DOI: 10.1038/s41591-022-01894-0.

[10] Henry KE, Adams R, Parent C, et al. Factors driving provider adoption of the TREWS machine learning-based early warning system and its effects on sepsis treatment timing. *Nat Med*. 2022;28(7):1447–1454. DOI: 10.1038/s41591-022-01895-z.

[11] Weissman GE, Hubbard RA, Himes BE, et al. Sepsis prediction models are trained on labels that diverge from clinician-recommended treatment times. *AMIA Annu Symp Proc*. 2024;2024:1215–1224.

[12] Kamran F, Tjandra D, Heiler A, et al. Evaluation of sepsis prediction models before onset of treatment. *NEJM AI*. 2024;1(3):AIoa2300032. DOI: 10.1056/AIoa2300032.

[13] Dutta S, McMurry R, Tasi MC, et al. Performance of a sepsis prediction model across different sepsis definitions. *JAMA Netw Open*. 2026;9(4):e265599. DOI: 10.1001/jamanetworkopen.2026.5599.

[14] Drew BJ, Harris P, Zègre-Hemsey JK, et al. Insights into the problem of alarm fatigue with physiologic monitor devices: a comprehensive observational study of consecutive intensive care unit patients. *PLoS One*. 2014;9(10):e110274. DOI: 10.1371/journal.pone.0110274.

[15] Ancker JS, Edwards A, Nosal S, et al. Effects of workload, work complexity, and repeated alerts on alert fatigue in a clinical decision support system. *BMC Med Inform Decis Mak*. 2017;17(1):36. DOI: 10.1186/s12911-017-0430-8.

[16] Paine CW, Goel VV, Ely E, et al. Systematic review of physiologic monitor alarm characteristics and pragmatic interventions to reduce alarm frequency. *J Hosp Med*. 2016;11(2):136–144. DOI: 10.1002/jhm.2520.

[17] Gu A, Dao T. Mamba: linear-time sequence modeling with selective state spaces. *arXiv preprint*. 2023;arXiv:2312.00752. DOI: 10.48550/arXiv.2312.00752.

[18] Gu A, Goel K, Ré C. Efficiently modeling long sequences with structured state spaces. In: *International Conference on Learning Representations (ICLR)*; 2022.

[19] Johnson A, Pollard T, Mark R. MIMIC-III Clinical Database, version 1.4. *PhysioNet*; 2016. DOI: 10.13026/C2XW26.

[20] Collins GS, Moons KGM, Dhiman P, et al. TRIPOD+AI statement: updated guidance for reporting clinical prediction models that use regression or machine learning methods. *BMJ*. 2024;385:e078378. DOI: 10.1136/bmj-2023-078378.

[21] von Elm E, Altman DG, Egger M, et al. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Lancet*. 2007;370(9596):1453–1457. DOI: 10.1016/S0140-6736(07)61602-X.

[22] Cohen SN, Foster J, Foster P, et al. Subtle variation in sepsis-III definitions markedly influences predictive performance within and across methods. *Sci Rep*. 2024;14:1920. DOI: 10.1038/s41598-024-51989-6.

Table 1: Cohort Progression and Denominator Accounting Ledger. Denominators traced from the base test cohort through surveillance eligibility, sepsis onset timing, prediction horizons, and pre-treatment actionability.

Progression Stage / Cohort Partition	Stays (N)	Evaluated Windows	Surveillance Volume	Prevalence	Clinical Definition & Measurement Convention
1. Base Prospective-Replay Test Cohort	8,254	97,700 / 99,048 sched.	4,070.8 pt-days	100.0%	Adult first ICU stays $\geq$ 30h; 1,348 censored (623 onset, 725 cessation)
2. Non-Sepsis Control ICU Stays	7,846	93,427 neg. windows	3,892.8 pt-days	95.06%	Stays without consensus Sepsis-3 during hospitalization
3. Consensus Sepsis-3 Stays ($S = 1$)	408	4,273 (678 pos, 3,595 neg)	178.0 pt-days	4.94%	Blood culture + antibiotic + acute ΔSOFA $\geq$ 2; median onset 49.0h
3a. Late-Onset Sepsis (Onset > 36h)	257	3,084 neg. windows	128.5 pt-days	63.0% of septic	Onset > 36h; lookback $[T_{S3} - 6, T_{S3}]$ does not intersect [19, 30] ($V = 0$)
3b. Surveillance-Covered ($V = 1$)	151	1,189 (678 pos, 511 neg)	49.5 pt-days	37.0% of septic	Onset in [20h, 36h]; lookback intersects surveillance window ($V = 1$)
4. Canonical Sepsis-3 Risk Set ($H = 6$h)	8,254	97,700 (678 pos / 97,022 neg)	4,070.8 pt-days	0.694%	Full unconstrained risk set ($t \in \mathcal{T}_i,\ t < T_{S3,i}$)
5. Outcome-Stratified Hybrid Set $\mathcal{R}_0^{\text{hybrid}}$ ($\delta = 0$h)	5,967	70,192 (674 pos / 69,518 neg)	2,924.7 pt-days	0.960%	Censoring post-anchor ($t \leq C_i$; T_{susp} cases, $T_{\text{Rx,start}}$ controls); 150/151 covered eligible
6. Outcome-Stratified Hybrid Set $\mathcal{R}_2^{\text{hybrid}}$ ($\delta = 2$h)	5,967	70,129 (663 pos / 69,466 neg)	2,922.0 pt-days	0.945%	2-hour clinical buffer ($t + 2\text{h} \leq C_i$); 149/151 covered eligible

Table 2: Benchmark Discrimination Across Outcome-Stratified Hybrid Risk Sets (Held-Out Point Estimates and Clustered 95% CIs). Point estimates calculated directly on held-out test data ($B = 0$); 95% CIs derived from $B = 1,000$ patient-clustered bootstrap resamples. Evaluated on hybrid risk sets $\mathcal{R}_0^{\text{hybrid}}$ and $\mathcal{R}_2^{\text{hybrid}}$ censoring cases at infection-suspicion anchor T_{susp} and controls at earliest prescription $T_{\text{Rx,start}}$. Sepsis cases and controls were restricted using different clinical-process anchors; differences from canonical discrimination reflect both evaluated observations and outcome-specific selection, and do not represent a uniform prospective eligibility policy.

Model Architecture / Metric	Canonical Sepsis-3 ($H = 6$h)	Hybrid $\mathcal{R}_0^{\text{hybrid}}$ ($\delta = 0$h)	Hybrid $\mathcal{R}_2^{\text{hybrid}}$ ($\delta = 2$h)	Paired ΔAUROC ($\mathcal{R}_0^{\text{hybrid}}$ − Canon)
BiMamba SSM (Proposed)	0.6326 (0.5872–0.6733)	**0.6433** (0.5998–0.6835)	**0.6447** (0.6002–0.6860)	**+0.0107** (+0.0060 to +0.0175)
AUPRC	*1.209% (0.892–1.769)*	***1.656%** (1.235–2.329)*	***1.647%** (1.220–2.331)*	—
Multi-Model Stacking Ensemble	0.6284 (0.5829–0.6699)	0.6351 (0.5897–0.6771)	0.6370 (0.5913–0.6799)	+0.0067 (+0.0021 to +0.0137)
AUPRC	*1.206% (0.882–1.748)*	*1.621% (1.207–2.263)*	*1.613% (1.198–2.255)*	—
Random Forest	0.6108 (0.5643–0.6535)	0.6143 (0.5684–0.6575)	0.6173 (0.5712–0.6612)	+0.0034 (-0.0008 to +0.0101)
AUPRC	*1.080% (0.821–1.434)*	*1.470% (1.138–1.920)*	*1.461% (1.120–1.916)*	—
Mask-Aware Clinical Transformer	0.5942 (0.5467–0.6410)	0.5961 (0.5490–0.6422)	0.5954 (0.5480–0.6426)	+0.0020 (-0.0024 to +0.0083)
AUPRC	*1.013% (0.766–1.303)*	*1.382% (1.048–1.760)*	*1.365% (1.032–1.752)*	—
Tabular GBDT (XGBoost)	0.5670 (0.5208–0.6102)	0.5643 (0.5186–0.6082)	0.5652 (0.5192–0.6099)	-0.0027 (-0.0070 to +0.0038)
AUPRC	*0.977% (0.739–1.357)*	*1.320% (1.008–1.817)*	*1.313% (0.998–1.811)*	—
Tabular GBDT (LightGBM)	0.5613 (0.5161–0.6026)	0.5587 (0.5139–0.6010)	0.5598 (0.5145–0.6033)	-0.0026 (-0.0070 to +0.0038)
AUPRC	*0.893% (0.694–1.195)*	*1.222% (0.957–1.629)*	*1.214% (0.949–1.630)*	—
Logistic Regression (L2)	0.5593 (0.5097–0.6048)	0.5654 (0.5157–0.6107)	0.5657 (0.5150–0.6127)	+0.0061 (+0.0023 to +0.0116)
AUPRC	*0.939% (0.716–1.218)*	*1.337% (1.012–1.740)*	*1.325% (1.002–1.731)*	—
Clinical Transformer (Baseline)	0.5239 (0.4808–0.5653)	0.5296 (0.4851–0.5712)	0.5286 (0.4834–0.5705)	+0.0057 (+0.0019 to +0.0110)
AUPRC	*0.808% (0.609–1.381)*	*1.147% (0.857–1.904)*	*1.131% (0.845–1.894)*	—

Table 3: Operational Burden-Matched Alert Frontiers Under 12h Refractory Suppression. Evaluated under Procedure B with exposure recomputation and first-emitted-alert timing verification.

Budget	Model Architecture	Cutoff (τ)	Alerts (K)	Episode PPV % [95% CI]	Det (N)	Sens. Cov %	Yield %	NNE [95% CI]
5.0 ep / 100 pt-d	Stacking Ensemble	0.0145	204	2.94 (0.98–5.88)%	6	3.97 (1.31–7.65)%	1.47 (0.49–2.86)%	34.0 (17.0–102.0)
	LightGBM	0.0117	204	2.94 (0.98–5.39)%	6	3.97 (1.26–7.10)%	1.47 (0.47–2.61)%	34.0 (18.5–102.0)
	BiMamba SSM	0.0187	204	1.96 (0.49–4.45)%	4	2.65 (0.65–6.20)%	0.98 (0.24–2.31)%	51.0 (22.5–204.0)
	XGBoost	0.0122	204	1.96 (0.49–3.94)%	4	2.65 (0.63–5.63)%	0.98 (0.23–2.09)%	51.0 (25.4–204.0)
	Random Forest	0.0177	204	1.47 (0.00–3.45)%	3	1.99 (0.00–4.49)%	0.74 (0.00–1.70)%	68.0 (29.0–∞)
	Clinical Transformer	0.0101	204	0.98 (0.00–2.46)%	2	1.32 (0.00–3.50)%	0.49 (0.00–1.27)%	102.0 (40.6–∞)
	Logistic Regression	0.0103	204	0.98 (0.00–2.94)%	2	1.32 (0.00–3.85)%	0.49 (0.00–1.45)%	102.0 (34.0–∞)
	Mask-Aware Transformer	0.0120	204	0.00 (0.00–0.00)%*	0	0.00 (0.00–0.00)%*	0.00 (0.00–0.00)%*	Undefined (∞)
10.0 ep / 100 pt-d	BiMamba SSM	0.0165	407	1.72 (0.49–2.95)%	7	4.64 (1.33–8.16)%	1.72 (0.49–2.93)%	58.1 (33.9–203.5)
	LightGBM	0.0110	407	1.72 (0.49–2.95)%	7	4.64 (1.38–7.74)%	1.72 (0.49–2.89)%	58.1 (33.9–203.5)
	Stacking Ensemble	0.0131	407	1.47 (0.25–2.46)%	6	3.97 (0.70–6.98)%	1.47 (0.26–2.54)%	67.8 (40.7–407.0)
	XGBoost	0.0114	407	1.47 (0.49–2.94)%	6	3.97 (1.32–7.65)%	1.47 (0.49–2.85)%	67.8 (34.0–203.5)
	Random Forest	0.0158	407	1.47 (0.49–2.70)%	6	3.97 (1.26–7.29)%	1.47 (0.47–2.70)%	67.8 (37.0–204.0)
	Mask-Aware Transformer	0.0110	407	1.23 (0.25–2.46)%	5	3.31 (0.74–6.67)%	1.23 (0.26–2.48)%	81.4 (40.7–407.0)
	Clinical Transformer	0.0094	407	1.23 (0.25–2.46)%	5	3.31 (0.68–6.38)%	1.23 (0.25–2.37)%	81.4 (40.7–407.0)
	Logistic Regression	0.0095	407	0.98 (0.25–1.97)%	4	2.65 (0.59–5.51)%	0.98 (0.23–2.07)%	101.8 (50.8–408.0)
20.0 ep / 100 pt-d	XGBoost	0.0105	814	1.84 (0.98–2.70)%	15	9.93 (5.26–14.47)%	3.68 (1.87–5.29)%	54.3 (37.0–101.9)
	Mask-Aware Transformer	0.0099	814	1.72 (0.74–2.58)%	14	9.27 (4.54–13.77)%	3.43 (1.59–5.19)%	58.1 (38.8–135.7)
	Logistic Regression	0.0088	814	1.35 (0.49–2.09)%	11	7.28 (2.74–11.11)%	2.70 (0.99–4.18)%	74.0 (47.8–203.7)
	BiMamba SSM	0.0142	814	1.23 (0.49–2.09)%	10	6.62 (3.07–10.89)%	2.45 (1.05–4.10)%	81.4 (47.9–203.2)
	LightGBM	0.0105	814	1.23 (0.61–2.09)%	10	6.62 (3.11–11.04)%	2.45 (1.15–4.10)%	81.4 (47.9–163.0)
	Random Forest	0.0131	814	1.11 (0.49–1.85)%	9	5.96 (2.70–9.68)%	2.21 (0.96–3.69)%	90.4 (54.2–203.7)
	Stacking Ensemble	0.0115	814	0.98 (0.37–1.72)%	8	5.30 (2.04–9.15)%	1.96 (0.74–3.43)%	101.8 (58.1–271.3)
	Clinical Transformer	0.0088	814	0.86 (0.25–1.48)%	7	4.64 (1.51–8.18)%	1.72 (0.52–3.04)%	116.3 (67.8–407.0)

*At zero observed detections, the empirical percentile-bootstrap interval was degenerate at zero. This does not establish zero population detection probability. NNE intervals derived by exact reciprocal transformation. 100% of detected sepsis patients satisfied two-hour pre-suspicion timing buffers ($t_{\text{first},i} + 2\text{h} \leq T_{\text{susp},i}$).

Table 4: Two-Axis Subgroup and Timestamp Uncertainty Matrix. Opportunity bounds conditional on fixed consensus Sepsis-3 timestamps and surveillance coverage.

Cohort & Timestamp Source	Sepsis Onset Category	Total (N)	Covered ($V = 1$)	Def. Elig (O^{def})	Point (O^{point})	Indet (O^{indet})	Poss (O^{poss})	Subgroup Bounds	Contribution to Cohort
MetaVision Prospective-Replay	Early Incident (≤ 24h)	47	47 (100.0%)	47	47	0	47	[100.0%, 100.0%]	[47, 47] / 408 (11.5%)
(Primary Benchmark: Suspicion Anchor T_{susp})	Intermediate (24–48h)	156	104 (66.7%)	103	103	1	104	[66.0%, 66.7%]	[103, 104] / 408 (25.2–25.5%)
	Late Incident (> 48h)	205	0 (0.0%)	0	0	0	0	[0.0%, 0.0%]	0 / 408 (0.0%)
	All Incident Sepsis-3	**408**	**151 (37.0%)**	**150**	**150**	**1**	**151**	**[36.8%, 37.0%]**	**[150, 151] / 408 (36.8–37.0%)**
MetaVision Sensitivity: Earliest Rx	Early Incident (≤ 24h)	47	47 (100.0%)	0	0	38	38	[0.0%, 80.9%]	[0, 38] / 408 (0.0–9.3%)
(Pre-Exposure Sensitivity: Earliest Rx $T_{\text{Rx,start}}$)	Intermediate (24–48h)	156	104 (66.7%)	0	0	9	9	[0.0%, 5.8%]	[0, 9] / 408 (0.0–2.2%)
	Late Incident (> 48h)	205	0 (0.0%)	0	0	0	0	[0.0%, 0.0%]	0 / 408 (0.0%)
	All Incident Sepsis-3	**408**	**151 (37.0%)**	**0**	**0**	**47**	**47**	**[0.0%, 11.5%]**	**[0, 47] / 408 (0.0–11.5%)**
Broad Historical MIMIC-III	Early Incident (≤ 24h)	2,641	102 (3.9%)	0	0	74	74	[0.00%, 2.80%]	[0, 74] / 3,581 (0.00–2.07%)
(Date-Level Proxy: Earliest Rx $T_{\text{Rx,start}}$)	Intermediate (24–48h)	361	235 (65.1%)	1	1	36	37	[0.28%, 10.25%]	[1, 37] / 3,581 (0.03–1.03%)
	Late Incident (> 48h)	579	0 (0.0%)	0	0	0	0	[0.00%, 0.00%]	[0, 0] / 3,581 (0.00%)
	All Incident Sepsis-3	**3,581**	**337 (9.4%)**	**1**	**1**	**110**	**111**	**[0.03%, 3.10%]**	**[1, 111] / 3,581 (0.03–3.10%)**