

Данные материалы были использованы для подготовки команды Казахстанского филиала МГУ на ИМС-2018, где ребята выступили очень успешно: Бекмаганбетов Бекарыс (ММ-2) взял золото (6 задач из 10), Аскергалиев Ануар (ВМК-2) взял серебро (4 задачи из 10), а Журавская Александра (ВМК-4) получила бронзовую награду (2 задачи из 10).

В брошюре собраны 219 задач из различных источников: задачи ИМС, задачи Putnam, Putnam And Beyond (Andreescu), Задачи студенческих олимпиад (Садовничий, Григорьян, Конягин) и другие.

Содержание

1 Telescope	3
1.1 Problems	3
1.2 Hints	4
2 Recurrent	5
2.1 Problems	5
2.2 Hints	6
3 Linear algebra	7
3.1 Problems	7
3.2 Hints	8
4 Polynomial	9
4.1 Problems	9
4.2 Hints	10
5 Calculus	11
5.1 Problems	11
5.2 Hints	12
6 Series	13
6.1 Problems	13
6.2 Hints	15
7 Linear algebra	16
7.1 Problems	16
7.2 Hints	18
8 Calculus	20
8.1 Problems	20
8.2 Hints	22
9 Sequences and series	23
9.1 Problems	23
9.2 Hints	26
10 Linear algebra	27
10.1 Problems	27
10.2 Hints	30
11 Number Theory	32
11.1 Problems	32

12 Train	34
12.1 Problems	34
12.2 Hints	36

1 Telescope

1.1 Problems

25.06.2018

1. (putnam-and-beyond-365) Find the sum $\frac{1}{\sqrt{1+\sqrt{2}}} + \frac{1}{\sqrt{2+\sqrt{3}}} + \cdots + \frac{1}{\sqrt{n+\sqrt{n+1}}}$.
2. (putnam-and-beyond-365) Let $a_0 = 1$, $a_1 = 3$, $a_{n+1} = \frac{a_n^2+1}{2}$, for $n \geq 1$. Prove that

$$\frac{1}{a_0+1} + \frac{1}{a_1+1} + \cdots + \frac{1}{a_n+1} + \frac{1}{a_{n+1}-1} = 1,$$

for $n \geq 1$.

3. (putnam-and-beyond-368) Let

$$a_n = \frac{4n + \sqrt{4n^2 - 1}}{\sqrt{2n+1} - \sqrt{2n-1}}$$

for $n \geq 1$. Prove that $a_1 + a_2 + \cdots + a_{40}$ is a positive integer.

4. (putnam-and-beyond-366) Let λ be a root of unity. Prove that

$$\lambda^{-1} = \sum_{n=0}^{\infty} \lambda^n (1 - \lambda)(1 - \lambda^2)(1 - \lambda^3) \cdots (1 - \lambda^n)$$

with the convention that the 0th term of the series is 1.

5. (putnam-2016-B1) Let x_0, x_1, x_2, \dots be the sequence such that $x_0 = 1$ and for $n \geq 0$,

$$x_{n+1} = \ln(e^{x_n} - x_n).$$

Show that the infinite series $x_0 + x_1 + x_2 + \dots$ converges and find its sum.

6. (putnam-2014-A3) Let $a_0 = \frac{5}{2}$ and $a_k = a_{k-1}^2 - 2$ for $k \geq 1$. Compute

$$\prod_{k=0}^{\infty} \left(1 - \frac{1}{a_k}\right)$$

in closed form.

1.2 Hints

1. (putnam-and-beyond-365)

$$\frac{1}{\sqrt{n} + \sqrt{n+1}} = \sqrt{n+1} - \sqrt{n}$$

2. (putnam-and-beyond-365)

$$2(a_{k+1} - 1) = (a_k - 1)(a_k + 1)$$

3. (putnam-and-beyond-368) $a = \sqrt{2n-1}$, $b = \sqrt{2n+1}$.

4. (putnam-and-beyond-369) Sum $a_{n+1} + a_n$.

5. (putnam-2016-B1) $e^x \geq x + 1$.

6. (putnam-2014-A3) $a_k + 1 = (a_{k-1} - 1)(a_{k-1} + 1)$, $a_k^2 - 4 = a_{k-1}^2(a_{k-1}^2 - 4)$.

2 Recurrent

2.1 Problems

26.06.2018

1. (putnam-and-beyond-299) Define the sequence $(a_n)_{n \geq 0}$ by $a_0 = 0$, $a_1 = 1$, $a_2 = 2$, $a_3 = 6$, and

$$a_{n+4} = 2a_{n+3} + a_{n+2} - 2a_{n+1} - a_n,$$

for $n \geq 0$. Prove that n divides a_n for all $n \geq 1$.

2. (putnam-and-beyond-302) A sequence u_n is defined by $u_0 = 2$, $u_1 = \frac{5}{2}$, $u_{n+1} = u_n(u_{n-1}^2 - 2) - u_1$, for $n \geq 1$. Prove that for all positive integers n ,

$$[u_n] = 2^{\frac{2^n - (-1)^n}{3}}.$$

3. (putnam-and-beyond-305) Find the general term of the sequence given by $x_0 = 3$, $x_1 = 4$, and

$$(n+1)(n+2)x_n = 4(n+1)(n+3)x_{n-1} - 4(n+2)(n+3)x_{n-2},$$

$n \geq 2$.

4. (putnam-and-beyond-307) Define the sequence (a_n) recursively by $a_1 = 1$ and

$$a_{n+1} = \frac{1 + 4a_n + \sqrt{1 + 24a_n}}{16}$$

for all $n \geq 1$. Find an explicit formula for a_n in terms of n .

5. (putnam-2015-A2) Let $a_0 = 1$, $a_1 = 2$, and $a_n = 4a_{n-1} - 1 - a_{n-2}$ for $n \geq 2$. Find an odd prime factor of a_{2015} .
6. (putnam-and-beyond-306) Let (x_n) , $n \geq 0$ be defined by the recurrence relation $x_{n+1} = ax_n + bx_{n-1}$, with $x_0 = 0$.

- Show that the expression

$$x_n^2 - x_{n-1}x_{n+1}$$

depends only on b and x_1 , but not on a .

- Show that the expression

$$f_n^2 - f_{n-1}f_{n+1}$$

where $f_n = f_{n-1} + f_{n-2}$, $f_0 = 0$, $f_1 = 1$.

7. (putnam-2017-A2) Let $Q_0(x) = 1$, $Q_1(x) = x$, and $Q_n(x) = \frac{(Q_{n-1}(x))^2 - 1}{Q_{n-2}(x)}$ for all $n \geq 2$. Show that, whenever n is a positive integer, $Q_n(x)$ is equal to a polynomial with integer coefficients.

2.2 Hints

1. (putnam-and-beyond-299) $a_n = n * f_n$.
2. (putnam-and-beyond-302) $u_n = 2^{x_n} + 2^{-x_n}$.
3. (putnam-and-beyond-305) $y_n = \frac{x_n}{n+3}$.
4. (putnam-and-beyond-307) $b_n = \sqrt{1 + 24a_n}$.
5. (putnam-2015-A2) $(x^5 + y^5)|(x^{2015} + y^{2015})$.
6. (putnam-and-beyond-306) Induction.
7. (putnam-2017-A2) See putnam-and-beyond-306. Note that $Q_n(x)$ has linear recurrent relation by $Q_{n-1}(x)$ and $Q_{n-2}(x)$.

3 Linear algebra

3.1 Problems

27.06.2018

1. (putnam-and-beyond-215) Let A, B, C, D be $n \times n$ matrices such that $AC = CA$. Prove that

$$\det \begin{pmatrix} A & B \\ C & D \end{pmatrix} = \det(AD - CB).$$

2. (putnam-2015-A6) Let n be a positive integer. Suppose that A, B , and M are $n \times n$ matrices with real entries such that $AM = MB$, and such that A and B have the same characteristic polynomial. Prove that $\det(A - MX) = \det(B - XM)$ for every $n \times n$ matrix X with real entries.

3. (putnam-and-beyond-216) Let X, Y be $n \times n$ matrices. Prove that

$$\det(I_n - XY) = \det(I_n - YX).$$

4. (putnam-and-beyond) Let A be $n \times n$ matrices. Prove that

$$\det(I_n + A^2) \geq 0$$

5. (putnam-and-beyond-217) Let A and B be $n \times n$ matrices that commute. Prove that if $\det(A + B) \geq 0$, then $\det(A^k + B^k) \geq 0$ for all $k \geq 1$.

6. (putnam-and-beyond-226) Let A, B, C be $n \times n$ matrices, $n \geq 1$, satisfying

$$ABC + AB + BC + AC + A + B + C = O_n.$$

7. (putnam-and-beyond-230) Let A and B be $n \times n$ matrices, $n \geq 1$, satisfying $AB - B^2A^2 = I_n$ and $A^3 + B^3 = O_n$. Prove that $BA - A^2B^2 = I_n$.

8. A is orthogonal matrix and $I + A$ is non singular. Prove that $(I + A)^{-1}(I - A)$ is skew symmetric. (косо-симметрическая)

9. A is orthogonal matrix, where n is even. Suppose that $\det(A) = -1$. Show that $\det(I - A) = 0$.

10. Let A, B is two invertible $n \times n$ real matrices. Assume that $A + B$ is invertible. Then prove that $A^{-1} + B^{-1}$ is also invertible.

11. Let X be any arbitrary $n \times n$ square matrix, n is even. Prove that

$$\text{tr}(X^n) \geq n \det(X).$$

12. Matrix $A, B, C, D \in R^{n \times n}$ such that

$$AC - BD = I_n, AD + BC = O_n$$

- Prove that $CA - DB = I_n, DA + CB = O_n$
- $\det(AC) \geq 0, (-1)^n \det(BD) \geq 0$.

3.2 Hints

1. (putnam-and-beyond-215) If A is invertable, then use matrix row transformation. If A is not invertable, $A_t = A + tI_n$ is invertable for all $t \neq -\lambda_k$. Checking equation is polynomial equation.
2. (putnam-2015-A6) If A and B is invertable, then checked it. If A or B is not invertable, $A_t = A + tI_n$, $B_t = B + tI_n$ is invertable for all $t \neq -\lambda_k$.
3. (putnam-and-beyond-216) Use problem 215.
4. (putnam-and-beyond) $I_n + A^2 = (I_n + iA)(I_n - iA)$, where $i = \sqrt{-1}$.
5. (putnam-and-beyond-217) Complex roots of $x^k = 1$ divides into complex conjugate pairs.
6. (putnam-and-beyond-226) If $XY = I$ then X, Y commute.
7. (putnam-and-beyond-230) $(A + iB^2)(B + iA^2) = I_n$.
8. B is skew symmetric $\Leftrightarrow B^T = -B$.
9. $\det(AB) = \det(A)\det(B) = \det(BA)$.
10. $\det(A^{-1} + B^{-1}) \neq 0$.
11. $\det(X) = \lambda_1 \cdot \lambda_2 \cdots \lambda_n$, $\text{tr}(X^n) = \lambda_1^n + \lambda_2^n + \cdots + \lambda_n^n$.
12. $(A + iB)(C + iD) = I_n$.

4 Polynomial

4.1 Problems

28.06.2018

1. Про многочлен $f(x) = x^{10} + a_9x^9 + \dots + a_0$ известно, что $f(1) = f(-1)$, $f(2) = f(-2)$, $f(3) = f(-3)$, $f(4) = f(-4)$, $f(5) = f(-5)$. Докажите, что $f(x) = f(-x)$ для любого действительного x .
2. Известно, что $P(x)$ – многочлен степени n такой, что для всех $t \in \{1, 2, 2^2, \dots, 2^n\}$ верно соотношение $P(t) = \frac{1}{t}$. Найдите $P(0)$.
3. Известно, что для многочлена степени n верно, что $f(0) = 1$, $f(1) = 2$, $f(2) = 4$, \dots , $f(n) = 2^n$. Найдите $f(n+1)$.
4. Пусть x, y – натуральные числа такие, что:

$$A = \frac{x^2 + y^2}{xy + 1}$$

– целое. Найдите все возможные значения A .

5. Пусть x, y – натуральные числа такие, что выражение $A = \frac{x^2 + y^2 + 1}{xy}$ – целое. Найдите все возможные значения A .
6. Произведение квадратных трехчленов $x^2 + a_1x + b_1$, $x^2 + a_2x + b_2$, \dots , $x^2 + a_nx + b_n$ равно многочлену $P(x) = x^{2n} + c_1x^{2n-1} + c_2x^{2n-2} + \dots + c_{2n-1}x + c_{2n}$, где коэффициенты c_1, c_2, \dots, c_{2n} положительны. Докажите, что для некоторого k ($1 \leq k \leq n$) коэффициенты a_k и b_k положительны.
7. (putnam-and-beyond-169) Let $P(x)$ be a polynomial with all roots real and distinct and such that none of its zeros is equal to 0. Prove that the polynomial $x^2P''(x) + 3xP'(x) + P(x)$ also has all roots real and distinct.
8. (putnam-and-beyond-172) Let $P(x)$ be a polynomial of degree $n > 3$ whose zeros $x_1 < x_2 < x_3 < \dots < x_{n-1} < x_n$ are real. Prove that

$$P' \left(\frac{x_1 + x_2}{2} \right) P' \left(\frac{x_{n-1} + x_n}{2} \right) \neq 0.$$

9. (putnam-and-beyond-173) Let a_1, a_2, \dots, a_n be positive real numbers. Prove that the polynomial $P(x) = x^n - a_1x^{n-1} - a_2x^{n-2} - \dots - a_n$ has a unique positive zero.
10. (putnam-and-beyond-174) Prove that the zeros of the polynomial

$$P(z) = z^7 + 7z^4 + 4z + 1$$

lie inside the disk of radius 2 centered at the origin.

11. (putnam-and-beyond-175) For $a \neq 0$ a real number and $n > 2$ an integer, prove that every nonreal root z of the polynomial equation $x^n + ax + 1 = 0$ satisfies the inequality

$$|z| \geq \sqrt[n]{\frac{1}{n-1}}.$$

12. Prove that for any distinct integers a_1, a_2, \dots, a_n the polynomial $P(x) = (x - a_1)(x - a_2) \cdots (x - a_n) - 1$ cannot be written as a product of two nonconstant polynomials with integer coefficients.

4.2 Hints

1. Многочлен $g(x) = f(x) - f(-x)$.
2. Многочлен $F(t) = P(t) * t - 1$.
3. Многочлен $f(x) = 1 + \frac{x}{1} + \frac{x(x-1)}{1 \cdot 2} + \frac{x(x-1)(x-2)}{1 \cdot 2 \cdot 3} + \dots + \frac{x(x-1) \dots (x-n+1)}{1 \cdot 2 \cdot 3 \dots n}$.
4. Идея: прыжки Виета. Если $x = y$, то легко решить явно. Если существует решение (x, y) , где $x > y$, то существует решение (y, z) , где $y > z$ (теорема Виета для квадратного уравнения).
5. Идея: прыжки Виета.
6. Теорема Виета для c_{2n} и c_1 .
7. (putnam-and-beyond-169) Total differential. Rolle's theorem:
If a real-valued function f is continuous on a proper closed interval $[a, b]$, differentiable on the open interval (a, b) , and $f(a) = f(b)$, then there exists at least one c in the open interval (a, b) such that $f'(c) = 0$.
8. (putnam-and-beyond-172) $P'(x) = P(x) \left(\frac{1}{x-x_1} + \frac{1}{x-x_2} + \dots + \frac{1}{x-x_n} \right)$.
9. (putnam-and-beyond-173) $\frac{P(x)}{x^n}$.
10. (putnam-and-beyond-174) $\left| \frac{P(z)}{z^r} \right|$. Triangle inequality.
11. (putnam-and-beyond-175) $z = r(\cos t + i \sin t)$.
12. (putnam-and-beyond-185) $P(x) = Q(x)R(x)$. What about $Q(x) + R(x)$?

5 Calculus

5.1 Problems

29.06.2018

1. (PUTNAM-1987-B1) Evaluate

$$\int_0^1 \frac{\sqrt{\ln(9-x)}dx}{\sqrt{\ln(9-x)} + \sqrt{\ln(x+3)}}$$

2. (PUTNAM-1989-A2) Evaluate

$$\int_0^a \int_0^b e^{\max\{b^2x^2, a^2y^2\}} dy dx,$$

where a and b are positive.

3. (PUTNAM-1990-B1) Find all real-valued continuously differentiable functions f on the real line such that for all x ,

$$f(x)^2 = \int_0^x (f(t)^2 + f'(t)^2) dt + 1990.$$

4. (PUTNAM-1998-B1) Find the minimum value of

$$\frac{(x + \frac{1}{x})^6 - x^6 - \frac{1}{x^6} - 2}{(x + \frac{1}{x})^3 + (x^3 + \frac{1}{x^3})}$$

for $x > 0$.

5. (PUTNAM-1998-A3) Let f be a real function on the real line with continuous third derivative. Prove that there exists a point a such that

$$f(a)f'(a)f''(a)f'''(a) \geq 0.$$

6. (PUTNAM-2006-B5) For each continuous function $f : [0, 1] \rightarrow \mathbb{R}$, let

$$I(f) = \int_0^1 x^2 f(x) dx$$

$$J(f) = \int_0^1 x(f(x))^2 dx.$$

Find the maximum value of $I(f) - J(f)$ over all such functions f .

7. (PUTNAM-2005-B3) Find all differentiable functions $f : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ for which there is a positive real number a such that

$$f'\left(\frac{a}{x}\right) = \frac{x}{f(x)}$$

for all $x > 0$.

8. Let function $f(x)$ is continuous on \mathbb{R} and $f^3(x) + f(x) = x$. Calculate

$$I = \int_0^2 f(x) dx.$$

9. If $f(x)$ is continuous and differentiable on $[0; 1]$ with $f(0) = 0$, $f(1) = 1$. Evaluate minimum value of

$$\int_0^1 (f'(x))^2 dx.$$

5.2 Hints

1. (PUTNAM-1987-B1) Symmetric variable substitution $9 - x = t + 3$.
2. (PUTNAM-1987-B1) From double integral to repeated integral.
3. (PUTNAM-1990-B1) Differentiate the equation.
4. (PUTNAM-1998-B1) Simplify. $x + \frac{1}{x} \geq 2$ for $x > 0$.
5. (PUTNAM-1998-A3) Without lose of generalization $f(x) > 0$. Without lose of generalization $f'(x) > 0$. It $f(x) > 0$, $f'(x) > 0$ for all x , then $f''(x) > 0$.
6. (PUTNAM-2006-B5) Cauchy-Schwarz inequality for $(I(f))^2$.
7. (PUTNAM-2005-B3) Variable substitution $t = \frac{a}{x}$. Differentiate the equation.
8. Variable substitution $f(x) = y$.
9. Cauchy-Schwarz inequality.

6 Series

6.1 Problems

02.07.2018

1. (putnam-and-beyond-352) Show that the series

$$\frac{1}{1+x} + \frac{2}{1+x^2} + \frac{4}{1+x^4} + \cdots + \frac{2^n}{1+x^{2^n}} + \cdots$$

converges when $|x| > 1$, and in this case find its sum.

2. (putnam-and-beyond-355) Let $S = \{x_1, x_2, \dots, x_n, \dots\}$ be the set of all positive integers that do not contain the digit 9 in their decimal representation. Prove that

$$\sum_{n=1}^{\infty} \frac{1}{x_n} < 80$$

3. (putnam-and-beyond-357) Does the series

$$\sum_{n=1}^{\infty} \sin(\pi \sqrt{n^2 + 1})$$

converges?

4. (putnam-and-beyond-369) Prove that

$$\sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{1^2 - 2^2 + 3^2 - \cdots + (-1)^{k+1} k^2} = \frac{2n}{n+1}$$

5. (putnam-and-beyond-370) Prove that

$$\sum_{n=1}^{9999} \frac{1}{(\sqrt{n} + \sqrt{n+1})(\sqrt[4]{n} + \sqrt[4]{n+1})} = 9$$

6. (putnam-and-beyond-371) Let $a_n = \sqrt{1 + (1 + \frac{1}{n})^2} + \sqrt{1 + (1 - \frac{1}{n})^2}$, $n \geq 1$. Prove that

$$\frac{1}{a_1} + \frac{1}{a_2} + \cdots + \frac{1}{a_{20}}$$

is a positive integer.

7. (putnam-and-beyond-373) Let $a_n = 3n + \sqrt{n^2 - 1}$ and $b_n = 2(\sqrt{n^2 - n} + \sqrt{n^2 + n})$, $n \geq 1$. Show that

$$\sqrt{a_1 - b_1} + \sqrt{a_2 - b_2} + \cdots + \sqrt{a_{49} - b_{49}} = A + B\sqrt{2}$$

for some integer A and B .

8. Compute the product

$$\prod_{n=2}^{\infty} \left(1 + \frac{(-1)^n}{F_n^2}\right)$$

where F_n is the n -th Fibonacci number.

9. (putnam-and-beyond-378) Let x be a positive number less than 1. Compute the product

$$\prod_{n=1}^{\infty} (1 + x^{2^n})$$

10. (ВШЭ-2012) Сходится ли ряд

$$\sum_{n=1}^{\infty} \frac{1}{n^{\frac{1}{n}+1}}?$$

11. (Садовничий-7-1) Последовательность задается: $x_1 = a > 1$, $x_{n+1} = x_n^2 - x_n + 1$. Найдите

$$\sum_{n=1}^{\infty} \frac{1}{x_n}$$

12. (Садовничий-7-4) Вычислить:

$$\sum_{n=1}^{\infty} \frac{1}{n(n+1)2^n}$$

13. (Садовничий-7-7) Вычислить:

$$\frac{1}{1 \cdot 2 \cdot 3} + \frac{1}{4 \cdot 5 \cdot 6} + \frac{1}{7 \cdot 8 \cdot 9} + \dots$$

14. (putnam-2016-B6) Evaluate

$$\sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k} \sum_{n=0}^{\infty} \frac{1}{k2^n + 1}.$$

15. (putnam-2015-A3) Compute

$$\log_2 \left(\prod_{a=1}^{2015} \prod_{b=1}^{2015} (1 + e^{2\pi i ab/2015}) \right)$$

Here i is the imaginary unit (that is, $i^2 = -1$).

16. (putnam-2014-A3) Let $a_0 = \frac{5}{2}$ and $a_k = a_{k-1}^2 - 2$ for $k \geq 1$. Compute

$$\prod_{k=0}^{\infty} \left(1 - \frac{1}{a_k} \right)$$

in closed form.

17. (putnam-2011-A2) Let a_1, a_2, \dots and b_1, b_2, \dots be sequences of positive real numbers such that $a_1 = b_1 = 1$ and $b_n = b_{n-1}a_n - 2$ for $n = 2, 3, \dots$. Assume that the sequence (b_j) is bounded. Prove that

$$S = \sum_{n=1}^{\infty} \frac{1}{a_1 \dots a_n}$$

converges, and evaluate S .

18. (putnam-2001-B3) For any positive integer n , let $\langle n \rangle$ denote the closest integer to \sqrt{n} . Evaluate

$$\sum_{n=1}^{\infty} \frac{2^{\langle n \rangle} + 2^{-\langle n \rangle}}{2^n}.$$

19. (putnam-1999-A4) Sum the series

$$\sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \frac{m^2 n}{3^m (n3^m + m3^n)}.$$

20. (Алфутова-устинов-11-65) Вычислите суммы:

- $C_n^1 + 2C_n^2 + 3C_n^3 + \dots + nC_n^n$.
- $C_n^1 + 2^2 C_n^2 + 3^2 C_n^3 + \dots + n^2 C_n^n$.

6.2 Hints

1. (putnam-and-beyond-352) Add $\frac{1}{1-x}$.
2. (putnam-and-beyond-355) Double series $\sum_{x_j < 10^n} \frac{1}{x_j} = \sum_{1 \leq i \leq n} \sum_{10^{i-1} \leq j < 10^i} \frac{1}{x_j}$
3. (putnam-and-beyond-357) $\sin(\pi\sqrt{n^2+1}) = (-1)^n \sin(\frac{\pi}{\sqrt{n^2+1}+n})$
4. (putnam-and-beyond-369) $1^2 - 2^2 + 3^2 - \dots + (-1)^{k+1} k^2 = (-1)^{k+1} \frac{k(k+1)}{2}$
5. (putnam-and-beyond-370) $a_n = \sqrt[4]{n+1} - \sqrt[4]{n}$
6. (putnam-and-beyond-371) $a_n = \frac{1}{4}(b_{n+1} - b_n)$
7. (putnam-and-beyond-373) $a_k - b_k = \frac{1}{2} \left((\sqrt{k} - \sqrt{k+1}) - (\sqrt{k-1} - \sqrt{k}) \right)$
8. Let (x_n) , $n \geq 0$ be defined by the recurrence relation $x_{n+1} = ax_n + bx_{n-1}$, with $x_0 = 0$.

- Show that the expression

$$x_n^2 - x_{n-1}x_{n+1}$$

depends only on b and x_1 , but not on a .

- Evaluate

$$f_n^2 - f_{n-1}f_{n+1}$$

where $f_n = f_{n-1} + f_{n-2}$, $f_0 = 0$, $f_1 = 1$.

9. (putnam-and-beyond-378) Multiply by $(1+x)$.
10. (ВШЭ-2012) $n^{n+1} < 2n$
11. (Садовничий-7-1) $\frac{1}{x_n} = \frac{1}{x_{n+1}-1} - \frac{1}{x_n-1}$
12. (Садовничий-7-4) Интегрируем дважды степенной ряд $\sum_{n=1}^{\infty} z^n$.
13. (Садовничий-7-7)
14. (putnam-2016-B6) $\frac{1}{k+1} = \int_0^1 x^k dx$
15. (putnam-2015-A3) $z^n - 1 = (z - \omega)(z - \omega^2) \cdots (z - \omega^{n-1})$. $(1 + \omega^b) = \frac{1 - \omega^{2b}}{1 - \omega^b}$.
16. (putnam-2014-A3) $a_k + 1 = (a_{k-1} - 1)(a_{k-1} + 1)$, $a_k^2 - 4 = a_{k-1}^2(a_{k-1}^2 - 4)$.
17. (putnam-2011-A2) $S_m - S_{m-1} = \frac{1}{a_1 a_2 \cdots a_m}$, where $S_m = \frac{b_1 b_2 \cdots b_n}{(b_1+2)(b_2+2) \cdots (b_n+2)}$
18. (putnam-2001-B3) Since $(k - 1/2)^2 = k^2 - k + 1/4$ and $(k + 1/2)^2 = k^2 + k + 1/4$, we have that $\langle n \rangle = k$ if and only if $k^2 - k + 1 \leq n \leq k^2 + k$. Hence

$$\sum_{n=1}^{\infty} \frac{2^{\langle n \rangle} + 2^{-\langle n \rangle}}{2^n} = \sum_{k=1}^{\infty} \sum_{n, \langle n \rangle = k} \frac{2^{\langle n \rangle} + 2^{-\langle n \rangle}}{2^n} = \sum_{k=1}^{\infty} \sum_{n=k^2-k+1}^{k^2+k} \frac{2^k + 2^{-k}}{2^n} = \sum_{k=1}^{\infty} (2^k + 2^{-k})(2^{-k^2+k} - 2^{-k^2-k}) = 3.$$

19. (putnam-1999-A4) Denote the series by S , and let $a_n = 3^n/n$:

$$S = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \frac{1}{a_m(a_m + a_n)} = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \frac{1}{a_n(a_m + a_n)},$$

Thus

$$2S = \left(\sum_{n=1}^{\infty} \frac{n}{3^n} \right)^2.$$

$$S = 9/32.$$

20. (Алфутова-устинов-11-65) $f(x) = (1+x)^n$, $f'(x)$, $f''(x)$.

7 Linear algebra

7.1 Problems

03.07.2018

1. (putnam-and-beyond-199) Let M be an $n \times n$ complex matrix. Prove that there exist Hermitian matrices A and B such that $M = A + iB$. (A matrix X is called Hermitian if $X^T = \overline{X}$).
2. (putnam-and-beyond-200) Do there exist $n \times n$ matrices A and B such that $AB - BA = I_n$?
3. (putnam-and-beyond-201) Let A and B be 2×2 matrices with real entries satisfying $(AB - BA)^n = I_2$ for some positive integer n . Prove that n is even and $(AB - BA)^4 = I_2$.
4. (putnam-and-beyond-205) Let A and B be $n \times n$ matrices with real entries satisfying

$$\operatorname{tr}(AA^T + BB^T) = \operatorname{tr}(AB + A^T B^T).$$

Prove that $A = B^T$.

5. (putnam-and-beyond-206) Let x_1, x_2, \dots, x_n be arbitrary numbers ($n \geq 1$). Compute the determinant

$$\begin{vmatrix} 1 & 1 & \dots & 1 \\ x_1 & x_2 & \dots & x_n \\ x_1^2 & x_2^2 & \dots & x_n^2 \\ \dots & \dots & \dots & \dots \\ x_1^{n-1} & x_2^{n-1} & \dots & x_n^{n-1} \end{vmatrix}$$

6. (putnam-and-beyond-213) Let A and B be 3×3 matrices with real elements such that $\det A = \det B = \det(A + B) = \det(A - B) = 0$. Prove that $\det(xA + yB) = 0$ for any real numbers x and y .
7. (putnam-and-beyond-222) Let A and B be 2×2 matrices with integer entries such that $A, A + B, A + 2B, A + 3B$, and $A + 4B$ are all invertible matrices whose inverses have integer entries. Prove that $A + 5B$ is invertible and that its inverse has integer entries.
8. (putnam-and-beyond-243) Let A be the $n \times n$ matrix whose i, j entry is $i + j$ for all $i, j = 1, 2, \dots, n$. What is the rank of A ?
9. (putnam-and-beyond-244) For integers $n \geq 2$ and $0 \leq k \leq n - 2$, compute the determinant

$$\begin{vmatrix} 1^k & 2^k & 3^k & \dots & n^k \\ 2^k & 3^k & 4^k & \dots & (n+1)^k \\ 3^k & 4^k & 5^k & \dots & (n+2)^k \\ \dots & \dots & \dots & \dots & \dots \\ n^k & (n+1)^k & (n+2)^k & \dots & (2n-1)^k \end{vmatrix}$$

10. (putnam-and-beyond-248) Let $A : V \rightarrow W$ and $B : W \rightarrow V$ be linear maps between finite-dimensional vector spaces. Prove that the linear maps AB and BA have the same set of nonzero eigenvalues, counted with multiplicities.
11. (putnam-and-beyond-249) Let A, B be 2×2 matrices with integer entries, such that $AB = BA$ and $\det B = 1$. Prove that if $\det(A^3 + B^3) = 1$, then $A^2 = O_2$.
12. (putnam-and-beyond-258) Let A and B be 2×2 matrices with determinant equal to 1. Prove that

$$\operatorname{tr}(AB) - (\operatorname{tr} A)(\operatorname{tr} B) + \operatorname{tr}(AB^{-1}) = 0.$$

13. (putnam-and-beyond-259) Find the 2×2 matrices with real entries that satisfy the equation

$$X^3 - 3X^2 = \begin{pmatrix} -2 & -2 \\ -2 & -2 \end{pmatrix}$$

14. (putnam-and-beyond-261) Let A and B be 3×3 matrices. Prove that

$$3 \det(AB - BA) = \operatorname{tr}((AB - BA)^3)$$

15. (putnam-2015-B3) Let S be the set of all 2×2 real matrices

$$M = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

whose entries a, b, c, d (in that order) form an arithmetic progression. Find all matrices M in S for which there is some integer $k > 1$ such that M^k is also in S .

16. (putnam-2014-A2) Let A be the $n \times n$ matrix whose entry in the i -th row and j -th column is

$$\frac{1}{\min(i, j)}$$

for $1 \leq i, j \leq n$. Compute $\det(A)$.

17. (ncumc-2017-5) Find the maximal set of points in \mathbb{C} such that there are no complex Hermitian positively definite matrices of identical sizes A, B for which the point is an eigenvalue of matrix $(A + B)^{-1}(I + AB)$.

18. (Садовничий-2-1-11) Вычислить определитель

$$\begin{vmatrix} P(x) & P(x+1) & \dots & P(x+n) \\ P'(x) & P'(x+1) & \dots & P'(x+n) \\ \dots & \dots & \dots & \dots \\ P^{(n)}(x) & P^{(n)}(x+1) & \dots & P^{(n)}(x+n) \end{vmatrix}$$

где $P(x) = x(x+1)(x+2) \cdot (x+n)$.

19. (Садовничий-2-1-20) Пусть A, B — квадратные матрицы порядка 2017. Доказать, что если $AB = 0$, то хотя бы одна из матриц $A + A^T$ или $B + B^T$ вырождена.

20. (Садовничий-2-1-21) Пусть A, B, C — квадратные матрицы порядка $n \times n$. Доказать, что

$$\operatorname{rg}(AB) + \operatorname{rg}(BC) \leq \operatorname{rg}(B) + \operatorname{rg}(ABC)$$

21. (Садовничий-2-1-27) Пусть A, B — квадратные матрицы порядка $n \times n$. Верно ли, что матрицы AB и BA имеют одинаковый характеристический многочлен?

7.2 Hints

1. (putnam-and-beyond-199) $\overline{M^T} = \overline{A^T} - i\overline{B^T} = A - iB$.
2. (putnam-and-beyond-200) $\text{tr}(AB) = \text{tr}(BA)$.
3. (putnam-and-beyond-201) $(AB - B)^2 = \begin{pmatrix} a & b \\ c & -a \end{pmatrix}^2 = kI_2$, where k is a root of unity.
4. (putnam-and-beyond-205) $\text{tr}(XX^T)$ is the sum of the squares of the entries of X .

$$\text{tr}[(A - B^T)(A - B^T)^T] = 0$$

5. (putnam-and-beyond-206) The Vandermonde determinant is equal to $\prod_{i>j}(x_i - x_j)$.
6. (putnam-and-beyond-213) $\det(xA + yB) = a_0(x)y^3 + a_1(x)y^2 + a_2(x)y + a_3(x)$. Idea: $\det(xA + xB)$, $\det(xA - xB)$.
7. (putnam-and-beyond-222) $\det C = \pm 1$, $P(x) = \det(A + xB)$.
8. (putnam-and-beyond-243) $n \geq 2$, $\text{rank} = 2$
9. (putnam-and-beyond-244) The polynomials $P_j(x) = (x + j)^k$, $j = 0, 1, \dots, n - 1$, lie in the $(k + 1)$ -dimensional real vector space of polynomials of degree at most k .
10. (putnam-and-beyond-248) Let A, B, C, D be $n \times n$ matrices such that $AC = CA$. Prove that

$$\begin{vmatrix} A & B \\ C & D \end{vmatrix} = \det(AD - CB).$$

11. (putnam-and-beyond-249)

$$A^3 + B^3 = (C^3 + I)B^3.$$

$$C^3 + I = (C + I)(C + \varepsilon I)(C + \varepsilon^2 I).$$

$$P(-1)P(-\varepsilon)P(-\varepsilon^2) = 1$$

where $P(x) = x^2 - mx + n$.

12. (putnam-and-beyond-258) $P_A(\lambda) = \det(\lambda I_n - A)$, $P_A(A) = 0_n$.
13. (putnam-and-beyond-259) Get \det . If $\det X = 0$, then $X^2 = (\text{tr}(X))X$ and $\text{tr}(X) = 2$ or $\text{tr}(X) = -1$.
If $\det(X - 3I) = 0$, then 3 is eigenvalue. $\det(X^3 - 3X + 4I_2) = 0$.
14. (putnam-and-beyond-261) $(AB - BA)^3 - c_1(AB - BA)^2 + c_2(AB - BA) - c_3I_3 = O_3$ Take trace.
15. (putnam-2015-B3) If a, b, c, d are in arithmetic progression, then we may write

$$a = r - 3s, b = r - s, c = r + s, d = r + 3s$$

for some r, s . If $s = 0$, then clearly all powers of M are in S . Also, if $r = 0$, then one easily checks that M^3 is in S .

We now assume $rs \neq 0$, and show that in that case M cannot be in S . First, note that the characteristic polynomial of M is $x^2 - 2rx - 8s^2$, and since M is nonsingular (as $s \neq 0$), this is also the minimal polynomial of M by the Cayley-Hamilton theorem. By repeatedly using the relation $M^2 = 2rM + 8s^2I$, we see that for each positive integer, we have $M^k = t_k M + u_k I$ for unique real constants t_k, u_k (uniqueness follows from the independence of M and I). Since M is in S , we see that M^k lies in S only if $u_k = 0$.

On the other hand, we claim that if $k > 1$, then $rt_k > 0$ and $u_k > 0$ if k is even, and $t_k > 0$ and $ru_k > 0$ if k is odd (in particular, u_k can never be zero). The claim is true for $k = 2$ by the relation $M^2 = 2rM + 8s^2I$. Assuming the claim for k , and multiplying both sides of the relation $M^k = t_kM + u_kI$ by M , yields

$$M^{k+1} = t_k(2rM + 8s^2I) + u_kM = (2rt_k + u_k)M + 8s^2t_kI,$$

implying the claim for $k + 1$.

16. (putnam-2014-A2) Sub n from $(n - 1)$ row.
17. (ncumc-2017-5) Let c be an eigenvalue of the operator in question, i.e. $(A + B)^{-1}(I + AB)x = cx$ for some non-zero vector x and some complex number c . Then, $x + ABx = c(Ax + Bx)$.

Mark $Bx = y$. Hence, $(y, x) > 0$, and it is the only condition for x, y . Equation rewritten in the form $A(y - cx) = cy - x$. Moreover, $(cy - x, y - cx) > 0$ due to the fact that A be positively definite. It is also possible that $y = cx$, $cy = x$. this takes place for $x = y$, $c = 1$. Let us introduce a notation $c = a + bi$. Then,

$$a((x, x) + (y, y)) + bB((y, y) - (x, x)) - (1 + a^2 + b^2)(x, y) > 0.$$

Consequently, $a > 0$, and for any $a > 0$ and any b , one can find matrices and almost orthogonal vectors x, y of identical lengths such that the inequality takes place.

18. (Садовничий-2-1-11) Умножим k -ю строку на $\frac{(x+n)^{k-1}}{(k-1)!}$. Ряд Тейлора.
19. (Садовничий-2-1-20)
20. (Садовничий-2-1-21) $rg(A + B) \leq rg(A) + rg(B)$, $rg(A + B) \geq rg(AB) + n$.
21. (Садовничий-2-1-27)

8 Calculus

8.1 Problems

04.07.2018

1. (imc-1994-1-2) Let $f \in C^1(a, b)$, $\lim_{x \rightarrow a+} f(x) = +\infty$, $\lim_{x \rightarrow b-} f(x) = -\infty$ and

$$f'(x) + f^2(x) \geq -1$$

for $x \in (a, b)$. Prove that $b - a \geq \pi$ and give an example where $b - a = \pi$.

2. (imc-1994-2-1) Let $f \in C^1[a, b]$, $f(a) = 0$ and suppose that $\lambda \in \mathbb{R}$, $\lambda > 0$, is such that $|f'(x)| \leq \lambda|f(x)|$ for all $x \in [a, b]$. Is it true that $f(x) = 0$ for all $x \in [a, b]$?
3. (imc-1995-1-2) Let f be a continuous function on $[0, 1]$ such that for every $x \in [0, 1]$ we have $\int_x^1 f(t)dt \geq \frac{1-x^2}{2}$. Show that $\int_0^1 f^2(t)dt \geq \frac{1}{3}$.
4. (imc-1996-1-2) Evaluate the definite integral

$$\int_{-\pi}^{\pi} \frac{\sin(nx)}{(1+2^x)\sin x} dx$$

where n is a natural number.

5. (imc-1998-1-3) Let $f(x) = 2x(1-x)$, $x \in \mathbb{R}$. Define $f_n(x) = f(f(\dots f(x)\dots))$ (n times).
- a Find $\lim_{n \rightarrow \infty} \int_0^1 f_n(x)dx$.
- b Compute $\int_0^1 f_n(x)dx$ for $n = 1, 2, \dots$
6. (imc-1998-1-4) The function $f : \mathbb{R} \rightarrow \mathbb{R}$ is twice differentiable and satisfies $f(0) = 2$, $f'(0) = -2$ and $f(1) = 1$. Prove that there exists a real number $\xi \in (0, 1)$ for which

$$f(\xi)f'(\xi) + f''(\xi) = 0.$$

7. (imc-1998-1-5) Let P be an algebraic polynomial of degree n having only real zeros and real coefficients. Prove that for every real x the following inequality holds:

$$(n-1)(P'(x))^2 \geq nP(x)P''(x).$$

8. (imc-1998-1-6) Let $f : [0, 1] \rightarrow \mathbb{R}$ be a continuous function with the property that for any x and y in the interval, $xf(y) + yf(x) \leq 1$. Show that

$$\int_0^1 f(x)dx \leq \frac{\pi}{4}.$$

9. (imc-1999-1-3) Suppose that a function $f : \mathbb{R} \rightarrow \mathbb{R}$ satisfies the inequality

$$\left| \sum_{k=1}^n 3^k (f(x+ky) - f(x-ky)) \right| \leq 1.$$

for every positive integer n and for all $x, y \in \mathbb{R}$. Prove that f is a constant function.

10. (imc-2000-1-1) Is it true that if $f : [0, 1] \rightarrow [0, 1]$ is

- a monotone increasing;
a monotone decreasing

then there exists an $x \in [0, 1]$ for which $f(x) = x$?

11. (imc-2002-1-2) Does there exist a continuously differentiable function $f : \mathbb{R} \rightarrow \mathbb{R}$ such that for every $x \in \mathbb{R}$ we have $f(x) > 0$ and $f'(x) = f(f(x))$?

12. (imc-2003-1-2) Evaluate the limit

$$\lim_{x \rightarrow +0} \int_x^{2x} \frac{\sin^m(t)}{t^n} dt$$

$m, n \in \mathbb{N}$

13. (imc-2004-2-2) Let $f, g : [a, b] \rightarrow [0, \infty)$ be continuous and nondecreasing functions such that for each $x \in [a, b]$ we have

$$\begin{aligned} \int_a^x \sqrt{f(t)} dt &\leq \int_a^x \sqrt{g(t)} dt \\ \int_a^b \sqrt{f(t)} dt &\leq \int_a^b \sqrt{g(t)} dt \end{aligned}$$

Prove that

$$\int_a^b \sqrt{1+f(t)} dt \leq \int_a^b \sqrt{1+g(t)} dt$$

14. (imc-2005-1-3) Let $f : \mathbb{R} \rightarrow [0, \infty)$ be a continuously differentiable function. Prove that

$$\left| \int_0^1 f^3(x) dx - f^2(0) \int_0^1 f(x) dx \right| \leq \max_{0 \leq x \leq 1} |f'(x)| \left(\int_0^1 f(x) dx \right)^2$$

15. (imc-2005-2-4) Prove that if $f : \mathbb{R} \rightarrow \mathbb{R}$ is three times differentiable, then there exists a real number $\xi \in (-1, 1)$ such that

$$\frac{f'''(\xi)}{6} = \frac{f(1) - f(-1)}{2} - f'(0).$$

16. (imc-2006-2-3) Compare $\tan(\sin x)$ and $\sin(\tan x)$ for all $x \in (0, \frac{\pi}{2})$.

17. (imc-2007-2-3) Let C be a nonempty closed bounded subset of the real line and $f : C \rightarrow C$ be a nondecreasing continuous function. Show that there exists a point $p \in C$ such that $f(p) = p$.

18. (imc-2009-2-2) Suppose $f : \mathbb{R} \rightarrow \mathbb{R}$ is a two times differentiable function satisfying $f(0) = 1$, $f'(0) = 0$, and for all $x \in [0, \infty)$,

$$f''(x) - 5f'(x) + 6f(x) \geq 0.$$

Prove that for all $x \in [0, \infty)$,

$$f(x) \geq 3e^{2x} - 2e^{3x}.$$

19. (imc-2010-1-1) Let $0 < a < b$. Prove that

$$\int_a^b (x^2 + 1)e^{-x^2} dx \geq e^{-a^2} - e^{-b^2}$$

20. (imc-2013-1-2) Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a twice differentiable function. Suppose $f(0) = 0$. Prove that there exists $\xi \in (-\frac{\pi}{2}, \frac{\pi}{2})$ such that

$$f''(\xi) = f(\xi)(1 + 2 \tan^2(\xi)).$$

21. (imc-2014-2-3) Let $f(x) = \frac{\sin x}{x}$, for $x > 0$, and let n be a positive integer. Prove that

$$\left| f^{(n)}(x) \right| < \frac{1}{n+1},$$

where $f^{(n)}$ denotes the n -th derivative of f .

22. (imc-2016-2-2) Today, Ivan the Confessor prefers continuous functions $f : [0, 1] \rightarrow \mathbb{R}$ satisfying

$$f(x) + f(y) \geq |x - y|$$

for all pairs $x, y \in [0, 1]$. Find the minimum of $\int_0^1 f(x) dx$ over all preferred functions.

8.2 Hints

1. (imc-1994-1-2) $\arctg(f(x))$.
2. (imc-1994-2-1) $g(x) = \ln f(x) - \lambda x$ is not increasing. $f(x) \geq e^{\lambda(x-y)} f(y)$.
3. (imc-1995-1-2) $\int_0^1 (f(t) - t)^2 dt \geq 0$.
4. (imc-1996-1-2) $I_n = I_{n-2}$.
5. (imc-1998-1-3) $x_n = f_n(x_0)$ has 0 in limit. $f_n(x) = \frac{1}{2} - 2^{2^n-1} \left(x - \frac{1}{2}\right)^{2^n}$.
6. (imc-1998-1-4) $g(x) = \frac{1}{2}f(x)^2 + f'(x)$. $h(x) = \frac{x}{2} - \frac{1}{f(x)}$.
7. (imc-1998-1-5) Find $P'(x)/P(x)$ and $P''(x)/P(x)$.
8. (imc-1998-1-6) $I + \int_0^{\frac{\pi}{2}} f(\sin\theta)\cos\theta d\theta = \int_0^{\frac{\pi}{2}} f(\cos\theta)\sin\theta d\theta$.
9. (imc-1999-1-3) $3^n(f(x+ny) - f(x-ny)) \leq 2$.
10. (imc-2000-1-1) yes, no.
11. (imc-2002-1-2) $f(x) < f(0) + x \cdot f'(0) = (1+x)f'(0)$
12. (imc-2003-1-2) $f(x)/x$ is decreasing.
13. (imc-2004-2-2) The length of the graph of F is \geq the length of the graph of G . This is clear since both functions are convex.
14. (imc-2005-1-3) Integrate $-Mf(x) \leq f(x)f'(x) \leq Mf(x)$
15. (imc-2005-2-4) $g(x) = -\frac{f(-1)}{2}x^2(x-1) - f(0)(x^2-1) + \frac{f(1)}{2}x^2(x+1) - f'(0)x(x-1)(x+1)$
16. (imc-2006-2-3) $f(x) = \tan(\sin x) - \sin(\tan x)$ increase.
17. (imc-2007-2-3) Suppose $f(x) \neq x$. Let $[a, b]$ be the smallest closed interval that contains C .
18. (imc-2009-2-2) $g(x) = f'(x) - 2f(x)$, $f'(x) - 2f(x) \geq -2e^{3x}$
19. (imc-2010-1-1) $f(x) = \int_0^x (t^2 + 1)e^{-t^2} dt$, $g(x) = -e^{-x^2}$.
20. (imc-2013-1-2) $g(x) = f(x)\cos x$
21. (imc-2014-2-3) $g(x) = x^{n+1} \left(f^n(x) - \frac{1}{n+1} \right)$
22. (imc-2016-2-2) Triangle inequality.

9 Sequences and series

9.1 Problems

05.07.2018

1. (imc-1995-2-2) Let $\{b_n\}_{n=0}^{\infty}$ be a sequence of positive real numbers such that $b_0 = 1$,

$$b_n = 2 + \sqrt{b_{n-1}} - 2\sqrt{1 + \sqrt{b_{n-1}}}.$$

Calculate

$$\sum_{n=1}^{\infty} b_n 2^n.$$

2. (imc-1996-2-5) Prove that

$$\lim_{x \rightarrow +\infty} \sum_{n=1}^{\infty} \frac{nx}{(n^2 + x)^2} = \frac{1}{2}.$$

3. (imc-1997-1-1) Let $\{\varepsilon_n\}_{n=0}^{\infty}$ be a sequence of positive real numbers, such that $\lim_{n \rightarrow \infty} \varepsilon_n = 0$. Find

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n \ln \left(\frac{k}{n} + \varepsilon_n \right).$$

4. Дискретное преобразование Абеля

$$\sum_{k=m}^n a_k b_k = a_n B_n - a_m B_{m-1} - \sum_{k=m}^{n-1} (a_{k+1} - a_k) B_k$$

5. (imc-1999-1-2) Does there exist a bijective map $\pi : N \rightarrow N$ such that

$$\sum_{n=1}^{\infty} \frac{\pi(n)}{n^2} < \infty?$$

6. (imc-1999-2-3) Assume that $x_1, \dots, x_n \geq -1$ and $\sum_{i=1}^n x_i^3 = 0$. Prove that $\sum_{i=1}^n x_i \leq \frac{n}{3}$.

7. (imc-2001-1-3) Find

$$\lim_{t \rightarrow 1-0} (1-t) \sum_{n=1}^{\infty} \frac{t^n}{1+t^n}.$$

8. (imc-2002-1-3) Let n be a positive integer and let $a_k = \frac{1}{C_n^k}$, $b_k = 2^{k-n}$ for $k = 1, \dots, n$. Show that

$$\frac{a_1 - b_1}{1} + \frac{a_2 - b_2}{2} + \dots + \frac{a_n - b_n}{n} = 0.$$

9. (imc-2003-1-1) Let a_1, a_2, \dots be a sequence of real numbers such that $a_1 = 1$ and $a_{n+1} > \frac{3}{2}a_n$ for all n . Prove that the sequence

$$\frac{a_n}{\left(\frac{3}{2}\right)^{n-1}}$$

has a finite limit or tends to infinity.

10. (imc-2003-2-6) Let $(a_n)_{n \in N}$ be the sequence defined by $a_0 = 1$,

$$a_{n+1} = \frac{1}{n+1} \sum_{k=0}^n \frac{a_k}{n-k+2}$$

Find the limit

$$\lim_{n \rightarrow \infty} \sum_{k=0}^n \frac{a_k}{2^k}$$

11. (imc-2008-2-3) Let n be a positive integer. Prove that 2^{n-1} divides

$$\sum_{0 \leq k < n/2} C_n^{2k+1} 5^k$$

12. (imc-2010-1-2) Compute the sum of the series

$$\frac{1}{1 \cdot 2 \cdot 3 \cdot 4} + \frac{1}{5 \cdot 6 \cdot 7 \cdot 8} + \cdots$$

13. (imc-2010-1-3) Define the sequence x_1, x_2, \dots inductively by $x_1 = \sqrt{5}$, $x_{n+1} = x_n^2 - 2$ for $n \geq 1$. Compute

$$\lim_{n \rightarrow \infty} \frac{x_1 \cdot x_2 \cdot x_3 \cdots x_n}{x_{n+1}}$$

14. (imc-2010-2-1) A sequence x_1, x_2, \dots of real numbers satisfies $x_{n+1} = x_n \cos x_n$ for all $n \geq 1$. Does it follow that this sequence converges for all initial values x_1 ?

A sequence y_1, y_2, \dots of real numbers satisfies $y_{n+1} = y_n \sin y_n$ for all $n \geq 1$. Does it follow that this sequence converges for all initial values y_1 ?

15. (imc-2010-2-2) Let a_0, a_1, \dots, a_n be positive real numbers such that $a_{k+1} - a_k \geq 1$ for all $k = 0, 1, \dots, n-1$. Prove that

$$1 + \frac{1}{a_0} \left(1 + \frac{1}{a_1 - a_0}\right) \cdots \left(1 + \frac{1}{a_n - a_0}\right) \leq \left(1 + \frac{1}{a_0}\right) \left(1 + \frac{1}{a_1}\right) \cdots \left(1 + \frac{1}{a_n}\right)$$

16. (imc-2011-2-1) Let $\{a_n\}_{n=0}^{\infty}$ be a sequence with $\frac{1}{2} < a_n < 1$ for all $n \geq 0$. Define the sequence $\{x_n\}_{n=0}^{\infty}$ by $x_0 = a_0$ and

$$x_{n+1} = \frac{a_{n+1} + x_n}{1 + a_{n+1}x_n}$$

What are the possible values of $\lim_{n \rightarrow \infty} x_n$? Can such a sequence diverge?

17. (imc-2012-2-2) Define the sequence a_0, a_1, \dots inductively by $a_0 = 1$, $a_1 = \frac{1}{2}$,

$$a_{n+1} = \frac{na_n^2}{1 + (n+1)a_n}$$

for $n \geq 1$. Show that the series

$$\sum_{k=1}^{\infty} \frac{a_{k+1}}{a_k}$$

converges and determine its value.

18. (imc-2013-1-4) Let $n \geq 3$ and let x_1, \dots, x_n be nonnegative real numbers. Define $A = \sum_{i=1}^n x_i$, $B = \sum_{i=1}^n x_i^2$, $C = \sum_{i=1}^n x_i^3$. Prove that

$$(n+1)A^2B + (n-2)B^2 \geq A^4 + (2n-2)AC$$

19. (imc-2014-1-2) Consider the following sequence

$$(a_n)_{n=1}^{\infty} = (1, 1, 2, 1, 2, 3, 1, 2, 3, 4, 1, 2, 3, 4, 5, 1, \dots)$$

Find all pairs (α, β) of positive real numbers such that

$$\lim_{n \rightarrow \infty} \frac{\sum_{k=1}^{\infty} a_k}{n^\alpha} = \beta$$

20. (imc-2015-1-2) For a positive integer n , let $f(n)$ be the number obtained by writing n in binary and replacing every 0 with 1 and vice versa. For example, $n = 23$ is 10111 in binary, so $f(n)$ is 1000 in binary, therefore $f(23) = 8$. Prove that

$$\sum_{k=1}^n f(k) \leq \frac{n^2}{4}$$

When does equality hold?

21. (imc-2015-1-3) Let $F(0) = 0$, $F(1) = \frac{3}{2}$, and $F(n) = \frac{2}{5}F(n-1) - F(n-2)$ for $n \geq 2$. Determine whether or not

$$\sum_{n=0}^{\infty} \frac{1}{F(2^n)}$$

is a rational number.

22. (imc-2016-1-3) Let n be a positive integer. Also let a_1, a_2, \dots, a_n and b_1, b_2, \dots, b_n be real numbers such that $a_i + b_i > 0$ for $i = 1, 2, \dots, n$. Prove that

$$\sum_{i=1}^n \frac{a_i b_i - b_i^2}{a_i + b_i} \leq \frac{\sum_{i=1}^n a_i \cdot \sum_{i=1}^n b_i - (\sum_{i=1}^n b_i)^2}{\sum_{i=1}^n (a_i + b_i)}$$

23. (imc-2016-2-1) Let (x_1, x_2, \dots) be a sequence of positive real numbers satisfying

$$\sum_{n=1}^{\infty} \frac{x_n}{2n-1} = 1$$

Prove that

$$\sum_{k=1}^{\infty} \sum_{n=1}^k \frac{x_n}{k^2} \leq 2$$

9.2 Hints

1. $a_n = 1 + \sqrt{b_n}$, telescope.
2. (imc-1996-2-5) Riemann sum for $f(t) = \frac{t}{(1+t^2)^2}$, where $h = \frac{1}{\sqrt{x}}$.
3. (imc-1997-1-1) Riemann sum for $f(x) = \ln(x)$.
4. Проверить
5. (imc-1999-1-2) Дискретное преобразование Абеля. $\pi(1) + \dots + \pi(n) \geq 1 + \dots + n$.
6. (imc-1999-2-3) $(x+1)(x-1/2)^2 \geq 0$.
7. (imc-2001-1-3) $\int_0^\infty \frac{dx}{1+e^x}$.
8. (imc-2002-1-3)
$$\frac{2^n}{n} \left[\sum_{k=0}^{n-1} \frac{1}{C_{n-1}^k} \right] = \sum_{k=1}^n \frac{2^n}{n}$$
9. (imc-2003-1-1) $b_{n+1} > b_n$.
10. (imc-2003-2-6) $f(x) = \sum_{n=0}^\infty a_n x^n$. $f'(x) = ?$. $\ln(f(x)) = \int_0^x \frac{f'(t)}{f(t)} dt$
11. (imc-2008-2-3) Bine formula for Fibonacci.
12. (imc-2010-1-2) $\frac{x^{4k}}{4k}$
13. (imc-2010-1-3) $x_{n+1}^2 - 4 = x_n^2(x_n^2 - 4)$
14. (imc-2010-2-1) No, $x_n = (-1)^{n-1}\pi$. Yes, Since the function $t \rightarrow t \sin t$ is continuous, $y_{n+1} = |y_n| \sin |y_n| \rightarrow |a| \sin |a| = a$
15. (imc-2010-2-2) Apply induction on n .
16. (imc-2011-2-1) $u_n = \operatorname{arctanh} a_n$ (hyperbolic).
17. (imc-2012-2-2) $ka_k = \frac{a_{k+1}}{a_k} + (k+1)a_{k+1}$
18. (imc-2013-1-4) $p(x) = (x-x_1)(x-x_2)\cdots(x-x_n)$. The $(n-3)$ -th derivative of p has three nonnegative real roots $0 \leq u \leq v \leq w$.
19. (imc-2014-1-2) Let see first $n(n+1)/2$ addend.
20. (imc-2015-1-2) $2^{s-1} - 1 \leq n \leq 2^s - 1$.
21. (imc-2015-1-3) $F(n) = 2^n - 2^{-n}$
22. (imc-2016-1-3) $\frac{XY-Y^2}{X+Y} = Y - \frac{2Y^2}{X+Y}$
23. (imc-2016-2-1) interchang the sum, $\sum_{k=m}^\infty \frac{1}{k^2} < \frac{1}{n-\frac{1}{2}}$.

10 Linear algebra

10.1 Problems

09.07.2018

1. (imc-1994-1-1) Let A be a $n \times n$, $n \geq 2$, symmetric, invertible matrix with real positive elements. Show that $z_n \leq n^2 - 2n$, where z_n is the number of zero elements in A^{-1} .
2. (imc-1995-1-5) Let A and B be real $n \times n$ matrices. Assume that there exist $n + 1$ different real numbers t_1, t_2, \dots, t_{n+1} such that the matrices $C_i = A + t_i B$, $i = 1, 2, \dots, n + 1$, are nilpotent (i.e. $C_i^n = 0$). Show that both A and B are nilpotent.
3. (imc-1995-2-1) Let A be 3×3 real matrix such that the vectors Au and u are orthogonal for each column vector $u \in \mathbb{R}^3$. Prove that:

- $A^T = -A$;
- there exists a vector $v \in \mathbb{R}^3$ such that $Au = v \times u$ for every $u \in \mathbb{R}^3$.

4. (imc-1996-1-1) Let for $j = 0, \dots, n$, $a_j = a_0 + jd$, where a_0, d are fixed real numbers. Put

$$\begin{vmatrix} a_0 & a_1 & a_2 & \dots & a_n \\ a_1 & a_0 & a_1 & \dots & a_{n-1} \\ a_2 & a_1 & a_0 & \dots & a_{n-2} \\ \dots & \dots & \dots & \dots & \dots \\ a_n & a_{n-1} & a_{n-2} & \dots & a_0 \end{vmatrix}$$

Calculate $\det(A)$.

5. (imc-1996-1-3) The linear operator A on the vector space V is called an involution if $A^2 = E$ where E is the identity operator on V . Prove that for every involution A on V there exists a basis of V consisting of eigenvectors of A .
6. (imc-1997-1-3) Let A and B be real $n \times n$ matrices such that $A^2 + B^2 = AB$. Prove that if $BA - AB$ is an invertible matrix then n is divisible by 3.
7. (imc-1997-2-2) Let M be an invertible matrix of dimension $2n \times 2n$, represented in block form as

$$M = \begin{vmatrix} A & B \\ C & D \end{vmatrix}$$

$$M^{-1} = \begin{vmatrix} E & F \\ G & H \end{vmatrix}$$

Show that $\det M \cdot \det H = \det A$.

8. (imc-1999-1-1)

- Show that for any $m \in \mathbb{N}$ there exists a real $m \times m$ matrix A such that $A^3 = A + I$, where I is the $m \times m$ identity matrix.
- Show that $\det A > 0$ for every real $m \times m$ matrix satisfying $A^3 = A + I$.

9. (imc-2000-1-3) A and B are square complex matrices of the same size and $\text{rank}(AB - BA) = 1$. Show that $(AB - BA)^2 = 0$.
10. (imc-2002-2-1) Compute the determinant of the $n \times n$ matrix $A = a_{ij}$:

$$a_{ij} = \begin{cases} (-1)^{|i-j|}, & i \neq j \\ 2, & i = j \end{cases}$$

11. (imc-2003-1-3) Let A be an $n \times n$ real matrix such that $3A^3 = A^2 + A + I$ (I is the identity matrix). Show that the sequence A^k converges to an idempotent matrix. (A matrix B is called idempotent if $B^2 = B$.)
12. (imc-2003-2-1) Let A and B be $n \times n$ real matrices such that $AB + A + B = 0$. Prove that $AB = BA$.
13. (imc-2004-2-1) Let A be a real 4×2 matrix and B be a real 2×4 matrix such that

$$AB = \begin{bmatrix} 1 & 0 & -1 & 0 \\ 0 & 1 & 0 & -1 \\ -1 & 0 & 1 & 0 \\ 0 & -1 & 0 & 1 \end{bmatrix}$$

Find BA .

14. (imc-2005-1-1) Let A be the $n \times n$ matrix, whose (i, j) -th entry is $i + j$ for all $i, j = 1, 2, \dots, n$. What is the rank of A ?
15. (imc-2007-2-4) Let $n > 1$ be an odd positive integer and $A = (a_{ij})$ be an $n \times n$ matrix with:

$$a_{ij} = \begin{cases} 2, & i = j \\ 1, & i - j = \pm 2 \pmod{n} \\ 0, & \text{otherwise} \end{cases}$$

Find $\det(A)$.

16. (imc-2008-2-5) Let $n > 1$ be an odd positive integer and $A = (a_{ij})$ be an $n \times n$ matrix with:

$$a_{ij} = \begin{cases} 1, & i + j \text{ is prime} \\ 0, & \text{otherwise} \end{cases}$$

Prove that $\det A = k^2$ for some integer k .

17. (imc-2008-2-2) Let A , B and C be real square matrices of the same size, and suppose that A is invertible. Prove that if $(A - B)C = BA^{-1}$, then $C(A - B) = A^{-1}B$.
18. (imc-2009-2-3) Let $A, B \in M_n(C)$ be two $n \times n$ matrices such that $A^2B + BA^2 = 2ABA$. Prove that there exists a positive integer k such that $(AB - BA)^k = 0$.
19. (imc-2010-1-5) Suppose that a, b, c are real numbers in the interval $[-1, 1]$ such that

$$1 + 2abc \geq a^2 + b^2 + c^2$$

Prove that

$$1 + 2(abc)^n \geq a^{2n} + b^{2n} + c^{2n}$$

for all positive integers n .

20. (imc-2013-1-1) Let A and B be real symmetric matrices with all eigenvalues strictly greater than 1. Let λ be a real eigenvalue of matrix AB . Prove that $|\lambda| > 1$.
21. (imc-2014-1-1) Determine all pairs (a, b) of real numbers for which there exists a unique symmetric 2×2 matrix M with real entries satisfying $\text{trace}(M) = a$ and $\det(M) = b$.
22. (imc-2014-2-2) Let $A = (a_{ij})$ be a symmetric $n \times n$ matrix with real entries, and let $\lambda_1, \lambda_2, \dots, \lambda_n$ denote its eigenvalues. Show that

$$\sum_{1 \leq i < j \leq n} a_{ii}a_{jj} \geq \sum_{1 \leq i < j \leq n} \lambda_i \lambda_j$$

and determine all matrices for which equality holds.

23. (imc-2015-1-1) For any integer $n \geq 2$ and two $n \times n$ matrices with real entries A, B that satisfy the equation

$$A^{-1} + B^{-1} = (A + B)^{-1}$$

prove that $\det(A) = \det(B)$. Does the same conclusion follow for matrices with complex entries?

24. (imc-2017-1-1) Determine all complex numbers λ for which there exist a positive integer n and a real $n \times n$ matrix A such that $A^2 = A^T$ and λ is an eigenvalue of A .

10.2 Hints

1. (imc-1994-1-1) we have at least two non-zero elements in every column of A^{-1} .
2. (imc-1995-1-5) n^2 polynom for each matrix element
3. (imc-1995-2-1) Set $v_1 = -a_{23}$, $v_2 = a_{13}$, $v_3 = -a_{12}$.
4. (imc-1996-1-1) $(-1)^n a_0 + a_n 2^{n-1} d^n$
5. (imc-1996-1-3) The minimal polynomial of A is a divisor of $x^2 - 1$. Eigenvalue equal to ± 1 and its multiplicity is less or equal to 1. Matrix is diagonalizable.
6. (imc-1997-1-3) $S = A + e^{i\frac{2\pi}{3}} B$. Then $S\bar{S} = e^{i\frac{2\pi}{3}} (BA - AB)$.
7. (imc-1997-2-2) The definition. If A or H invertable. $\det(XY) = \det(YX)$.
8. (imc-1999-1-1) The minimal polynomial of A is a divisor of $p(x) = x^3 - x - 1$ has a positive real root and two conjugated complex roots. Matrix is diagonalizable. $\det A$ equal to product of eigenvalues.
9. (imc-2000-1-3) Jordan form.
10. (imc-2002-2-1) $\det(A) = n + 1$
11. (imc-2003-1-3) The minimal polynomial of A is a divisor of $3x^3 - x^2 - x - 1$. This polynomial has three different roots. This implies that A is diagonalizable. The eigenvalues of the matrices A and D are all roots of polynomial. One of the three roots is 1, the remaining two roots have smaller absolute value than 1. Hence, the diagonal elements of D^k , which are the k -th powers of the eigenvalues, tend to either 0 or 1 and the limit $M = \lim D^k$ is idempotent. Then $\lim A^k = C^{-1} M C$ is idempotent as well.
Минимальный многочлен делит характеристический многочлен матрицы. Любой аннулирующий многочлен делится на минимальный[1]. Минимальный многочлен единственен. Множество корней минимального многочлена совпадает с множеством корней характеристического многочлена матрицы.
12. (imc-2003-2-1) $(A + I)(B + I) = I = (B + I)(A + I)$
13. (imc-2004-2-1) $A = (A_1 A_2)^T$, $B = (B_1 B_2)$.
14. (imc-2005-1-1) $rank = 2$
15. (imc-2007-2-4) $A = B^2$.
16. (imc-2008-2-5) $|C00C| = k^2$ Now let X' be the matrix obtained from A by replacing the first row by $(100 \dots 0$, and let Y be the matrix obtained from A by replacing the entry a_{11} by 0. By multilinearity of the determinant, $\det(A) = \det(X') + \det(Y)$. Note that X' can be written as $|10vX|$ for some $(n-1) \times (n-1)$ -matrix X and some column vector v . Then $\det(A) = \det(X) + \det(Y)$. Now consider two cases. If n is odd, then X is of type (C), and Y is of type (B). Therefore, $|\det(A)| = |\det(X)|$ is a perfect square. If n is even, then X is of type (B), and Y is of type (C); hence $|\det(A)| = |\det(Y)|$ is a perfect square. The set of primes can be replaced by any subset of $\{2\} \cup \{3, 5, 7, 9, 11, \dots\}$.
17. (imc-2008-2-2) $XY = I = YX$.
18. (imc-2009-2-3) $\delta^2 B = 0$. $\delta^k(B^k) = k!(\delta B)^k$. Cayley-Hamilton Theorem.
Set $X = AB - BA$. The matrix X commutes with A . $X^{m+1} = AX^m B - BX^m A$. Take trace.
19. (imc-2010-1-5) $\begin{vmatrix} 1 & a & b \\ a & 1 & c \\ b & c & 1 \end{vmatrix}$

20. (imc-2013-1-1) The transforms given by A and B strictly increase the length of every nonzero vector, this can be seen easily in a basis where the matrix is diagonal with entries greater than 1 in the diagonal. Hence their product AB also strictly increases the length of any nonzero vector, and therefore its real eigenvalues are all greater than 1 or less than -1 .
21. (imc-2014-1-1) Note that $\text{trace}(M) = a$ and $\det(M) = b$ if and only if the two eigenvalues of M are solutions of $x^2 - ax + b = 0$.
22. (imc-2014-2-2) Eigenvalues of a real symmetric matrix are real, hence the inequality makes sense. Similarly, for Hermitian matrices diagonal entries as well as eigenvalues have to be real.

$$\sum_i a_{ii}^2 \geq \sum_{i,j} a_{ij}^2 = \sum_i \lambda_i^2$$

23. (imc-2015-1-1) $X = AB^{-1}$, $X^3 = I$. In case of complex matrices the statement is false.
24. (imc-2017-1-1) $A^4 - A = 0$. The matrix represents a rotation by $2\pi/3$.

11 Number Theory

11.1 Problems

10.07.2018

1. Докажите, что существует бесконечно много чисел, не представимых в виде суммы 3 точных квадратов.
2. Доказать, что существует бесконечно много чисел, не представимых в виде суммы трёх кубов.
3. Докажите, что уравнение не имеет решений в целых числах: $x^3 + 21y^2 + 5 = 0$.
4. Докажите, что $x^2 + xy - y^2 = 3$ не имеет целых решений.
5. Существует ли такое натуральное n , что $n^2 + n + 1$ делится на 2015?
6. Решите в натуральных числах $x! - 1 = y^2$.
7. Докажите, что если $a^2 + b^2 + ab$ делится на 11, то a и b делятся на 11.
8. Найдите все пятерки простых чисел, для которых число $q_1^4 + q_2^4 + q_3^4 + q_4^4 + q_5^4$ равно произведению двух последовательных четных натуральных чисел.
9. Докажите, что уравнение $x^3 + y^3 = 4(x^2y + xy^2 + 1)$ не имеет решений в целых числах.
10. Найдите все такие пары простых чисел p и q , что $p^3 - q^5 = (p + q)^2$.
11. Дана последовательность $a_n = 1 + 2^n + 3^n + 4^n + 5^n + 6^n + 7^n$. Существуют ли a_n , делящиеся на 2016?
12. Докажите, что число $60^{2017} + 2017^{60}$ – составное.
13. Пусть p – простое число, $p > 2$. Докажите, что любой простой делитель числа $2^p - 1$ имеет вид $2kp + 1$.
14. Докажите, что ни при каком целом k число $k^2 + k + 1$ не делится на 101.
15. Пусть $(a, b) = 1$.
 - Пусть $a|bc$. Докажите, что $a|c$.
 - Докажите, что $(a, bc) = (a, c)$.
16. Докажите, что дробь несократима при всех натуральных n :
а) $\frac{21n+4}{14n+3}$; б) $\frac{7n+16}{3n+7}$;
17. Докажите, что если a делится на b , то $(a, b) = b$. При каких целых n следующая дробь будет целой:
а) $\frac{n^2+1}{n+1}$; б) $\frac{n^3+n+1}{n^2-n+1}$; в) $\frac{n^4+1}{n^2+n+1}$.
18. Докажите, что при $m \neq n$ выполняется равенство: $(a^m - 1, a^n - 1) = a^{(m,n)} - 1$ ($a > 1$).
19. Пусть $f_k = 2^{2^k} + 1$ – числа Ферма.
 - Составьте рекуррентную формулу f_k и f_{k-1} .
 - Докажите, что $(f_n, f_m) = 1$, где $n \neq m$.
20. Пусть F_n – числа Фибоначчи ($F_0 = 0, F_1 = 1, F_n = F_{n-1} + F_{n-2}$).
 - Докажите, что $F_n = F_k \cdot F_{n-k+1} + F_{k-1} \cdot F_{n-k}$.

- Докажите, что $(F_n, F_m) = F_{(n,m)}$.
21. Пусть $b \neq d$ и $\frac{a}{b}, \frac{c}{d}$ – несократимые дроби. Докажите, что $\frac{a}{b} + \frac{c}{d}$ – не целое число.
 22. Докажите, что $\frac{[a,b,c]^2}{[a,b]*[b,c]*[a,c]} = \frac{(a,b,c)^2}{(a,b)*(b,c)*(c,a)}$.
 23. При каком наибольшем n число $n^3 + 100$ делится на $n + 10$.
 24. Найдите все такие натуральные n , что $80^n - 1$ делится на $8^n - 1$ без остатка.
 25. Найти $(5^m + 7^m, 5^n + 7^n)$.
 26. Докажите, что число $(36a + b)(36b + a)$ не может быть точной степенью двойки.
 27. Известно, что $\frac{1}{x} - \frac{1}{y} = \frac{1}{z}$. Пусть $h = (x, y, z)$. Докажите, что $hxyz$ – полный квадрат.

12 Train

12.1 Problems

15-20.07.2018

1. Доказать, что если матрицы A и B действительные положительно определенные симметричные порядка n , то

$$\det(A + B) \geq \det(A) + \det(B)$$

2. Пусть функция $f(x)$ интегрируема по Риману на отрезке $[0; 1]$, причем $|f(x)| \leq 1$. Доказать неравенство

$$\int_0^1 \sqrt{1 - f(x)^2} dx \leq \sqrt{1 - \left(\int_0^1 f(x) dx \right)^2}$$

3. Функция $f(x)$ дважды непрерывно дифференцируема на отрезке $[a, b]$ и имеет на $[a, b]$ не менее трех различных нулей. Доказать, что существует точка $\xi \in [a, b]$ такая, что

$$f(\xi) + f''(\xi) = 2f'(\xi)$$

4. Let $a_1 = a_2 = 97$ and

$$a_{n+1} = a_n a_{n-1} + \sqrt{(a_n^2 - 1)(a_{n-1}^2 - 1)}$$

Prove that

- $2 + 2a_n$ is a perfect square;
- $2 + \sqrt{2 + 2a_n}$ is a perfect square.

5. Evaluate

$$\lim_{n \rightarrow \infty} \sum_{k=0}^n \frac{k(n-k)! + (k+1)}{(k+1)!(n-k)!}$$

6. Determine whether or not there exist integers $a, b, c > 2018$ satisfying the equation

$$a^3 + 2b^3 + 4c^3 = 6abc + 1.$$

7. Let $f : [0, 1] \rightarrow \mathbb{R}$ be a differentiable function with continuous derivative such that

$$\int_0^1 (f'(x))^2 dx = 1$$

Prove that $|f(1) - f(0)| < 1$.

8. Let $A \in \mathbb{R}^{m,n}$, $B \in \mathbb{R}^{n,p}$, $C \in \mathbb{R}^{p,q}$, $D \in \mathbb{R}^{q,m}$ be such that

$$A^T = BCD, B^T = CDA, C^T = DAB, D^T = ABC$$

Prove that $(ABCD)^2 = ABCD$.

9. Let $P(x)$ be a polynomial with real coefficients of degree at least 2. Prove that if there is a real number a such that

$$P(a)P''(a) > (P'(a))^2,$$

then P has at least two nonreal zeros.

10. Let $A, B \in \mathbb{R}^{2018, 2018}$ such that $AB = BA$, $A^{2018} = I$, $B^{2018} = I$ and $\text{tr}(AB) = 2018$. Prove that $\text{tr}(A) = \text{tr}(B)$.

11. Let $f : \mathbb{C} \rightarrow \mathbb{C}$ be a function such that $f(z)f(iz) = z^2$ for all $z \in \mathbb{C}$. Prove that $f(-z) = -f(z)$ for all $z \in \mathbb{C}$.
12. a) k is even positive integer and A is a real symmetric $n \times n$ matrix such that

$$(tr(A^k))^{k+1} = (tr(A^{k+1}))^k.$$

Prove that $A^n = tr(A)A^{n-1}$.

- b) Does the assertion from a) also hold for odd positive integers k .

12.2 Hints

1. (Садовничий, стр. 229, №23) Докажем при $A = I$. Докажем при $A = C^2$, где C — симметрическая положительно определенная.
2. (Садовничий, стр. 120, №23) Неравенство Коши-Шварца.
3. (Садовничий, стр. 101, №5) Теорема Ролля для функции $g(x) = e^{-x}f(x)$.
4. (Titu 360, p.80, №55) The expression $a_1^2 - 1 = x^2$ and $2a_1 + 2 = y^2$ ask for substitution: $a_1 = \frac{1}{2}(b^2 + \frac{1}{b^2})$, hence $b + \frac{1}{b} = 14$. Setting $b = c^2$, and $c = 2 + \sqrt{3}$.
We will prove that $a_n = \frac{1}{2}(c^{4F_n} + c^{-4F_n})$.
5. (Titu 360, p.195, №13) Is equal to $1 - \frac{1}{(n+1)!} + \frac{2^n}{n!}$.
6. (AOPS, USA December TST for IMO 2012, Problem 3)

$$x^3 + y^3 + z^3 - 3xyz = \det \begin{pmatrix} x & y & z \\ z & x & y \\ y & z & x \end{pmatrix}$$

Let $w^3 = 2$. Then

$$x^3 + 2y^3 + 4z^3 - 6xyz = \det \begin{pmatrix} x & y & z \\ z & x & y \\ y & z & x \end{pmatrix} \cdot \det \begin{pmatrix} 1 & w & w^2 \\ w^2 & 1 & w \\ w & w^2 & 1 \end{pmatrix}$$

7. (Titu 360, p.218, №56)

$$\int_0^1 (f'(x) - 1)^2 dx \geq 0$$

8. (AOPS) $X = ABCD$.

$$A = D^T C^T B^T = ABCDABCD A = X^2 A \Leftrightarrow X = X^3$$

Минимальный аннулирующий многочлен матрицы X делит многочлен $t^3 - t$, то есть имеет корни среди $\{-1, 0, 1\}$, причем все они имеют степень не выше 1. Значит, матрица диагонализирема.

$$AA^T = ABCD = X \Rightarrow (Xe, e) = (A^T e, A^T e) \geq 0 \Rightarrow$$

$$X \geq 0 \Rightarrow \lambda(X) \geq 0$$

На диагонали стоят только 0 или 1, то есть $X^2 = X$.

9. (Titu 360, p.45, №57) Differentiate the equation:

$$\frac{P'(x)}{P(x)} = \sum_{i=1}^n \frac{1}{x - x_i}$$

10. (AOPS) $\lambda_1 + \lambda_2 + \dots + \lambda_n = \text{tr}(AB)$.

11. (Titu 360, p.36, №39) Substitute z with zi .

12. (AOPS)

$$(\lambda_1^k + \lambda_2^k + \dots + \lambda_n^k)^{\frac{1}{k}} = (\lambda_1^{k+1} + \lambda_2^{k+1} + \dots + \lambda_n^{k+1})^{\frac{1}{k+1}}$$