

a
Adjusted standardized mean difference vs reference (SMD)
Hidden
0.3
0.0
−0.3
SMD
−0.5
0.0
0.5
0.1
0.3
0.5
Domain
Painful neuropathy
Painless neuropathy
Hidden neuropathy
Hand symptoms only
Back pain only
26
19
22
16
10
0
10
20
n, CI ≠ 0
HbA1c
BMI
Waist
Glucose
Weight
ALP
CRP
RDW
WBC
Triglycerides
GGT
ALT
Phosphorus
Neutrophils %
LDH
Uric acid
BUN
eGFR
AST
Potassium
MPV
Systolic BP
Calcium
Platelets
Creatinine
Eosinophils %
Bilirubin
Height
Chloride
Monocytes %
MCV
Cholesterol
Lymphocytes %
Diastolic BP
Sodium
Iron
Bicarbonate
Haemoglobin
Total protein
Albumin
Glucose & lipids
Body size
Blood pressure
Liver & protein
Kidney & electrolytes
Blood count & inflammation
b
Hidden vs painless neuropathy
SMD, hidden neuropathy
SMD, painless neuropathy
0.50
0.25
0.00
−0.25
−0.50
HbA1c
BMI
Waist
Glucose
Weight
ALP
CRP
RDW
WBC
Triglycerides
GGT
ALT
Phosphorus
Neutrophils %
LDH
Diastolic BP
Sodium
Iron
Bicarbonate
Haemoglobin
Total protein
Albumin
r = 0.81, P = 3 × 10−10
slope 1.15 (95% CI 0.87–1.42)
cosine similarity 0.83
14 biomarkers shifted in both
c
Hidden vs back-pain control
SMD, hidden neuropathy
SMD, back pain only
HbA1c
BMI
Waist
Glucose
Weight
ALP
CRP
RDW
WBC
Triglycerides
GGT
ALT
Phosphorus
Neutrophils %
LDH
Diastolic BP
Sodium
Iron
Bicarbonate
Haemoglobin
Total protein
Albumin
r = 0.66, P = 3 × 10−6
slope 1.91 (95% CI 1.20–2.62)
cosine similarity 0.71
8 biomarkers shifted in both
d
Profile cosine similarity
Painless
Painful
Hands only
Back pain
0.83
0.85
0.79
0.74
0.57
0.61
0.71
0.69
0.57
0.77
Hidden
Painless
Painful
Hands only
e
Similarity with each domain left out
n
All biomarkers
Without glucose & lipids
Without body size
Without blood pressure
Without liver & protein
Without kidney & electrolytes
Without blood count & inflammation
40
35
36
38
32
31
28
0.4
0.6
0.8
1.0
Cosine similarity with the hidden-group profile
Painless neuropathy
Painful neuropathy
Hand symptoms only
Back pain only