
a
Neuropathic foot symptoms
Foot pain or tingling
D10/D1 2.3× · OR/SD 1.64, P = 4 × 10⁻¹²
Foot numbness
D10/D1 2.8× · OR/SD 1.61, P = 9 × 10⁻⁹
Prevalence (%)
PU-score decile (within age band × sex)
b
Negative-control symptoms
Back pain
D10/D1 1.1× · OR/SD 1.19, P = 2 × 10⁻⁴
Hand-only symptoms
D10/D1 1.2× · OR/SD 1.02, P = 0.84
Prevalence (%)
PU-score decile (within age band × sex)
c
All-cause mortality (crude)
Died
D10/D1 1.9× · HR/SD 1.45, P = 2 × 10⁻²⁴
Died by 2019 (%)
PU-score decile (within age band × sex)
d
Balance across deciles
Mean age (y)
Female (%)
n per decile 642–652
PU-score decile
e
Per-SD associations (adjusted for age, sex, race)
HR (95% CI)
P
Mortality
1.45 (1.35–1.55)
<.001
Mortality, + ABI/CVD
1.36 (1.26–1.47)
<.001
OR: all / held-out
P: all / held-out
Foot pain or tingling
1.64 / 1.30
<.001 / .005
Foot numbness
1.61 / 1.54
<.001 / .001
Back pain
1.19 / 1.09
<.001 / .23
Hand-only symptoms
1.02 / 1.21
.84 / .15
Odds or hazard ratio per SD of PU score
All cycles 1999–2004
Held-out 2003–04
Cox HR