

a
Transfer-model contrast
22/23 intervals exclude 0
SMN – back pain only (logit)
0.4
0.2
0
P value
<.001 <.001 .009 <.001 <.001 <.001 <.001 <.001 <.001 <.001 <.001 <.001 <.001 <.001 <.001 <.001 <.001 <.001 .002 .15 <.001 .03
Learner
Logistic regression (L2)
Elastic net
Scoring
Single-fold scoring
Features
No glycemic features
No body-size features
Adjustment
Adjusted for BMI and waist
Sample
Adults without diabetes
Held-out cycle 2003–04
Case definition
Corroborated symptoms
Exclusions
Excl. sciatica
Excl. low B12
Excl. PAD / claudication
Excl. arthritis, cancer, alcohol, CKD
Excl. high urate / gout drugs
Excl. neurotoxic drugs
Excl. hepatitis C
Excl. thyroid disease
b
All-cause mortality
17/20 intervals exclude 1
Hazard ratio, SMN vs reference
5
3
2
1.5
1
0.6
P value
.86 .56 .25 .04 .01 .04 .007 .007 .02 .006 .01 .01 .004 .03 .002 .009 .03 <.001 <.001 <.001
Sample
Adults without diabetes
Cycles 1999–2002 only
Cycle 2003–04 only
Excl. deaths < 2 y
Case definition
Corroborated symptoms
Exclusions
Excl. sciatica
Excl. low B12
Excl. arthritis, cancer, alcohol, CKD
Excl. high urate / gout drugs
Excl. neurotoxic drugs
Excl. hepatitis C
Excl. thyroid disease
Circulation
Excl. PAD / claudication
+ ABI, claudication, CVD
Normal circulation only
SMN, 95% CI excludes null
95% CI includes null
Main analysis
Back pain only
Circulation-related analysis