

a
Transfer-model contrast
24 of 25 sensitivity analyses exclude 0
SMN − back pain only (logit)
0.4
0.2
0
P value
<.001
<.001
.009
<.001
<.001
<.001
<.001
<.001
<.001
<.001
<.001
<.001
<.001
<.001
<.001
<.001
<.001
<.001
.002
.01
<.001
.03
Learner
Logistic regression (L2)
Elastic net
Survey-weighted, PSU-grouped training
Scoring
Single-fold scoring
Features
No glycemic features
No body-size features
No demographic features
Adjustment
Adjusted for BMI and waist
Sample
Adults without diabetes
Held-out cycle 2003–04
Case definition
Corroborated symptoms
Isolated foot symptoms
Exclusions
Excl. sciatica
Excl. low B12
Excl. PAD / claudication
Excl. arthritis, cancer, alcohol, CKD
Excl. high urate / gout drugs
Excl. neurotoxic drugs
Excl. hepatitis C
Excl. thyroid disease
b
All-cause mortality
19 of 22 sensitivity analyses exclude 1
Hazard ratio, SMN vs reference
5
3
2
1.5
1
0.6
P value
.90
.56
.25
.04
.01
.04
.01
.007
.007
.02
.006
.004
.01
.01
.002
.004
.03
.002
.009
.03
.03
<.001
<.001
<.001
Sample
Adults without diabetes
Cycles 1999–2002 only
Cycle 2003–04 only
Excl. deaths < 2 y
Case definition
Corroborated symptoms
Isolated foot symptoms
Missing covariates
Without income-to-poverty ratio
Multiple imputation
Exclusions
Excl. sciatica
Excl. low B12
Excl. arthritis, cancer, alcohol, CKD
Excl. high urate / gout drugs
Excl. neurotoxic drugs
Excl. hepatitis C
Excl. thyroid disease
Circulation
Excl. PAD / claudication
+ ABI, claudication, CVD
Normal circulation only (refitted)
SMN, 95% CI excludes null
95% CI includes null
Main analysis
Back pain only
Circulation-related analysis