

a
NHANES 1999–2004
n = 31,126
Excluded, n = 21,156
aged <40 years
Aged ≥40 years
n = 9,970
Excluded, n = 825
not examined
Examined
n = 9,145
Excluded, n = 1,243
no valid monofilament
Valid monofilament
n = 7,902
Excluded, n = 8
symptom question unanswered
Included
n = 7,894
Monofilament normal (all sites felt), n = 6,474
≥1 insensate site, n = 1,420
b
SMN
Study group
n 644
Prev., % 8.2 (7.3–9.2)
Died by 2019 267/642
front
back
Feet Hands Low back Feet only
100% 57% 67% 15.8%
Glycemic status, %
64.6 13.7 21.7
Back pain only
Negative control
n 1,716
Prev., % 23.8 (22.7–24.9)
Died by 2019 615/1,713
front
back
Feet Hands Low back Feet only
0% 0% 100% –
Glycemic status, %
73.3 16.8 10.0
Hand symptoms only
Negative control
n 645
Prev., % 9.0 (8.0–10.1)
Died by 2019 233/645
front
back
Feet Hands Low back Feet only
0% 100% 57% –
Glycemic status, %
73.9 14.7 11.4
Reference
Training, negative
n 3,469
Prev., % 45.3 (43.7–46.9)
Died by 2019 1,204/3,463
front
back
Feet Hands Low back Feet only
0% 0% 0% –
Glycemic status, %
75.2 15.9 8.9
Insensate with symptoms
Comparison group
n 267
Prev., % 2.7 (2.3–3.1)
Died by 2019 182/266
front
back
Feet Hands Low back Feet only
100% 49% 64% 19.9%
Glycemic status, %
42.8 14.6 42.6
Insensate, no symptoms
Training, positive
n 1,153
Prev., % 11.0 (10.0–12.0)
Died by 2019 722/1,153
front
back
Feet Hands Low back Feet only
0% 12% 39% –
Glycemic status, %
58.6 20.3 21.1
HbA1c <5.7%
HbA1c 5.7–6.4%
Diabetes