





Trial-to-criterion vs. session — window=3 trials, criterion=0.7
All subjects — Short→Long, Long trials
slope=-0.011
p=0.739, R²=0.01, n=13
All subjects — Short→Long, Short trials
slope=+0.013
p=0.298, R²=0.11, n=12
All subjects — Long→Short, Long trials
slope=-0.023
p=0.523, R²=0.04, n=13
All subjects — Long→Short, Short trials
slope=+0.007
p=0.902, R²=0.00, n=13
MC06 — Short→Long, Long trials
slope=-0.011
p=0.739, R²=0.01, n=13
MC06 — Short→Long, Short trials
slope=+0.013
p=0.298, R²=0.11, n=12
MC06 — Long→Short, Long trials
slope=-0.023
p=0.523, R²=0.04, n=13
MC06 — Long→Short, Short trials
slope=+0.007
p=0.902, R²=0.00, n=13
Occurrences to criterion (P≥0.7, window=3)
Session (chronological order)
MC06


Trial-to-criterion vs. session — window=2 trials, criterion=0.7 (robustness)

























Exponential adaptation time constant τ vs. session (fits with R²≥0.15 only)
























Early / Middle / Late training-session tertiles — trial-to-criterion (window=3, criterion=0.7)
Bracket: Early vs Late (paired Wilcoxon across subjects / Mann-Whitney within subject)

























Early / Middle / Late training-session tertiles — exponential τ (fits with $R^2 \geq 0.15$ only)
Bracket: Early vs Late (paired Wilcoxon across subjects / Mann-Whitney within subject)






















Trial-to-criterion robustness: rolling window (2 vs 3 trials) × criterion threshold, pooled across sessions
window=2 trials — Short→Long, Long trials
window=2 trials — Short→Long, Short trials
window=2 trials — Long→Short, Long trials
window=2 trials — Long→Short, Short trials
window=3 trials — Short→Long, Long trials
window=3 trials — Short→Long, Short trials
window=3 trials — Long→Short, Long trials
window=3 trials — Long→Short, Short trials
Mean occurrences to criterion
Criterion threshold
0.60
0.70
0.75
0.80
100% reached
78% reached
89% reached
93% reached
85% reached
64% reached
44% reached

Occurrence-indexed accuracy curves: Early vs Late training-session tertile

