




Trial-to-criterion vs. session — window=3 trials, criterion=0.7

























Trial-to-criterion vs. session — window=2 trials, criterion=0.7 (robustness)

























Exponential adaptation time constant τ vs. session (fits with $R^2 \geq 0.15$ only)

























Early / Middle / Late training-session tertiles — trial-to-criterion (window=3, criterion=0.7)
Bracket: Early vs Late (paired Wilcoxon across subjects / Mann-Whitney within subject)
























Early / Middle / Late training-session tertiles — exponential τ (fits with $R^2 \geq 0.15$ only)
Bracket: Early vs Late (paired Wilcoxon across subjects / Mann-Whitney within subject)






















Trial-to-criterion robustness: rolling window (2 vs 3 trials) × criterion threshold, pooled across sessions
window=2 trials — Short→Long, Long trials
window=2 trials — Short→Long, Short trials
window=2 trials — Long→Short, Long trials
window=2 trials — Long→Short, Short trials
window=3 trials — Short→Long, Long trials
window=3 trials — Short→Long, Short trials
window=3 trials — Long→Short, Long trials
window=3 trials — Long→Short, Short trials
Mean occurrences to criterion
Criterion threshold
0.60
0.70
0.75
0.80
95% reached
70% reached
85% reached
65% reached
55% reached
60% reached

Occurrence-indexed accuracy curves: Early vs Late training-session tertile

