Common/rare choice bias relative to non-naive neutral trials — Exact session ISI extremes

All subjects — pooled — all blocks
All subjects — pooled — short block
All subjects — pooled — long block
All subjects — grand average — all blocks
All subjects — grand average — short block
All subjects — grand average — long block
MC06 — pooled — all blocks
MC06 — pooled — short block
MC06 — pooled — long block
MC06 — grand average — all blocks
MC06 — grand average — short block
MC06 — grand average — long block
Choice bias: ΔP(Choose Long-Associated Response)
(block − matched neutral)
mean ± SEM
ns
n/a
Common easy
Common difficult
Rare easy
Rare difficult
1 sess 43 trials
1 sess 30 trials
1 sess 10 trials
1 sess 7 trials
1 sess 23 trials
1 sess 13 trials
1 sess 6 trials
1 sess 1 trials
1 sess 20 trials
1 sess 17 trials
1 sess 4 trials
1 sess 6 trials

Block versus matched non-naive neutral P(Choose Long-Associated Response) — Exact session ISI extremes
Pooled: Fisher exact; grand average: paired Wilcoxon

All subjects — pooled — all blocks
All subjects — pooled — short block
All subjects — pooled — long block
All subjects — grand average — all blocks
All subjects — grand average — short block
All subjects — grand average — long block
MC06 — pooled — all blocks
MC06 — pooled — short block
MC06 — pooled — long block
MC06 — grand average — all blocks
MC06 — grand average — short block
MC06 — grand average — long block
Matched neutral
Block
P(Choose Long-Associated Response)
Common easy
Common difficult
Rare easy
Rare difficult
1 sess
ns
n/a
0.0
0.2
0.4
0.6
0.8
1.0
1.2
0.5
0.6
0.7
0.8
0.9

Bias distributions — Exact session ISI extremes: median + IQR and two-sided tests versus zero
All visual summaries use median + IQR; Shapiro p < 0.01 selects Wilcoxon, otherwise one-sample t-test

All subjects — pooled — all blocks
All subjects — pooled — short block
All subjects — pooled — long block
All subjects — grand average — all blocks
All subjects — grand average — short block
All subjects — grand average — long block
MC06 — pooled — all blocks
MC06 — pooled — short block
MC06 — pooled — long block
MC06 — grand average — all blocks
MC06 — grand average — short block
MC06 — grand average — long block
Bias distribution
ΔP(Long-associated response)
0.1
0.0
−0.1
−0.2
−0.3
−0.4
−0.5
0.06
0.04
0.02
0.00
−0.02
−0.04
−0.06
ns
ns
n/a
ns
Common easy
Common difficult
Rare easy
Rare difficult
n=2
one-sample t

Common/rare choice bias relative to non-naive neutral trials — Four psychometric bins (outer bins define easy/hard)

All subjects — pooled — all blocks
All subjects — pooled — short block
All subjects — pooled — long block
All subjects — grand average — all blocks
All subjects — grand average — short block
All subjects — grand average — long block
MC06 — pooled — all blocks
MC06 — pooled — short block
MC06 — pooled — long block
MC06 — grand average — all blocks
MC06 — grand average — short block
MC06 — grand average — long block
Choice bias: ΔP(Choose Long-Associated Response) (block − matched neutral)
0.8
0.6
0.4
0.3
0.2
0.1
0.0
−0.1
−0.2
−0.3
−0.4
−0.6
−0.8
ns
n/a
○ mean ± SEM
Common easy
Common difficult
Rare easy
Rare difficult
1 sess 51 trials
1 sess 49 trials
1 sess 12 trials
1 sess 10 trials
1 sess 28 trials
1 sess 22 trials
1 sess 7 trials
1 sess 2 trials
1 sess 23 trials
1 sess 27 trials
1 sess 5 trials
1 sess 8 trials

Block versus matched non-naive neutral P(Choose Long-Associated Response) — Four psychometric bins (outer bins define easy/hard)
Pooled: Fisher exact; grand average: paired Wilcoxon

All subjects — pooled — all blocks
All subjects — pooled — short block
All subjects — pooled — long block
All subjects — grand average — all blocks
All subjects — grand average — short block
All subjects — grand average — long block
MC06 — pooled — all blocks
MC06 — pooled — short block
MC06 — pooled — long block
MC06 — grand average — all blocks
MC06 — grand average — short block
MC06 — grand average — long block
Matched neutral
Block
P(Choose Long-Associated Response)
Common easy
Common difficult
Rare easy
Rare difficult
1 sess
ns
n/a

Bias distributions — Four psychometric bins (outer bins define easy/hard): median + IQR and two-sided tests versus zero
All visual summaries use median + IQR; Shapiro $p < 0.01$ selects Wilcoxon, otherwise one-sample t-test

All subjects — pooled — all blocks
Bias distribution
ΔP(Long-associated response)
0.4
0.3
0.2
0.1
0.0
−0.1
−0.2
−0.3
ns
ns
ns
ns
Common
easy
Common
difficult
Rare
easy
Rare
difficult
n=2
one-sample t
n=2
one-sample t
n=2
one-sample t
n=2
one-sample t
All subjects — pooled — short block
All subjects — pooled — long block
All subjects — grand average — all blocks
All subjects — grand average — short block
All subjects — grand average — long block
0.06
0.04
0.02
0.00
−0.02
−0.04
−0.06
MC06 — pooled — all blocks
MC06 — pooled — short block
MC06 — pooled — long block
MC06 — grand average — all blocks
MC06 — grand average — short block
MC06 — grand average — long block

Bin-resolved psychometric choice bias: block minus matched non-naive neutral

All subjects — pooled — all blocks
All subjects — pooled — short block
All subjects — pooled — long block
All subjects — grand average — all blocks
All subjects — grand average — short block
All subjects — grand average — long block
MC06 — pooled — all blocks
MC06 — pooled — short block
MC06 — pooled — long block
MC06 — grand average — all blocks
MC06 — grand average — short block
MC06 — grand average — long block
Common — Short ISI
Common — Long ISI
Rare — Short ISI
Rare — Long ISI
Choice bias: ΔP(Long-associated response) (block − matched neutral)
ISI (s)
0.5
1.0
1.5
2.0
0.6
0.4
0.2
0.0
-0.2
-0.4
-0.6
-0.8
0.3
0.1
-0.1
-0.3