





Trial-to-criterion vs. session — window=3 trials, criterion=0.7
All subjects — Short→Long, Long trials
slope=+0.054
p=0.179, R²=0.33, n=7
MC06
All subjects — Short→Long, Short trials
slope=-0.030
p=0.338, R²=0.18, n=7
MC06
All subjects — Long→Short, Long trials
slope=+0.000
p=1, R²=0.00, n=7
MC06
All subjects — Long→Short, Short trials
slope=-0.030
p=0.123, R²=0.41, n=7
MC06
MC06 — Short→Long, Long trials
slope=+0.054
p=0.179, R²=0.33, n=7
MC06 — Short→Long, Short trials
slope=-0.030
p=0.338, R²=0.18, n=7
MC06 — Long→Short, Long trials
slope=+0.000
p=1, R²=0.00, n=7
MC06 — Long→Short, Short trials
slope=-0.030
p=0.123, R²=0.41, n=7
Occurrences to criterion (P≥0.7, window=3)
Session (chronological order)


Trial-to-criterion vs. session — window=2 trials, criterion=0.7 (robustness)

























Exponential adaptation time constant τ vs. session (fits with $R^2 \geq 0.15$ only)

























Early / Middle / Late training-session tertiles — trial-to-criterion (window=3, criterion=0.7)
Bracket: Early vs Late (paired Wilcoxon across subjects / Mann-Whitney within subject)

























Early / Middle / Late training-session tertiles — exponential τ (fits with $R^2 \geq 0.15$ only)
Bracket: Early vs Late (paired Wilcoxon across subjects / Mann-Whitney within subject)






















Trial-to-criterion robustness: rolling window (2 vs 3 trials) × criterion threshold, pooled across sessions
window=2 trials — Short→Long, Long trials
window=2 trials — Short→Long, Short trials
window=2 trials — Long→Short, Long trials
window=2 trials — Long→Short, Short trials
window=3 trials — Short→Long, Long trials
window=3 trials — Short→Long, Short trials
window=3 trials — Long→Short, Long trials
window=3 trials — Long→Short, Short trials
Mean occurrences to criterion
Criterion threshold
0.60
0.70
0.75
0.80
100% reached
71% reached
86% reached
74% reached
44% reached
70% reached
51% reached

Occurrence-indexed accuracy curves: Early vs Late training-session tertile

