Block-specific PSE and lapse rates with shared slope: pooled and equal-session psychometrics by subject

$P(\text{Long} \mid x, b) = \lambda_b + (1-\lambda_b-\gamma_b)\,\sigma[\beta(x-\alpha_b)]$; $\alpha_N=\alpha_0,\ \alpha_L=\alpha_0+\delta_L,\ \alpha_S=\alpha_0+\delta_S$

All subjects

MC06

Block-specific PSE, slope, and lapse rates: pooled and equal-session psychometrics by subject

$$P(\text{Long} \mid x, b) = \lambda_b + (1-\lambda_b-\gamma_b)\,\sigma[\beta_b(x-\alpha_b)];\quad \alpha_L = \alpha_N - \delta\alpha_L,\ \alpha_S = \alpha_N - \delta\alpha_S,\ \beta_L = \beta_N - \delta\beta_L,\ \beta_S = \beta_N - \delta\beta_S$$

All subjects

Pooled trials
Neutral (n=347)
Short (n=961)
Long (n=960)
Neutral: λ=0.127, γ=0.096
Short: λ=0.025, γ=0.041
Long: λ=0.000, γ=0.000
P(Choose Long-Associated Response)
ISI (s)
Sessions weighted equally
Neutral (n=347)
Short (n=961)
Long (n=960)
Neutral: λ=0.138, γ=0.068
Short: λ=0.027, γ=0.000
Long: λ=0.038, γ=0.000
Pooled trials — fitted PSE
Sessions weighted equally — fitted PSE
PSE
Pooled trials — fitted slope
Sessions weighted equally — fitted slope
Slope
Pooled trials — fitted lower lapse
Sessions weighted equally — fitted lower lapse
Lower lapse
Pooled trials — fitted upper lapse
Sessions weighted equally — fitted upper lapse
Upper lapse
Neutral
Short
Long

MC06

Pooled trials
Neutral (n=347)
Short (n=961)
Long (n=960)
Neutral: λ=0.127, γ=0.096
Short: λ=0.025, γ=0.041
Long: λ=0.000, γ=0.000
P(Choose Long-Associated Response)
ISI (s)
Sessions weighted equally
Neutral (n=347)
Short (n=961)
Long (n=960)
Neutral: λ=0.138, γ=0.068
Short: λ=0.027, γ=0.000
Long: λ=0.038, γ=0.000
Pooled trials — fitted PSE
Sessions weighted equally — fitted PSE
PSE
Pooled trials — fitted slope
Sessions weighted equally — fitted slope
Slope
Pooled trials — fitted lower lapse
Sessions weighted equally — fitted lower lapse
Lower lapse
Pooled trials — fitted upper lapse
Sessions weighted equally — fitted upper lapse
Upper lapse
Neutral
Short
Long