

## Supplement 4. Studies included in the review.

1. Alosco ML, Spitznagel MB, Miller L, Raz N, Cohen R, Sweet LH, et al. Depression is associated with reduced physical activity in persons with heart failure. *Health Psychol.* 2012 Nov;31(6):754–62.
2. Chien H-C, Chen H-M, Garet M, Wang R-H. Predictors of physical activity in patients with heart failure: a questionnaire study. *J Cardiovasc Nurs.* 2014 Jul;29(4):324–31.
3. Corvera-Tindel T, Doering LV, Gomez T, Dracup K. Predictors of noncompliance to exercise training in heart failure. *J Cardiovasc Nurs.* 2004 Aug;19(4):269–77; quiz 278.
4. Dontje ML, van der Wal MHL, Stolk RP, Brügemann J, Jaarsma T, Wijtvliet PEPJ, et al. Daily physical activity in stable heart failure patients. *J Cardiovasc Nurs.* 2014 Jun;29(3):218–26.
5. Evangelista LS, Berg J, Dracup K. Relationship between psychosocial variables and compliance in patients with heart failure. *Heart Lung.* 2001 Aug;30(4):294–301.
6. Evangelista L, Doering LV, Dracup K, Westlake C, Hamilton M, Fonarow GC. Compliance behaviors of elderly patients with advanced heart failure. *J Cardiovasc Nurs.* 2003 Aug;18(3):197–206; quiz 207.
7. Gallagher R, Luttik M-L, Jaarsma T. Social support and self-care in heart failure. *J Cardiovasc Nurs.* 2011 Dec;26(6):439–45.
8. Klompstra L, Jaarsma T, Strömberg A. Self-efficacy Mediates the Relationship Between Motivation and Physical Activity in Patients With Heart Failure. *J Cardiovasc Nurs.* 2018;33(3):211–6.
9. Lee H, Boo S, Yu J, Suh S-R, Chun KJ, Kim JH. Physical Functioning, Physical Activity, Exercise Self-Efficacy, and Quality of Life Among Individuals With Chronic Heart Failure in Korea: A Cross-Sectional Descriptive Study. *J Nurs Res.* 2016 Jun 27.
10. Moreno-Suarez I, Liew S, Dembo LG, Larbalestier R, Maiorana A. Physical Activity Is Higher in Patients with LVADs Compared to Chronic Heart Failure. *Med Sci Sports Exerc.* 2019 Jul 24.
11. Oka RK, Gortner SR, Stotts NA, Haskell WL. Predictors of physical activity in patients with chronic heart failure secondary to either ischemic or idiopathic dilated cardiomyopathy. *Am J Cardiol.* 1996 Jan 15;77(2):159–63.
12. Pihl E, Fridlund B, Mårtensson J. Patients' experiences of physical limitations in daily life activities when suffering from chronic heart failure; a phenomenographic analysis. *Scand J Caring Sci.* 2011 Mar;25(1):3–11.
13. Pozehl BJ, McGuire R, Duncan K, Hertzog M, Deka P, Norman J, et al.

Accelerometer-Measured Daily Activity Levels and Related Factors in Patients With Heart Failure. *J Cardiovasc Nurs*. 2018;33(4):329–35.

14. Snipelisky D, Kelly J, Levine JA, Koepp GA, Anstrom KJ, McNulty SE, et al. Accelerometer-Measured Daily Activity in Heart Failure With Preserved Ejection Fraction: Clinical Correlates and Association With Standard Heart Failure Severity Indices. *Circ Heart Fail*. 2017 Jun;10(6):e003878.
15. Tierney S, Elwers H, Sange C, Mamas M, Rutter MK, Gibson M, et al. What influences physical activity in people with heart failure?: a qualitative study. *Int J Nurs Stud*. 2011 Oct;48(10):1234–43.
16. Tierney S, Mamas M, Skelton D, Woods S, Rutter MK, Gibson M, et al. What can we learn from patients with heart failure about exercise adherence? A systematic review of qualitative papers. *Health Psychol*. 2011 Jul;30(4):401–10.
17. van der Wal MHL, Jaarsma T, Moser DK, Veeger NJGM, van Gilst WH, van Veldhuisen DJ. Compliance in heart failure patients: the importance of knowledge and beliefs. *Eur Heart J*. 2006 Feb;27(4):434–40.  
  
van der Wal MHL, van Veldhuisen DJ, Veeger NJGM, Rutten FH, Jaarsma T. Compliance with non-pharmacological recommendations and outcome in heart failure patients. *Eur Heart J*. 2010 Jun;31(12):1486–93.
18. Werhahn SM, Dathe H, Rottmann T, Franke T, Vahdat D, Hasenfuß G, et al. Designing meaningful outcome parameters using mobile technology: a new mobile application for telemonitoring of patients with heart failure. *ESC Heart Fail*. 2019 Mar 13;6(3):516–25.
19. Witham MD, Argo IS, Johnston DW, Struthers AD, McMurdo MET. Predictors of exercise capacity and everyday activity in older heart failure patients. *Eur J Heart Fail*. 2006 Mar;8(2):203–7.