

The goal is to analyse the Differentiation Matrix properties

Given a large set of nodes  $\{\vec{x}_i\}_{i=1}^n$  located in  $\mathbb{R}^d$ .

1. Infinitely smooth Radial function  $\phi(\cdot)$

A local interpolant takes the form  $S(\vec{x}) = \sum_{j=1}^m \lambda_j \phi((\vec{x} - \vec{x}_j))$   
where  $\vec{x}_j$  is the central node and  $\{\vec{x}_{i,j}\}_{j=1}^m$  are its  $m$  neighbours

We can impose the interpolation condition at each neighbouring node.

Let's denote  $\{\phi(\vec{x}_{i,j})\}_{j=1}^m$  by  $\vec{F}$

Define the collocation matrix  $A_i = [\phi((\vec{x}_{i,j} - \vec{x}_i))]_{j=1}^m$ , s.t.  $A_i \vec{F} = \vec{f}$  (1)

We wish to find the weights  $\{w_j\}_{j=1}^m$  s.t.  $\sum_{j=1}^m w_j \phi(\vec{x}_{i,j}) = L_i g(\vec{x})|_{\vec{x}=\vec{x}_i}$

New denote  $\{w_j\}_{j=1}^m$  by  $\vec{w}$ , and  $(L_i g(\vec{x})|_{\vec{x}=\vec{x}_i})_{j=1}^m$  by  $\vec{g}_i$ , so, in matrix form, we seek  $\vec{w}$  s.t.  $\vec{F}^\top \vec{w} = \vec{g}_i$  (2)

We can apply the differential operator  $L$  to the local interpolant.

$L_i S(\vec{x}) = \sum_{j=1}^m \lambda_j L_i \phi((\vec{x} - \vec{x}_{i,j}))$  and evaluate it at the central point  $\vec{x} = \vec{x}_i$ .

$$L_i g(\vec{x}_i) \approx L_i S(\vec{x}_i) = \sum_{j=1}^m \lambda_j L_i \phi((\vec{x}_i - \vec{x}_{i,j})) \Big|_{\vec{x}=\vec{x}_i} = \left( L_i \phi((\vec{x}_i - \vec{x}_{i,1})) \Big|_{\vec{x}=\vec{x}_i}, L_i \phi((\vec{x}_i - \vec{x}_{i,2})) \Big|_{\vec{x}=\vec{x}_i}, \dots, L_i \phi((\vec{x}_i - \vec{x}_{i,m})) \Big|_{\vec{x}=\vec{x}_i} \right)^\top \vec{\lambda}$$

$$= B_i \vec{\lambda}$$

From (1) (Since  $A$  is invertible)  $\vec{g}_i = B_i \cdot A^{-1} \vec{F}$

$$\text{from (2), } \vec{w} = (B_i \cdot A^{-1})^\top = (A^{-1})^\top B_i^\top = (A^\top)^{-1} B_i^\top$$

$$\text{So } A^\top \vec{w} = B_i^\top$$

Note :  $A$  is a local collocation matrix, associated to its cluster center  $\vec{x}_i$ .


$B_i$  is  $L$  applied to the local interpolant and evaluated at  $\vec{x}_i$ .

### Global Representation :

Consider a graph of each center with its  $m$  neighbours

A sort of adjacency matrix can be built.

For instance :  $E = \begin{pmatrix} 0 & 1 & 1 & \\ 1 & 0 & 1 & 1 \\ 1 & 1 & 0 & \\ & & & \ddots \end{pmatrix}$




Properties: (1)  $E$  is  $N \times N$  ( $N$  is the total number of nodes)

(2)  $\sum_{j=1}^N E_{ij} = m, \forall i$  ( $m$  is fixed : the number of neighbours per cluster)

(3)  $E$  is not necessarily symmetric

(4)  $E_{ii} = 1, \forall i$



Let  $A$  be the global collocation matrix, with all the nodes. Similarly, let  $B$  be the global differentiated interpolant evaluated at each node.

Let  $A_i$  be the local collocation matrix associated to the cluster of which the central node is  $\bar{x}_i$ . Similarly,  $B_i$  is the local interpolant differentiated and evaluated at the central node (recall it is a row vector).

The neighbors of  $\bar{x}_i$  are  $\{\bar{x}_{i_1}, \bar{x}_{i_2}, \dots, \bar{x}_{i_m}\}$

The indices of the neighbors are the positions of the non-zero entries of  $E_{i,:}$ .

Let  $P_i$  be the  $n \times N$  reduction matrix, so

$$\boxed{A_i = P_i A P_i^T}$$

$$\text{For instance, } E_{i,:} = [0 \ 1 \ 0 \ 0 \ 1 \ 1] \quad , \quad P_i = \begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix} \quad N=6, m=3 \quad 3 \times 6$$

$$A_i \cdot P_i A P_i^T = \begin{bmatrix} 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} t_{i_1} & t_{i_2} & t_{i_3} & \cdots & t_{i_6} \\ t_{i_1} & t_{i_2} & & & \\ \vdots & & & & \\ t_{i_1} & t_{i_2} & t_{i_3} & t_{i_4} & t_{i_5} & t_{i_6} \\ t_{i_1} & t_{i_2} & t_{i_3} & t_{i_4} & t_{i_5} & t_{i_6} \\ t_{i_1} & t_{i_2} & t_{i_3} & t_{i_4} & t_{i_5} & t_{i_6} \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} = \begin{bmatrix} t_{i_1} & t_{i_2} & t_{i_3} \\ t_{i_1} & t_{i_2} & t_{i_3} \end{bmatrix}$$

*central node index*

$$\text{Similarly, } B_i = P_i (C_i) B P_i^T = [0 \ 0 \ 0 \ 0 \ 0 \ 1 \ 0] \begin{bmatrix} L & t_{i_1} & t_{i_2} & \cdots & t_{i_6} \\ t_{i_1} & L & t_{i_2} & \cdots & t_{i_6} \\ \vdots & \vdots & \vdots & & \vdots \\ t_{i_1} & t_{i_2} & t_{i_3} & L & t_{i_5} \\ t_{i_1} & t_{i_2} & t_{i_3} & t_{i_4} & L \\ t_{i_1} & t_{i_2} & t_{i_3} & t_{i_4} & t_{i_5} & L \end{bmatrix} = [L \ t_{i_1} \ t_{i_2} \ t_{i_3} \ t_{i_4} \ t_{i_5}]$$

so the local cluster differentiation weights satisfy

$$A_i^T \bar{w}_i = B_i^T \quad \text{Here, } A_i \text{ is symmetric, so } A_i \bar{w}_i = B_i^T$$

$$P_i A P_i^T \bar{w}_i = P_i B^T P_i (C_i)^T$$

$$\bar{w}_i = (P_i A P_i^T)^{-1} (P_i B^T P_i (C_i)^T)$$

Building the Differentiation Matrix:

$\bar{w}_i$  contains the entries of the  $i^{\text{th}}$  row, along the non-zero columns of  $E_{i,:}$ .

$$\text{So } D_i = P_i^T (\cdot, c) \bar{w}_i^T P_i$$

$N \times 1 \quad m \times m \quad m \times N$

$$\begin{aligned} \text{So } D &= \sum_{i=1}^N P_i^T (\cdot, c) \bar{w}_i^T P_i \\ &\quad \text{i.e. row of } B \\ &= \sum_{i=1}^N \underbrace{P_i^T (\cdot, c) (P_i (\cdot, c) B P_i^T)}_{\text{zero matrix}} \underbrace{(P_i A^T P_i^T)^{-1} P_i}_{\bar{w}_i^T} \\ &\quad \text{except 1 at } (i,i) \end{aligned}$$

Note: For the symmetric version of  $D$ :

$$W_i = (P_i A P_i^T)^{-1} (P_i B P_i^T)$$

$m \times m \quad m \times n \quad n \times m$

$$D = \sum_{i=1}^N P_i^T (P_i A P_i^T)^{-1} (P_i B P_i^T) P_i$$