

Name: \_\_\_\_\_

Show an appropriate amount of work. You can (and probably should) check your work with a computer.

**1. Compute the magnitude and argument for**

**1.1.**  $z = \left(\frac{i}{1+i}\right)^3$

$\text{abs}(z) =$

$\text{arg}(z) =$

**1.2.**  $z = \left(\frac{1+i}{\sqrt{1+3i}}\right)^3$

$\text{abs}(z) =$

$\text{arg}(z) =$

**1.3.** The roots of  $z^4 = -64$

$\text{abs}(z) =$

$\text{arg}(z) =$

**1.4.** The roots of  $z^2 + 2iz + 5 = 0$

$\text{abs}(z) =$

$\text{arg}(z) =$

**1.5.**  $z = i^i$

$\text{abs}(z) =$

$\text{arg}(z) =$

2. Compute the real and imaginary parts for

2.1.  $z = \left(\frac{i}{1+i}\right)^3$

$$z = \boxed{\phantom{000}} + i \boxed{\phantom{000}}$$

2.2.  $z = \left(\frac{1+i}{\sqrt{1+3i}}\right)^3$

$$z = \boxed{\phantom{000}} + i \boxed{\phantom{000}}$$

2.3. The roots of  $z^3 = -64$

$$z_1 = \boxed{\phantom{000}} + i \boxed{\phantom{000}}$$

$$z_2 = \boxed{\phantom{000}} + i \boxed{\phantom{000}}$$

$$z_3 = \boxed{\phantom{000}} + i \boxed{\phantom{000}}$$

2.4. The roots of  $z^2 + 2iz + 5 = 0$

$$z_1 = \boxed{\phantom{000}} + i \boxed{\phantom{000}}$$

$$z_2 = \boxed{\phantom{000}} + i \boxed{\phantom{000}}$$

2.5.  $z = i^i$

$$z = \boxed{\phantom{000}} + i \boxed{\phantom{000}}$$


### 3. Sketch Points

**3.1.** Sketch the set of points  $C_1$  that satisfy  $|z - 2| < 1$ . Make sure you label the set.

**3.2.** Sketch the set of points  $C_2$  that satisfy  $\text{im}(z - 2) = 1$ . Make sure you label the set.

**3.3.** Sketch the set of points  $C_3$  that satisfy  $\arg(z - 2) = \pi/4$ . Make sure you label the set.

Out[ ]=



**4.** For  $f(z) = u(x, y) + i v(x, y)$  and write down the CR equations

and

**4.1.** For  $\cos(z)$  complete the following

$u(x, y) =$   and  $v(x, y) =$

$\partial_x u =$   and  $\partial_x v =$

$\partial_y u =$   and  $\partial_y v =$

Analytic Yes or No

**4.2.** For  $i x^2 - 2 x y - i y^2$  complete the following

$u(x, y) =$   and  $v(x, y) =$

$\partial_x u =$   and  $\partial_x v =$

$\partial_y u =$   and  $\partial_y v =$

Analytic Yes or No  .

If yes  $f(z) =$

**4.3.** For  $i x^2 - 2 x y - y^2$  complete the following

$u(x, y) =$   and  $v(x, y) =$

$\partial_x u =$   and  $\partial_x v =$

$\partial_y u =$   and  $\partial_y v =$

Analytic Yes or No  .

If yes  $f(z) =$

5. Is  $e^{x-iy}$  analytic? Yes or No . Show your work below.

6. Label poles with  $P_1, P_2, \dots$  Label zeros with  $Z_1, Z_2, \dots$


$$\text{Res}(f, P_1) = \boxed{\phantom{000000}}$$

$$\text{Res}(f, P_2) = \boxed{\phantom{000000}}$$

$$\text{In}[*]:= \mathbf{f[z\_]} := 3 \frac{\mathbf{z - 1 + i}}{(1 + z^2)}$$

**ComplexPlot3D[ f[z],  
{z, 2}, PlotLegends → Automatic]**

Out[\*]=



**7. Complete the following. The TS**

$$\arctan(z) = \sum_{k=0}^{\infty} a_k z^k \text{ where } a_k =$$

converges for . Justify your

convergence statement using a Calculus II theorem. Give a simpler complex analytic justification below.

**8. Complete the following. The TS**

$$\log(2+z) = \sum_{k=0}^{\infty} a_k z^k \text{ where } a_k =$$

converges for . Justify your


convergence statement using a Calculus II theorem. Give a simpler complex analytic justification below.



9. Label and explain the singularity and color discontinuity visible in the plot below using appropriate language. The TS  $f(z) = \sum_{k=0}^{\infty} a_k(z - z_0)^k$  with  $z_0 = -2 + 2i$  has  $a_k =$   and converges for  $|z - z_0| <$  . Explain what the TS converges to around the color discontinuity.

```
In[46]:= f[z_] := Log[1 + z]
ComplexPlot3D[f[z],
  {z, 2}, PlotLegends -> Automatic]
```

Out[47]=



- 10.** Label and explain the singularity and color discontinuity visible in the plots below using appropriate language. Explain why you might expect  $f_1$  to match  $f_2$ . Explain why they do not match using appropriate language.


In[138]:=

$$f1[z_] := \sqrt{1 - z^2}$$

$$f2[z_] := \pm \sqrt{z^2 - 1}$$

```
{ComplexPlot3D[ f1[z], {z, 2}, PlotLabel → "f1"],  
  ComplexPlot3D[ f2[z], {z, 2}, PlotLabel → "f2"] }
```

Out[140]=



**11.** For the closed counter clockwise contour  $\Gamma$  shown below

$$p_1(t) = \boxed{\phantom{0}} \text{ for } 0 \leq t < 1$$


$$p_2(t) = \boxed{\phantom{0}} \text{ for } 0 \leq t < 1$$

$$\int_{\Gamma} f(z) dz = \int_0^1 \boxed{\phantom{0}} dt + \int_0^1 \boxed{\phantom{0}} dt.$$

For  $f(z) = z^2$  evaluate  $\int_{\Gamma} f(z) dz = \boxed{\phantom{0}}$  and

$\int_{\Gamma_1} f(z) dz = \boxed{\phantom{0}}$ . Show appropriate work below.

Out[93]=



**12.** For the closed counter clockwise contour  $\Gamma$  shown below

$$p_1(t) = \boxed{\phantom{0}} \text{ for } 0 \leq t < 1$$

$$p_2(t) = \boxed{\phantom{0}} \text{ for } 0 \leq t < 1$$

$$p_3(t) = \boxed{\phantom{0}} \text{ for } 0 \leq t < 1 \quad 0 \leq t < 1$$


$$\int_{\Gamma} f(z) dz = \int_0^1 \boxed{\phantom{0}} dt +$$

$$\int_0^1 \boxed{\phantom{0}} dt + \int_0^1 \boxed{\phantom{0}} dt$$

For  $f(z) = \frac{i}{z-(1+2i)}$  evaluate  $\int_{\Gamma} f(z) dz = \boxed{\phantom{0}}$  and

$\int_{\Gamma_3} f(z) dz = \boxed{\phantom{0}}$ . Show appropriate work below.

Out[117]=



**13.** Compute  $\int_{\Gamma} \frac{e^z}{(z^2+4)(z-1)} dz = \boxed{\phantom{000000}}$  for the counter clockwise contour  $\Gamma$ . Show appropriate work.

`p1[t_] := (E2 $\pi$ i t + 1.2)2`

`ParametricPlot[ReIm[p1[t]], {t, 0, 1}, PlotLegends → {"Γ"}]`

Out[124]=

