

Complex Integration

4410

February 16, 2026

Cauchy's Integral Formula

Theorem (Cauchy Integral Formula).

Let f be analytic on and inside a simple closed positively oriented contour γ . Let z_0 be a point inside γ . Then

$$f(z_0) = \frac{1}{2\pi i} \int_{\gamma} \frac{f(z)}{z - z_0} dz.$$

Why this is astonishing:

- The value of f at z_0 is completely determined by its values on the boundary.
- Analytic functions are incredibly rigid.
- Integration produces evaluation.

Holomorphic functions are controlled by their boundary behavior.

Cauchy Formula for Derivatives

Theorem (Higher Derivative Formula).

If f is analytic on and inside γ and z_0 is inside γ , then for $n \geq 1$:

$$f^{(n)}(z_0) = \frac{n!}{2\pi i} \int_{\gamma} \frac{f(z)}{(z - z_0)^{n+1}} dz.$$

Consequences:

- Analytic $\Rightarrow$ infinitely differentiable.
- Derivatives are given by integral formulas.
- Leads to power series expansions:

$$f(z) = \sum_{n=0}^{\infty} \frac{f^{(n)}(z_0)}{n!} (z - z_0)^n.$$

Differentiation becomes a byproduct of integration.

Riemann Integral: Limit Definition

Let $f : [a, b] \rightarrow \mathbb{R}$ be bounded. Take a partition

$$P = \{a = x_0 < x_1 < \cdots < x_n = b\}$$

and choose sample points $\xi_i \in [x_{i-1}, x_i]$. Then the **Riemann sum** is

$$S(f, P) = \sum_{i=1}^n f(\xi_i) \Delta x_i, \quad \Delta x_i = x_i - x_{i-1}.$$

If the limit exists the Riemann integral is

$$\int_a^b f(x) dx := \lim_{\|P\| \rightarrow 0} S(f, P).$$

Alternate version with $m_i = \inf_{[x_{i-1}, x_i]} f$ and $M_i = \sup_{[x_{i-1}, x_i]} f$. If

$$\sup_P L(f, P) = \inf_P U(f, P).$$

then f is Riemann integrable where

$$L(f, P) = \sum m_i \Delta x_i, \quad U(f, P) = \sum M_i \Delta x_i,$$

Properties of the Riemann Integral

- Linearity: $\int_a^b (cf + g) = c \int_a^b f + \int_a^b g$.
- Monotonicity: $f \leq g \Rightarrow \int_a^b f \leq \int_a^b g$.
- Additivity: $\int_a^b f = \int_a^c f + \int_c^b f$.
- Continuous f are Riemann integrable.

Fundamental Theorem of Calculus (Calc II)

Let f be continuous on $[a, b]$ and define

$$F(x) = \int_a^x f(t) dt.$$

- Then $F'(x) = f(x)$ for all $x \in [a, b]$.
- For any antiderivative F of f ,

$$\int_a^b f(x) dx = F(b) - F(a).$$

This theorem connects differentiation and integration and is the basis for evaluating definite integrals.

Green's Theorem

$$\oint_C (P \, dx + Q \, dy) = \iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dA$$

for positively oriented simple closed curve C and D its interior.

Interpretation: circulation around C = double integral of curl over D .

Example (Green's Theorem)

$\oint_C (x^2 dx + y^2 dy)$ around unit circle:

$$\iint_D \left(\frac{\partial}{\partial x} y^2 - \frac{\partial}{\partial y} x^2 \right) dA = 0$$

Divergence Theorem (Gauss)

$$\iint_S \mathbf{F} \cdot \mathbf{n} \, dS = \iiint_V \nabla \cdot \mathbf{F} \, dV$$

for vector field $\mathbf{F}$ on solid V with boundary S .

Stokes' Theorem

$$\oint_{\partial S} \mathbf{F} \cdot d\mathbf{r} = \iint_S (\nabla \times \mathbf{F}) \cdot d\mathbf{S}.$$

Relates circulation along a boundary to curl over surface.

Complex Integrals (Section 6.1)

Let $f : [a, b] \rightarrow \mathbb{C}$, $f(t) = u(t) + iv(t)$. Define the integral by

$$\int_a^b f(t) dt := \int_a^b u(t) dt + i \int_a^b v(t) dt.$$

Riemann sum definition:

$$\int_a^b f(t) dt = \lim_{\|P\| \rightarrow 0} \sum_{i=1}^n f(\xi_i) \Delta t_i.$$

Chopping up explanation: Partition $[a, b]$ into small intervals. On each interval $[t_{i-1}, t_i]$, approximate the integral by $f(\xi_i) \Delta t_i$. Sum over all intervals. Take the limit as the maximum interval length $\|P\| \rightarrow 0$.

Contours and Contour Integrals (Section 6.2)

A contour $\gamma : [a, b] \rightarrow \mathbb{C}$, $\gamma(t) = x(t) + iy(t)$, defines

$$\int_{\gamma} f(z) dz := \int_a^b f(\gamma(t)) \gamma'(t) dt.$$

Riemann sum version:

$$\int_{\gamma} f(z) dz = \lim_{\|P\| \rightarrow 0} \sum_{i=1}^n f(\gamma(t_i^*)) (\gamma(t_i) - \gamma(t_{i-1})).$$

Chopping up explanation:

- Partition $[a, b]$ into tiny intervals.
- Over each small interval, approximate the curve by a line segment: $\gamma(t_i) - \gamma(t_{i-1})$.
- Multiply by $f(\gamma(t_i^*))$ at a sample point.
- Sum over all intervals; refine partition to get the integral.

Why parameterization works: $\gamma'(t)\Delta t \approx \gamma(t_i) - \gamma(t_{i-1})$ for small Δt .

Example: Contour Integral

Unit circle: $\gamma(t) = e^{it}$, $t \in [0, 2\pi]$:

$$\int_{\gamma} z^n dz = \int_0^{2\pi} e^{int} i e^{it} dt = i \int_0^{2\pi} e^{i(n+1)t} dt$$

which vanishes when $n \neq -1$.

Cauchy–Goursat Theorem (Section 6.3)

If f is analytic inside and on a simple closed contour γ ,

$$\int_{\gamma} f(z) dz = 0.$$

Allows deformation of contours for analytic functions.

Where Does Cauchy's Formula Come From?

Let f be analytic inside and on a simple closed contour γ , and let z_0 lie inside γ .

Consider the function

$$g(z) = \frac{f(z) - f(z_0)}{z - z_0}.$$

Key observations:

- $g(z)$ is analytic for $z \neq z_0$.
- Because f is differentiable at z_0 ,

$$\lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0} = f'(z_0),$$

so the singularity is removable.

- Therefore g is analytic everywhere inside γ .

By Cauchy–Goursat,

$$\int_{\gamma} g(z) dz = 0.$$

Summary

- FTC links derivatives and integrals.
- Green's, Divergence, Stokes' theorems: local derivatives $\rightarrow$ global integrals.
- Riemann sum and sup/inf definitions give rigor.
- Complex contour integrals generalize line integrals; chopping explains the limit.
- Cauchy–Goursat: integral around closed loop vanishes for analytic functions.