

Paths of analysis*

C30

Synthia

March 3, 2022

1 Analysis parameters

Analysis type: Automatic Retrosynthesis

Rules: none selected

Filters: FGI, FGI with protections

Max. paths returned: 5

Max. iterations: 300

Commercial:

1. Max. molecular weight - 1000 g/mol
2. Max. price - 1000 \$/g

Published:

1. Max. molecular weight - 1000 g/mol
2. Popularity - 10

My Stockroom:

1. Max. molecular weight - 1000 g/mol

Reaction scoring formula: $\text{TUNNEL_COEF} * \text{FGI_COEF} * \text{STEP} * 20 + 1000000 * (\text{CONFLICT} + \text{NON_SELECTIVITY} + \text{FILTERS} + \text{PROTECT})$

Chemical scoring formula: $\text{SMALLER}^3, \text{SMALLER}^{1.5}$

Min. search width: 400

Max. reactions per product: 60

Strategies: none selected

*The results stated herein were generated using the proprietary platform owned and maintained by Grzybowski Scientific Inventions, Inc., a subsidiary of Merck KGaA, Darmstadt Germany. The results are provided on an as is basis, and shall be used solely in connection with the rights afforded in the license agreement and for no other purpose.

FGI Coeff: 0

JSON Parameters: {}

2 Paths

1 path found. *Paths are sorted by score. Reactions are sorted in appearance order for each path.*

2.1 Path 1

Score: 169.06





Figure 1: Outline of path 1

2.1.1 Amide coupling



Substrates:

1. Z-Hyp-OH - *available at Sigma-Aldrich*
2. 1-Amino-cyclopropanecarbonitrile - *Combi-Blocks*

Products:

1. N#CC1(NC(=O)C2CC(O)CN2C(=O)OCc2ccccc2)CC1

Typical conditions: DCC.DCM or EDC.DCM or SOCl₂.DCM


Protections: none

Yield: good

Reference: [10.1021/cr100048w](#) and [10.1039/B701677H](#) and [10.1039/C5RA24527C](#) and [10.3727/00000006783981206](#) and [10.1021/np060007f](#) and [10.1021/jo00012a058](#) and [10.1016/j.bmcl.2007.08.037](#) and [10.1039/C0OB00355G](#) and [10.1021/jm500031w](#) (p.3056) and [10.1016/j.tet.2011.03.046](#)

Retrosynthesis ID: 10087

2.1.2 Hydrolysis of nitriles to amides



Substrates:

1. N#CC1(NC(=O)C2CC(O)CN2C(=O)OCc2ccccc2)CC1

Products:

1. NC(=O)C1(NC(=O)C2CC(O)CN2C(=O)OCc2ccccc2)CC1

Typical conditions: H₂SO₄.DCM.rt


Protections: none

Yield: moderate

Reference: [10.1021/jo00040a061](#) and [10.1016/0223-5234\(90\)90132-M](#) and [10.1016/j.tetlet.2011.09.012](#)

Retrosynthesis ID: 11358

2.1.3 Hydrolysis of nitriles to amides



Substrates:

1. 4-Fluoro-3-methoxybenzonitrile - *available at Sigma-Aldrich*

Products:

1. 4-fluoro-3-methoxybenzamide

Typical conditions: H₂SO₄.DCM.rt


Protections: none

Yield: moderate

Reference: [10.1021/jo00040a061](#) and [10.1016/0223-5234\(90\)90132-M](#) and [10.1016/j.tetlet.2011.09.012](#)

Retrosynthesis ID: 11358

2.1.4 Cleavage of benzyloxycarbamates



Substrates:

1. NC(=O)C1(NC(=O)C2CC(O)CN2C(=O)OCc2ccccc2)CC1

Products:

1. NC(=O)C1(NC(=O)C2CC(O)CN2)CC1

Typical conditions: H₂.Pd/C


Protections: none

Yield: good

Reference: [10.1021/jm070755h](#) and [10.1021/jm2016057](#) and [10.1055/s-0033-1340215](#) and [10.1016/S0040-4039\(03\)01181-X](#)

Retrosynthesis ID: 9990024

2.1.5 Sulfonylation of amides



Substrates:

1. 4-fluoro-3-methoxybenzamide
2. 2-Propanesulfonyl chloride - *available at Sigma-Aldrich*

Products:

1. COc1cc(C(=O)NS(=O)(=O)C(C)C)ccc1F

Typical conditions: Py.RSO₂Cl


Protections: none

Yield: good

Reference: [10.1021/ja9945313](#) AND [10.1016/j.ejmech.2013.04.028](#)
AND [10.1039/c5ra14001c](#) AND [10.1016/j.bmcl.2013.12.043](#) AND
[10.1016/j.tetasy.2012.08.013](#)

Retrosynthesis ID: 14787

2.1.6 Amide coupling



Substrates:

1. NC(=O)C1(NC(=O)C2CC(O)CN2)CC1
2. 2-[(tert-butoxy)carbonyl]amino-3,3-dimethylbutanoic acid - *Enamine*

Products:

1. CC(C)(C)OC(=O)NC(C(=O)N1CC(O)CC1C(=O)NC1(C(N)=O)CC1)C(C)(C)C

Typical conditions: DCC.DCM or EDC.DCM or SOCl₂.DCM


Protections: none

Yield: good

Reference: [10.1021/ol400686f](#) and [10.1021/jo00200a057](#) and [10.1021/cr100048w](#) and [10.1039/B701677H](#) and [10.1039/C5RA24527C](#) and [10.3727/000000006783981206](#) and [10.1021/np060007f](#) and [10.1021/jo00012a058](#) and [10.1016/j.bmcl.2007.08.037](#) and [10.1039/C0OB00355G](#) and [10.1021/jm500031w](#) (p.3056) and [10.1016/j.tet.2011.03.046](#)

Retrosynthesis ID: 9147

2.1.7 Nucleophilic aromatic substitution



Substrates:

1. COc1cc(C(=O)NS(=O)(=O)C(C)C)ccc1F
2. CC(C)(C)OC(=O)NC(C(=O)N1CC(O)CC1C(=O)NC1(C(N)=O)CC1)C(C)(C)C

Products:

1. COc1cc(C(=O)NS(=O)(=O)C(C)C)ccc1OC1CC(C(=O)NC2(C(N)=O)CC2)N(C(=O)C(NC(=O)OC(C

Typical conditions: NaH.THF.0-80 C or K2CO3.DMF.110 C

Protections: none

Yield: good

Reference: [10.1016/j.tetlet.2015.10.008](#) p. 6479, 6483 and
[10.1016/j.ejmech.2016.06.056](#) p. 82, 85

Retrosynthesis ID: 49475