Paths of analysis*

Synthia

March 3, 2022

1 Analysis parameters

Analysis type: Automatic Retrosynthesis

Rules: none selected

Filters: FGI, FGI with protections

Max. paths returned: 5

Max. iterations: 300

Commercial:

- 1. Max. molecular weight 1000 g/mol
- 2. Max. price 1000 \$/g

Published:

- 1. Max. molecular weight 1000 g/mol
- 2. Popularity 10

My Stockroom:

1. Max. molecular weight - 1000 g/mol

Reaction scoring formula: TUNNEL_COEF*FGI_COEF*STEP*20+1000 000*(CONFLICT+NON SELECTIVITY+FILTERS+PROTECT)

Chemical scoring formula: SMALLER^ 3,SMALLER^ 1.5

Min. search width: 400

Max. reactions per product: 60

Strategies: none selected

^{*}The results stated herein were generated using the proprietary platform owned and maintained by Grzybowski Scientific Inventions, Inc., a subsidiary of Merck KGaA, Darmstadt Germany. The results are provided on an as is basis, and shall be used solely in connection with the rights afforded in the license agreement and for no other purpose.

FGI Coeff: 0

JSON Parameters: {}

2 Paths

 $1\ \mathrm{path}$ found. Paths are sorted by score. Reactions are sorted in appearance order for each path.

2.1 Path 1

Score: 136.23




Figure 1: Outline of path 1

2.1.1 Alkylation of secondary alcohols

Substrates:

- 1. 3-chlormethyl-benzo[b]thiophen
- 2. 1-(2,4-dichloro-phenyl)-2-imidazol-1-yl-ethanol $available \ at \ Sigma-Aldrich$

Products:

 $1. \ 1\hbox{-}[2\hbox{-}(benzo[b]thiophen-3\hbox{-}ylmethoxy)-2\hbox{-}(2,4\hbox{-}dichloro\hbox{-}phenyl)\hbox{-}ethyl]-1h-imidazole$

Typical conditions: K2CO3.acetone.heat

Protections: none
Yield: moderate

Reference: 10.1016/S0022-1139(00)85021-6 and

Retrosynthesis ID: 31011106