Paths of analysis*

Synthia

March 3, 2022

1 Analysis parameters

Analysis type: Automatic Retrosynthesis

Rules: none selected

Filters: FGI, FGI with protections

Max. paths returned: 5

Max. iterations: 300

Commercial:

1. Max. molecular weight - 1000 g/mol

2. Max. price - 1000 \$/g

Published:

- 1. Max. molecular weight 1000 g/mol
- 2. Popularity 10

My Stockroom:

1. Max. molecular weight - 1000 g/mol

Reaction scoring formula: TUNNEL_COEF*FGI_COEF*STEP*20+1000 000*(CONFLICT+NON SELECTIVITY+FILTERS+PROTECT)

Chemical scoring formula: SMALLER^ 3,SMALLER^ 1.5

Min. search width: 400

Max. reactions per product: 60

Strategies: none selected

^{*}The results stated herein were generated using the proprietary platform owned and maintained by Grzybowski Scientific Inventions, Inc., a subsidiary of Merck KGaA, Darmstadt Germany. The results are provided on an as is basis, and shall be used solely in connection with the rights afforded in the license agreement and for no other purpose.

FGI Coeff: 0

JSON Parameters: {}

2 Paths

3 paths found. Paths are sorted by score. Reactions are sorted in appearance order for each path.

2.1 Path 1

Score: 626.83




Figure 1: Outline of path 1

2.1.1 Sulfonylation of amides

- $\begin{array}{ll} \hbox{1. 1-(Aminocarbonyl)-1-cyclopropane} \hbox{acad -} & \textit{available at Sigma-Aldrich} \\ \end{array}$
- 2. Cyclopropanesulfonyl chloride available at Sigma-Aldrich

Products:

Substrates:

1. O=C(O)C1(C(=O)NS(=O)(=O)C2CC2)CC1

Typical conditions: Py.RSO2Cl

Protections: none

Yield: good

10.1016/j.tetasy.2012.08.013

Retrosynthesis ID: 14787

2.1.2 Schmidt Reaction

Substrates:

1. hydrazoic acid

2. O=C(O)C1(C(=O)NS(=O)(=O)C2CC2)CC1

Products:

1. NC1(C(=O)NS(=O)(=O)C2CC2)CC1

Typical conditions: azide.H+.40C

Protections: none
Yield: moderate

Reference: 10.1039/B505080D Retrosynthesis ID: 11704

2.1.3 Amide coupling

Substrates:

- 1. NC1(C(=O)NS(=O)(=O)C2CC2)CC1
- 2. Z-Hyp-OH available at Sigma-Aldrich

Products:

 $1. \ O = C(NC1(C(=O)NS(=O)(=O)C2CC2)CC1)C1CC(O)CN1C(=O)OCc1ccccc1$

Typical conditions: DCC.DCM or EDC.DCM or SOC12.DCM

Protections: none

Yield: good

Reference: 10.1021/cr100048w and 10.1039/B701677H and 10.1039/C5RA24527C and 10.3727/0000000006783981206 and 10.1021/np060007f and 10.1021/jo00012a058 and 10.1016/j.bmcl.2007.08.037 and 10.1039/C00B00355G and 10.1021/jm500031w (p.3056) and 10.1016/j.tet.2011.03.046

Retrosynthesis ID: 10087

2.1.4 Mitsunobu reaction

Substrates:

- $1. \ O = C(NC1(C(=O)NS(=O)(=O)C2CC2)CC1)C1CC(O)CN1C(=O)OCc1ccccc1$
- 2. Methyl 3-chloro-4-hydroxybenzoate available at Sigma-Aldrich

Products:

Typical conditions: DEAD.or.DCAD.or.DIAD.PPh3

Protections: none

Yield: good

Reference: DOI: 10.1021/jo0345751 AND 10.1021/ol0618757

2.1.5Cleavage of benzyloxycarbamates

Substrates:

Products:

 $1. \ \ COC(=O)c1ccc(OC2CNC(C(=O)NC3(C(=O)NS(=O)(=O)C4CC4)CC3)C2)c(Cl)c1$

Typical conditions: H2.Pd/C

Protections: none

Yield: good

1340215 and 10.1016/S0040-4039(03)01181-X

Retrosynthesis ID: 9990024

2.1.6Amide coupling

Substrates:

- $1. \ COC(=O)c1ccc(OC2CNC(C(=O)NC3(C(=O)NS(=O)(=O)C4CC4)CC3)C2)c(Cl)c1\\$
- 2. 2-[(tert-butoxy)carbonyl]amino-3-methylpentanoic acid available at $Sigma\hbox{-}Aldrich$

Products:

Typical conditions: DCC.DCM or EDC.DCM or SOCl2.DCM

Protections: none

Yield: good

Reference: 10.1021/o1400686f and 10.1021/jo00200a057 and 10.1021/cr100048w and 10.1039/B701677H and 10.1039/C5RA24527C and 10.3727/000000006783981206 and 10.1021/np060007f and 10.1021/jo00012a058 and 10.1016/j.bmcl.2007.08.037 and 10.1039/C00B00355G and 10.1021/jm500031w (p.3056) and 10.1016/j.tet.2011.03.046

Retrosynthesis ID: 9147

2.2 Path 2

Score: 657.80




Figure 2: Outline of path 2

2.2.1 Sulfonylation of amides

Substrates:

1. 1-(Aminocarbonyl)-1-cyclopropanecarboxylic acid - available at Sigma-Aldrich

2. Cyclopropanesulfonyl chloride - available at Sigma-Aldrich

Products:

1. O=C(O)C1(C(=O)NS(=O)(=O)C2CC2)CC1

Typical conditions: Py.RSO2Cl

Protections: none

Yield: good

10.1016/j. tetasy. 2012. 08.013

Retrosynthesis ID: 14787

2.2.2 Schmidt Reaction

Substrates:

1. hydrazoic acid

 $2. \ \mathrm{O=C(O)C1(C(=O)NS(=O)(=O)C2CC2)CC1}$

Products:

1. NC1(C(=O)NS(=O)(=O)C2CC2)CC1

Typical conditions: azide.H+.40C

Protections: none
Yield: moderate

Reference: 10.1039/B505080D

2.2.3 Amide coupling

Substrates:

- 1. NC1(C(=O)NS(=O)(=O)C2CC2)CC1
- 2. Z-Hyp-OH available at Sigma-Aldrich

Products:

 $1. \ O = C(NC1(C(=O)NS(=O)(=O)C2CC2)CC1)C1CC(O)CN1C(=O)OCc1ccccc1$

Typical conditions: DCC.DCM or EDC.DCM or SOC12.DCM

Protections: none

Yield: good

Retrosynthesis ID: 10087

10.1016/j.tet.2011.03.046

2.2.4 Cleavage of benzyloxycarbamates

Substrates:

 $1. \ O = C(NC1(C(=O)NS(=O)(=O)C2CC2)CC1)C1CC(O)CN1C(=O)OCc1ccccc1$

Products:

 $1. \ O=C(NC1(C(=O)NS(=O)(=O)C2CC2)CC1)C1CC(O)CN1$

Typical conditions: H2.Pd/C

Protections: none

Yield: good

1340215 and 10.1016/S0040-4039(03)01181-X

Retrosynthesis ID: 9990024

2.2.5 Amide coupling

Substrates:

- 1. O=C(NC1(C(=O)NS(=O)(=O)C2CC2)CC1)C1CC(O)CN1
- 2. 2-[(tert-butoxy)carbonyl]amino-3-methylpentanoic acid available at Sigma-Aldrich

Products:

 $1. \ \ CCC(C)C(NC(=O)OC(C)(C)C)C(=O)N1CC(O)CC1C(=O)NC1(C(=O)NS(=O)(=O)C2CC2)CC1$

Typical conditions: DCC.DCM or EDC.DCM or SOCl2.DCM

Protections: none

Yield: good

Reference: 10.1021/ol400686f and 10.1021/jo00200a057 and 10.1021/cr100048w and 10.1039/B701677H and 10.1039/C5RA24527C and

10.3727/000000006783981206 and 10.1021/np060007f and 10.1021/jo00012a05810.1016/j.bmcl.2007.08.037and 10.1039/C0OB00355G10.1021/jm500031w (p.3056) and 10.1016/j.tet.2011.03.046

Retrosynthesis ID: 9147

2.2.6 Mitsunobu reaction

Substrates:

- $1. \ \ CCC(C)C(NC(=O)OC(C)(C)C)C(=O)N1CC(O)CC1C(=O)NC1(C(=O)NS(=O)(=O)C2CC2)CC1$
- 2. Methyl 3-chloro-4-hydroxybenzoate available at Sigma-Aldrich

Products:

Typical conditions: DEAD.or.DCAD.or.DIAD.PPh3

Protections: none

Yield: good

Reference: DOI: 10.1021/jo0345751 AND 10.1021/ol0618757

Retrosynthesis ID: 7562

Path 3

Score: 750.01




Figure 3: Outline of path 3

2.3.1 Sulfonylation of amides

Substrates:

- 1. 1-(Aminocarbonyl)-1-cyclopropanecarboxylic acid available at Sigma-Aldrich
- 2. Cyclopropanesulfonyl chloride available at Sigma-Aldrich

Products:

1. O=C(O)C1(C(=O)NS(=O)(=O)C2CC2)CC1

Typical conditions: Py.RSO2Cl

Protections: none

Yield: good

Reference: 10.1021/ja9945313 AND 10.1016/j.ejmech.2013.04.028 AND 10.1039/c5ra14001c AND 10.1016/j.bmcl.2013.12.043 AND 10.1016/j.tetasy.2012.08.013

Retrosynthesis ID: 14787

2.3.2 Schmidt Reaction

Substrates:

1. hydrazoic acid

2. O=C(O)C1(C(=O)NS(=O)(=O)C2CC2)CC1

Products:


1. NC1(C(=O)NS(=O)(=O)C2CC2)CC1

Typical conditions: azide.H+.40C

Protections: none
Yield: moderate

Reference: 10.1039/B505080D Retrosynthesis ID: 11704

2.3.3 Amide coupling



Substrates:

 $1. \ 2\hbox{-}[(\text{tert-butoxy})\text{carbonyl}]\\ a mino-3\hbox{-}methyl pentanoic acid - } \\ available \ at \\ Sigma-Aldrich$

2. cis-4-hydroxy-proline lactone - Enamine

Products:

 $1. \ \mathrm{CCC}(\mathrm{C})\mathrm{C}(\mathrm{NC}(=\mathrm{O})\mathrm{OC}(\mathrm{C})(\mathrm{C})\mathrm{C})\mathrm{C}(=\mathrm{O})\mathrm{N}1\mathrm{CC}2\mathrm{CC}1\mathrm{C}(=\mathrm{O})\mathrm{O}2$

Typical conditions: DCC.DCM or EDC.DCM or SOC12.DCM

Protections: none

Yield: good

Reference: 10.1021/ol400686f and 10.1021/jo00200a057 and 10.1021/cr100048w and 10.1039/B701677H and 10.1039/C5RA24527C and 10.3727/000000006783981206 and 10.1021/np060007f and 10.1021/jo00012a058 and 10.1016/j.bmcl.2007.08.037 and 10.1039/C0OB00355G and 10.1021/jm500031w (p.3056) and 10.1016/j.tet.2011.03.046

Retrosynthesis ID: 9147

2.3.4 Intramolecular amidation of esters

Substrates:

1. NC1(C(=O)NS(=O)(=O)C2CC2)CC1

 $2. \ \mathrm{CCC}(\mathrm{C})\mathrm{C}(\mathrm{NC}(=\mathrm{O})\mathrm{OC}(\mathrm{C})(\mathrm{C})\mathrm{C})\mathrm{C}(=\mathrm{O})\mathrm{N}1\mathrm{CC}2\mathrm{CC}1\mathrm{C}(=\mathrm{O})\mathrm{O}2$

Products:

 $1. \ \ CCC(C)C(NC(=O)OC(C)(C)C)C(=O)N1CC(O)CC1C(=O)NC1(C(=O)NS(=O)(=O)C2CC2)CC1$

Typical conditions: DABAL-(Me)3.THF.cooling

Protections: none

Yield: good

Reference: 10.1016/j.tetlet.2006.06.004 and 10.1246/cl.1987.803 and 10.1016/j.polymer.2013.01.040 and 10.1016/j.tetasy.2003.11.026 and 10.1021/ol050773y

2.3.5 Mitsunobu reaction

Substrates:

- $1. \ \ CCC(C)C(NC(=O)OC(C)(C)C)C(=O)N1CC(O)CC1C(=O)NC1(C(=O)NS(=O)(=O)C2CC2)CC1$
- 2. Methyl 3-chloro-4-hydroxybenzoate available at Sigma-Aldrich

Products:

 $\textbf{Typical conditions:} \ \ DEAD.or.DCAD.or.DIAD.PPh3$

Protections: none

Yield: good

Reference: DOI: 10.1021/jo0345751 AND 10.1021/ol0618757