

Paths of analysis*

C122

Synthia

March 3, 2022

1 Analysis parameters

Analysis type: Automatic Retrosynthesis

Rules: none selected

Filters: FGI, FGI with protections

Max. paths returned: 5

Max. iterations: 300

Commercial:

1. Max. molecular weight - 1000 g/mol
2. Max. price - 1000 \$/g

Published:

1. Max. molecular weight - 1000 g/mol
2. Popularity - 10

My Stockroom:

1. Max. molecular weight - 1000 g/mol

Reaction scoring formula: $\text{TUNNEL_COEF} * \text{FGI_COEF} * \text{STEP} * 20 + 1000000 * (\text{CONFLICT} + \text{NON_SELECTIVITY} + \text{FILTERS} + \text{PROTECT})$

Chemical scoring formula: $\text{SMALLER}^3, \text{SMALLER}^{1.5}$

Min. search width: 400

Max. reactions per product: 60

Strategies: none selected

*The results stated herein were generated using the proprietary platform owned and maintained by Grzybowski Scientific Inventions, Inc., a subsidiary of Merck KGaA, Darmstadt Germany. The results are provided on an as is basis, and shall be used solely in connection with the rights afforded in the license agreement and for no other purpose.

FGI Coeff: 0

JSON Parameters: {}

2 Paths

2 paths found. *Paths are sorted by score. Reactions are sorted in appearance order for each path.*

2.1 Path 1

Score: 138.85





Figure 1: Outline of path 1

2.1.1 Suzuki coupling of arylboronic acids with aryl bromides



Substrates:

1. 5-Bromo-7-azaindole - *available at Sigma-Aldrich*
2. (p-Fluorophenyl)boric acid - *available at Sigma-Aldrich*

Products:

1. Fc1ccc(-c2cnc3[nH]ccc3c2)cc1

Typical conditions: Pd catalyst.base.solvent


Protections: none

Yield: good

Reference: [10.1021/cr00039a007](#) and [10.1007/3418_2012_32](#) and [10.1021/cr0505268](#) and [10.1016/j.jfluchem.2016.01.018](#) and [10.1039/C3CS60197H](#) and [10.1016/j.ejmech.2018.08.092](#) and [10.1038/s41929-020-00564-z](#) (metal-free coupling)

Retrosynthesis ID: 25150

2.1.2 N-Sulfonylation



Substrates:

- 1-Propanesulfonyl chloride - [available at Sigma-Aldrich](#)
- 3-Bromo-2,4-difluoroaniline - [AstaTech](#)

Products:

- N-(3-Bromo-2,4-difluorophenyl)-1-propanesulfonamide - [available at Sigma-Aldrich](#)

Typical conditions: THF.rt


Protections: none

Yield: good

Reference: [10.1055/s-0029-1217565](#) and [10.1002/\(SICI\)1099-0690\(199806\)1998:6<945::AID-EJOC945>3.0.CO;2-3](#) and [10.1055/s-2001-14567](#) and [10.1016/j.bmc.2014.07.022](#)

Retrosynthesis ID: 14718

2.1.3 Iodination of aromatic compounds



Substrates:

1. Fc1ccc(-c2cnc3[nH]ccc3c2)cc1

Products:

1. Fc1ccc(-c2cnc3[nH]cc(I)c3c2)cc1

Typical conditions: I₂ or other iodinating agent e.g. NIS


Protections: none

Yield: good

Reference: DOI: [10.1039/C5SC00964B](https://doi.org/10.1039/C5SC00964B) and [10.1016/j.tetlet.2005.05.117](https://doi.org/10.1016/j.tetlet.2005.05.117) and [10.1007/s11178-005-0256-1](https://doi.org/10.1007/s11178-005-0256-1)

Retrosynthesis ID: 10697

2.1.4 Pd-catalyzed conversion of aryl bromides to Weinreb amides



Substrates:

1. N-(3-Bromo-2,4-difluorophenyl)-1-propanesulfonamide - *available at Sigma-Aldrich*

- Carbon monoxide - *available at Sigma-Aldrich*
- n-methoxymethylamine - *ChemImpexInternational*

Products:

- CCCC(=O)(=O)Nc1ccc(F)c(C(=O)N(C)OC)c1F

Typical conditions: Pd(OAc)₂.Xantphos.CO(1 atm).Na₂CO₃.toluene.80C


Protections: none

Yield: moderate

Reference: DOI: [10.1021/ol061902t](https://doi.org/10.1021/ol061902t)

Retrosynthesis ID: 1688

2.1.5 Synthesis of ketones from Weinreb amides



Substrates:

- CCCC(=O)(=O)Nc1ccc(F)c(C(=O)N(C)OC)c1F
- Fc1ccc(-c2cnc3[nH]cc(I)c3c2)cc1

Products:

- CCCC(=O)(=O)Nc1ccc(F)c(C(=O)c2c[nH]c3ncc(-c4ccc(F)cc4)cc23)c1F

Typical conditions: 1.RmgBr.THF 2.TFA.DCM

Protections: none

Yield: good

Reference: [10.1021/jm051185t](https://doi.org/10.1021/jm051185t) and [10.1021/ol101021v](https://doi.org/10.1021/ol101021v) (supporting info)

Retrosynthesis ID: 5060

2.2 Path 2

Score: 162.57





Figure 2: Outline of path 2

2.2.1 Suzuki coupling with aryl chlorides



Substrates:

1. N-[3-[(5-Chloro-1H-pyrrolo[2,3-b]pyridin-3-yl)carbonyl]-2,4-difluorophenyl]-1-propanesulfonamide - *AstaTech*
2. (p-Fluorophenyl)boric acid - *available at Sigma-Aldrich*

Products:

1. CCCS(=O)(=O)Nc1ccc(F)c(C(=O)c2c[nH]c3ncc(-c4ccc(F)cc4)cc23)c1F

Typical conditions: [Pd].catalyst.base.

Protections: none

Yield: good

Reference: [10.1002/anie.201108608](#) and [10.1002/anie.200801465](#) and [10.1055/s-0033-1338293](#) and [10.1039/c1cc10708a](#) and [10.1055/s-0030-1260169](#) and [10.1016/j.tet.2005.05.071](#) and [10.1038/s41929-020-00564-z](#) (metal-free coupling)

Retrosynthesis ID: 26284