

Paths of analysis*

C138

Synthia

March 3, 2022

1 Analysis parameters

Analysis type: Automatic Retrosynthesis

Rules: none selected

Filters: FGI, FGI with protections

Max. paths returned: 5

Max. iterations: 300

Commercial:

1. Max. molecular weight - 1000 g/mol
2. Max. price - 1000 \$/g

Published:

1. Max. molecular weight - 1000 g/mol
2. Popularity - 10

My Stockroom:

1. Max. molecular weight - 1000 g/mol

Reaction scoring formula: $\text{TUNNEL_COEF} * \text{FGI_COEF} * \text{STEP} * 20 + 1000000 * (\text{CONFLICT} + \text{NON_SELECTIVITY} + \text{FILTERS} + \text{PROTECT})$

Chemical scoring formula: $\text{SMALLER}^3, \text{SMALLER}^{1.5}$

Min. search width: 400

Max. reactions per product: 60

Strategies: none selected

*The results stated herein were generated using the proprietary platform owned and maintained by Grzybowski Scientific Inventions, Inc., a subsidiary of Merck KGaA, Darmstadt Germany. The results are provided on an as is basis, and shall be used solely in connection with the rights afforded in the license agreement and for no other purpose.

FGI Coeff: 0

JSON Parameters: {}

2 Paths

1 path found. *Paths are sorted by score. Reactions are sorted in appearance order for each path.*

2.1 Path 1

Score: 89.36





Figure 1: Outline of path 1

2.1.1 Alkylation of amines with alkyl chlorides



Substrates:

1. 3-methoxypyrrolidine - *Combi-Blocks*
2. 2-Chloro-5-(chloromethyl)pyridine - *available at Sigma-Aldrich*

Products:

1. COC1CCN(Cc2ccc(Cl)nc2)C1

Typical conditions: KOH. toluene. PTC. catalyst or KI. base e.g. K₂CO₃


Protections: none

Yield: moderate

Reference: [10.1016/S0040-4020\(01\)00989-9](#) and [10.1021/acs.oprd.8b00074](#)
and [10.1016/s0040-4039\(00\)74286-9](#) and [10.1080/00397911.2013.828077](#) and
[10.1016/j.bmcl.2012.08.032](#)

Retrosynthesis ID: 4784

2.1.2 Nucleophilic aromatic substitution



Substrates:

1. COC1CCN(Cc2ccc(Cl)nc2)C1

Products:

1. COC1CCN(Cc2ccc(N)nc2)C1

Typical conditions: solvent. Heating or pressure


Protections: none


Yield: good

Reference: [10.1021/jm00040a009](#) or [10.1111/bph.12233](#) or [10.1246/cl.1987.1187](#)

Retrosynthesis ID: 5003

2.1.3 Buchwald-Hartwig amination





Substrates:

1. 6-(2-Chloro-5-fluoropyrimidin-4-yl)-4-fluoro-1-isopropyl-2-methyl-1H-benzo[d]imidazole - *available at Sigma-Aldrich*
2. COC1CCN(Cc2ccc(N)nc2)C1

Products:

1. COC1CCN(Cc2ccc(Nc3nc(F)c(-c4cc(F)c5nc(C)n(C(C)C)c5c4)n3)nc2)C1

Typical conditions: PdCl₂.NaOtBu.dioxane.heat

Protections: none

Yield: good

Reference: [10.1021/acs.oprd.9b00161](https://doi.org/10.1021/acs.oprd.9b00161) and [10.1002/anie.201904795](https://doi.org/10.1002/anie.201904795) and [10.1021/acs.chemrev.6b00512](https://doi.org/10.1021/acs.chemrev.6b00512)

Retrosynthesis ID: 10319