

Paths of analysis*

C55

Synthia

March 3, 2022

1 Analysis parameters

Analysis type: Automatic Retrosynthesis

Rules: none selected

Filters: FGI, FGI with protections

Max. paths returned: 5

Max. iterations: 300

Commercial:

1. Max. molecular weight - 1000 g/mol
2. Max. price - 1000 \$/g

Published:

1. Max. molecular weight - 1000 g/mol
2. Popularity - 10

My Stockroom:

1. Max. molecular weight - 1000 g/mol

Reaction scoring formula: $\text{TUNNEL_COEF} * \text{FGI_COEF} * \text{STEP} * 20 + 1000000 * (\text{CONFLICT} + \text{NON_SELECTIVITY} + \text{FILTERS} + \text{PROTECT})$

Chemical scoring formula: $\text{SMALLER}^3, \text{SMALLER}^{1.5}$

Min. search width: 400

Max. reactions per product: 60

Strategies: none selected

*The results stated herein were generated using the proprietary platform owned and maintained by Grzybowski Scientific Inventions, Inc., a subsidiary of Merck KGaA, Darmstadt Germany. The results are provided on an as is basis, and shall be used solely in connection with the rights afforded in the license agreement and for no other purpose.

FGI Coeff: 0

JSON Parameters: {}

2 Paths

1 path found. *Paths are sorted by score. Reactions are sorted in appearance order for each path.*

2.1 Path 1

Score: 373.35





Figure 1: Outline of path 1

2.1.1 Suzuki coupling of arylboronic acids with aryl iodides



Substrates:

- 1-Bromo-4-iodobenzene - *available at Sigma-Aldrich*
- (1H-Pyrrolo[2,3-b]pyridin-5-yl)boronic acid - *Combi-Blocks*

Products:

- Brc1ccc(-c2cnc3[nH]ccc3c2)cc1

Typical conditions: Pd catalyst.base.solvent


Protections: none

Yield: good

Reference: [10.1021/cr00039a007](https://doi.org/10.1021/cr00039a007) and [10.1007/3418_2012_32](https://doi.org/10.1007/3418_2012_32) and [10.1021/cr0505268](https://doi.org/10.1021/cr0505268) and [10.1016/j.jfluchem.2016.01.018](https://doi.org/10.1016/j.jfluchem.2016.01.018) and [10.1039/C3CS60197H](https://doi.org/10.1039/C3CS60197H)

Retrosynthesis ID: 25149

2.1.2 Iodination of aromatic compounds



Substrates:

1. BrC1CCC(-C2CNC3[NH]CCC3C2)CC1

Products:

1. BrC1CCC(-C2CNC3[NH]CC(I)C3C2)CC1

Typical conditions: I₂ or other iodinating agent e.g. NIS


Protections: none

Yield: good

Reference: DOI: [10.1039/C5SC00964B](https://doi.org/10.1039/C5SC00964B) and [10.1016/j.tetlet.2005.05.117](https://doi.org/10.1016/j.tetlet.2005.05.117) and [10.1007/s11178-005-0256-1](https://doi.org/10.1007/s11178-005-0256-1)

Retrosynthesis ID: 10697

2.1.3 N-alkylation of Heterocycles



Substrates:

1. a-Chlorotoluene - *available at Sigma-Aldrich*

2. BrC1CCC(-C2CNC3[nH]CC(I)C3C2)CC1

Products:

1. BrC1CCC(-C2CNC3C(C2)C(I)CN3Cc2ccccc2)CC1

Typical conditions: NaH.DMF


Protections: none

Yield: moderate

Reference: [10.1021/ol503625z](#) and [10.1081/SCC-120022467](#) (experimental) and [10.1021/ol2018328](#) (SI, p.5) and [10.1021/jo8026565](#) (SI, p.2)

Retrosynthesis ID: 28538

2.1.4 Synthesis of aryl Grignard reagents



Substrates:

1. magnesium
2. BrC1CCC(-C2CNC3C(C2)C(I)CN3Cc2ccccc2)CC1

Products:

1. Br[Mg]c1cn(Cc2ccccc2)c2ncc(-c3ccc(Br)cc3)cc12

Typical conditions: iPrMgCl.LiCl.THF or other conditions Mg.THF or tBuLi.MgBr2


Protections: none

Yield: moderate

Reference: DOI: [10.1016/S0040-4039\(99\)01404-5](#) and [10.1021/jo0000574](#) and WO2014123793 p.137 and [10.1021/jm400491x](#) and [10.3762/bjoc.12.36](#)

Retrosynthesis ID: 10011460

2.1.5 Arylation of sulfonamides with aryl iodides



Substrates:

1. 6-Bromo-2-fluoro-3-iodobenzaldehyde - *AOBChem*
2. Propane-1-sulfonamide - *Combi-Blocks*

Products:

1. CCCCS(=O)(=O)Nc1ccc(Br)c(C=O)c1F

Typical conditions: Cu.salt.diamine.base.DMF.heat


Protections: none

Yield: good

Reference: [10.1016/j.tetlet.2006.04.041](#) and [10.1016/j.tetlet.2011.10.113](#)
and [10.1016/j.tetlet.2005.08.149](#) and [10.1021/ol035942y](#) and [10.1021/acs.jmedchem.6b00685](#)

Retrosynthesis ID: 10012567

2.1.6 Grignard-Type Reaction



Substrates:

1. Br[Mg]c1cn(Cc2ccccc2)c2ncc(-c3ccc(Br)cc3)cc12
2. CCCS(=O)(=O)Nc1ccc(Br)c(C=O)c1F

Products:

1. CCCS(=O)(=O)Nc1ccc(Br)c(C(O)c2cn(Cc3ccccc3)c3ncc(-c4ccc(Br)cc4)cc23)c1F


Typical conditions: Mg or Li.ether

Protections: none

Yield: good

Reference: [10.1055/s-0030-1260809](#) or [10.1021/jm061429p](#) or [10.1021/jo0621423](#) or [10.1021/ja00373a036](#) or [10.1016/S0040-4020\(01\)00457-4](#)

Retrosynthesis ID: 25123

2.1.7 N-debenzylation of indoles**Substrates:**

1. CCCS(=O)(=O)Nc1ccc(Br)c(C(O)c2cn(Cc3ccccc3)c3ncc(-c4ccc(Br)cc4)cc23)c1F

Products:

1. CCCS(=O)(=O)Nc1ccc(Br)c(C(O)c2c[nH]c3ncc(-c4ccc(Br)cc4)cc23)c1F

Typical conditions: Na.NH₃


Protections: none

Yield: good

Reference: DOI: [10.1021/jo0110597](#)

Retrosynthesis ID: 356

2.1.8 Parikh-Doering Oxidation



Substrates:

1. CCCC(=O)(=O)Nc1ccc(Br)c(C(O)c2c[nH]c3ncc(-c4ccc(Br)cc4)cc23)c1F

Products:

1. CCCC(=O)(=O)Nc1ccc(Br)c(C(=O)c2c[nH]c3ncc(-c4ccc(Br)cc4)cc23)c1F

Typical conditions: DMSO, sulfur trioxide pyridine complex, NEt₃

Protections: none

Yield: good

Reference: [10.1021/ja00997a067](https://doi.org/10.1021/ja00997a067)

Retrosynthesis ID: 10257