How to Calculate the Total Amount of Your Salary?

- Jin Cai
- 2. Zeyu Dong
- 3. Yi'an Yu
- 4. Daoyuan Lai
- 5. Guoyang Qin
- 6. Iydon Liang

Department of Mathematics Southern University of Science and Technology

October 8, 2019

Outline



- Introduction
- 2 The Geometric Progression
- 3 Summation of Geometric Progression (Finite Case)
- Summation of Geometric Progression (Infinite Case)
- 5 Application to the Salary Example
- 6 Conclusion
- Acknowledgments
- 8 References



- Many jobs offer an annual cost-of-living increase to keep salaries consistent with inflation.
- A recent college graduate finds a position as a sales manager earning an annual salary of \$ 26,000.
- He is promised a 2% cost of living increase each year.
- So how could we determine how much this college graduate will get after *n*, which is a natural integer, years' hard work?
- This question arises naturally as payment is one of the most significant factors people will consider when finding a job.



- Many jobs offer an annual cost-of-living increase to keep salaries consistent with inflation.
- A recent college graduate finds a position as a sales manager earning an annual salary of \$ 26,000.
- He is promised a 2% cost of living increase each year.
- So how could we determine how much this college graduate will get after *n*, which is a natural integer, years' hard work?
- This question arises naturally as payment is one of the most significant factors people will consider when finding a job.



- Many jobs offer an annual cost-of-living increase to keep salaries consistent with inflation.
- A recent college graduate finds a position as a sales manager earning an annual salary of \$ 26,000.
- He is promised a 2% cost of living increase each year.
- So how could we determine how much this college graduate will get after *n*, which is a natural integer, years' hard work?
- This question arises naturally as payment is one of the most significant factors people will consider when finding a job.



- Many jobs offer an annual cost-of-living increase to keep salaries consistent with inflation.
- A recent college graduate finds a position as a sales manager earning an annual salary of \$ 26,000.
- He is promised a 2% cost of living increase each year.
- So how could we determine how much this college graduate will get after *n*, which is a natural integer, years' hard work?
- This question arises naturally as payment is one of the most significant factors people will consider when finding a job.



- Many jobs offer an annual cost-of-living increase to keep salaries consistent with inflation.
- A recent college graduate finds a position as a sales manager earning an annual salary of \$ 26,000.
- He is promised a 2% cost of living increase each year.
- So how could we determine how much this college graduate will get after *n*, which is a natural integer, years' hard work?
- This question arises naturally as payment is one of the most significant factors people will consider when finding a job.



- Many jobs offer an annual cost-of-living increase to keep salaries consistent with inflation.
- A recent college graduate finds a position as a sales manager earning an annual salary of \$ 26,000.
- He is promised a 2% cost of living increase each year.
- So how could we determine how much this college graduate will get after *n*, which is a natural integer, years' hard work?
- This question arises naturally as payment is one of the most significant factors people will consider when finding a job.

Outline



- Introduction
- The Geometric Progression
- 3 Summation of Geometric Progression (Finite Case)
- Summation of Geometric Progression (Infinite Case)
- 5 Application to the Salary Example
- 6 Conclusion
- Acknowledgments
- References



 Let's review the definition of arithmetic progression (Doe(2019)) first.

Definition 1

In an **arithmetic progression**, the difference between n^{th} term and $(n-1)^{th}$ term will be a constant which is known as the common difference of the arithmetic progression.

 What if the ratio of nth term to (n - 1)th term in a sequence is a constant?



 Let's review the definition of arithmetic progression (Doe(2019)) first.

Definition 1

In an **arithmetic progression**, the difference between n^{th} term and $(n-1)^{\text{th}}$ term will be a constant which is known as the common difference of the arithmetic progression.

What if the ratio of nth term to (n - 1)th term in a sequence is a constant?



 Let's review the definition of arithmetic progression (Doe(2019)) first.

Definition 1

In an **arithmetic progression**, the difference between n^{th} term and $(n-1)^{th}$ term will be a constant which is known as the common difference of the arithmetic progression.

• What if the ratio of n^{th} term to $(n-1)^{th}$ term in a sequence is a constant?



Definition 2

A **geometric progression** is a sequence in which each term is derived by multiplying or dividing the preceding term by a fixed number called the **common ratio**.

Definition 3

If we define the initial term of a sequence $\{a_n\}$ as a_1 , then $\{a_n\}$ is a **geometric progression** $(a_n \neq 0 \text{ for any } n)$ if and only if for any positive integer n,

$$\frac{a_{n+1}}{a_n} = r_1$$

where r is a constant.



Definition 2

A **geometric progression** is a sequence in which each term is derived by multiplying or dividing the preceding term by a fixed number called the **common ratio**.

Definition 3

If we define the initial term of a sequence $\{a_n\}$ as a_1 , then $\{a_n\}$ is a **geometric progression** $(a_n \neq 0 \text{ for any } n)$ if and only if for any positive integer n,

$$\frac{a_{n+1}}{a_n} = r,$$

where *r* is a constant.

Outline



- Introduction
- The Geometric Progression
- Summation of Geometric Progression (Finite Case)
- Summation of Geometric Progression (Infinite Case)
- 5 Application to the Salary Example
- 6 Conclusion
- Acknowledgments
- References



Definition 4

The sum of the terms in a geometric sequence is called a **geometric** series.

Theorem 1

The formula for S_n is

$$S_n = \frac{a_1(1-r^n)}{1-r}, \quad r \neq 1,$$

where S_n is the geometric series.

• How to derive this theorem?



Definition 4

The sum of the terms in a geometric sequence is called a **geometric** series.

Theorem 1

The formula for S_n is

$$S_n = \frac{a_1(1-r^n)}{1-r}, \quad r \neq 1,$$

where S_n is the geometric series.

• How to derive this theorem?



Definition 4

The sum of the terms in a geometric sequence is called a **geometric** series.

Theorem 1

The formula for S_n is

$$S_n = \frac{a_1(1-r^n)}{1-r}, \quad r \neq 1,$$

where S_n is the geometric series.

• How to derive this theorem?



Proof.

 Recall that a geometric progression is a sequence in which the ratio of any two consecutive terms is the common ratio, r. We can write the sum of the first n terms of a geometric series as

$$S_n = a_1 + ra_1 + r^2a_1 + \dots + r^{n-1}a_1.$$
 (3.1)

• We will begin by multiplying both sides of the equation by *r*,

$$rS_n = ra_1 + r^2a_1 + r^3a_1 + \dots + r^na_1.$$
 (3.2)

$$S_n - rS_n = (a_1 + ra_1 + r^2a_1 + \dots + r^{n-1}a_1) - (ra_1 + r^2a_1 + r^3a_1 + \dots + r^na_1).$$





Proof.

 Recall that a geometric progression is a sequence in which the ratio of any two consecutive terms is the common ratio, r. We can write the sum of the first n terms of a geometric series as

$$S_n = a_1 + ra_1 + r^2a_1 + \dots + r^{n-1}a_1.$$
 (3.1)

We will begin by multiplying both sides of the equation by r,

$$rS_n = ra_1 + r^2a_1 + r^3a_1 + \dots + r^na_1.$$
 (3.2)

$$S_n - rS_n = (a_1 + ra_1 + r^2a_1 + \dots + r^{n-1}a_1) - (ra_1 + r^2a_1 + r^3a_1 + \dots + r^na_1).$$





Proof.

 Recall that a geometric progression is a sequence in which the ratio of any two consecutive terms is the common ratio, r. We can write the sum of the first n terms of a geometric series as

$$S_n = a_1 + ra_1 + r^2a_1 + \dots + r^{n-1}a_1.$$
 (3.1)

• We will begin by multiplying both sides of the equation by *r*,

$$rS_n = ra_1 + r^2a_1 + r^3a_1 + \dots + r^na_1.$$
 (3.2)

$$S_n - rS_n = (a_1 + ra_1 + r^2a_1 + \dots + r^{n-1}a_1) - (ra_1 + r^2a_1 + r^3a_1 + \dots + r^na_1).$$



Proof.

 Recall that a geometric progression is a sequence in which the ratio of any two consecutive terms is the common ratio, r. We can write the sum of the first n terms of a geometric series as

$$S_n = a_1 + ra_1 + r^2a_1 + \dots + r^{n-1}a_1.$$
 (3.1)

• We will begin by multiplying both sides of the equation by r,

$$rS_n = ra_1 + r^2a_1 + r^3a_1 + \dots + r^na_1.$$
 (3.2)

$$S_n - rS_n = (a_1 + ra_1 + r^2a_1 + \dots + r^{n-1}a_1) - (ra_1 + r^2a_1 + r^3a_1 + \dots + r^na_1).$$





Proof.

- Notice that when we subtract, all but the first term of the top equation and the last term of the bottom equation are canceled.
- And it is trivial to obtain the formula for S_n,

$$S_n = \frac{a_1(1 - r^n)}{1 - r}, \quad r \neq 1.$$
 (3.3)





Proof.

- Notice that when we subtract, all but the first term of the top equation and the last term of the bottom equation are canceled.
- And it is trivial to obtain the formula for S_n ,

$$S_n = \frac{a_1(1 - r^n)}{1 - r}, \quad r \neq 1. \tag{3.3}$$





Proof.

- Notice that when we subtract, all but the first term of the top equation and the last term of the bottom equation are canceled.
- And it is trivial to obtain the formula for S_n,

$$S_n = \frac{a_1(1 - r^n)}{1 - r}, \quad r \neq 1.$$
 (3.3)





Proof.

- Notice that when we subtract, all but the first term of the top equation and the last term of the bottom equation are canceled.
- And it is trivial to obtain the formula for S_n,

$$S_n = \frac{a_1(1 - r^n)}{1 - r}, \quad r \neq 1.$$
 (3.3)



Outline



- Introduction
- The Geometric Progression
- Summation of Geometric Progression (Finite Case)
- Summation of Geometric Progression (Infinite Case)
- 5 Application to the Salary Example
- Conclusion
- Acknowledgments
- References



- The number of terms in infinite geometric progression will approach infinity $(n = \infty)$.
- The formal definition of infinity has been discussed in Zhang(2018).
- Sum of infinity geometric progression can only be defined at the range of $-1 < r < 1(r \neq 0)$ exclusive.

Theorem 2

$$S = \frac{a_1}{1 - r}$$



- The number of terms in infinite geometric progression will approach infinity $(n = \infty)$.
- The formal definition of infinity has been discussed in Zhang(2018).
- Sum of infinity geometric progression can only be defined at the range of $-1 < r < 1 (r \neq 0)$ exclusive.

Theorem 2

$$S = \frac{a_1}{1 - r}.$$



- The number of terms in infinite geometric progression will approach infinity $(n = \infty)$.
- The formal definition of infinity has been discussed in Zhang(2018).
- Sum of infinity geometric progression can only be defined at the range of $-1 < r < 1(r \neq 0)$ exclusive.

Theorem 2

$$S = \frac{a_1}{1 - r}$$



- The number of terms in infinite geometric progression will approach infinity $(n = \infty)$.
- The formal definition of infinity has been discussed in Zhang(2018).
- Sum of infinity geometric progression can only be defined at the range of $-1 < r < 1(r \neq 0)$ exclusive.

Theorem 2

$$S = \frac{a_1}{1 - r}.$$

Outline



- Introduction
- 2 The Geometric Progression
- Summation of Geometric Progression (Finite Case)
- Summation of Geometric Progression (Infinite Case)
- Application to the Salary Example
- 6 Conclusion
- Acknowledgments
- References



- There will be a 102% annual increment. So the common ratio *r* is 1.02.
- If we consider the first 5 years' total amount of the salary, we could calculate S_5 by directly substituting n = 5, r = 1.02, $a_1 = 26,000$ into Equation (3.3).
- We will get S_5 = 137, 907.06. I.e.This graduate will get \$137,907.06 after 5 years' working.



- There will be a 102% annual increment. So the common ratio *r* is 1.02.
- If we consider the first 5 years' total amount of the salary, we could calculate S_5 by directly substituting n = 5, r = 1.02, $a_1 = 26,000$ into Equation (3.3).
- We will get S_5 = 137, 907.06. I.e.This graduate will get \$137,907.06 after 5 years' working.



- There will be a 102% annual increment. So the common ratio *r* is 1.02.
- If we consider the first 5 years' total amount of the salary, we could calculate S_5 by directly substituting n = 5, r = 1.02, $a_1 = 26,000$ into Equation (3.3).
- We will get $S_5 = 137,907.06$. I.e.This graduate will get \$137,907.06 after 5 years' working.



- There will be a 102% annual increment. So the common ratio *r* is 1.02.
- If we consider the first 5 years' total amount of the salary, we could calculate S_5 by directly substituting n = 5, r = 1.02, $a_1 = 26,000$ into Equation (3.3).
- We will get $S_5 = 137,907.06$. I.e.This graduate will get \$137,907.06 after 5 years' working.

Outline



- Introduction
- The Geometric Progression
- Summation of Geometric Progression (Finite Case)
- Summation of Geometric Progression (Infinite Case)
- 5 Application to the Salary Example
- Conclusion
- Acknowledgments
- References

Conclusion



- We derived the formula of the geometric series.
- Furthermore, with a proper choice of the common ratio, we showed that this formula can be generalized to infinite case.
- A real problem is analyzed to illustrate the proposed formula and answered the question in the title: by using the formula proposed in this paper, we could calculate the total amount of the salary if it has a constant multiple of the common ratio each payment period.

Conclusion



- We derived the formula of the geometric series.
- Furthermore, with a proper choice of the common ratio, we showed that this formula can be generalized to infinite case.
- A real problem is analyzed to illustrate the proposed formula and answered the question in the title: by using the formula proposed in this paper, we could calculate the total amount of the salary if it has a constant multiple of the common ratio each payment period.

Conclusion



- We derived the formula of the geometric series.
- Furthermore, with a proper choice of the common ratio, we showed that this formula can be generalized to infinite case.
- A real problem is analyzed to illustrate the proposed formula and answered the question in the title: by using the formula proposed in this paper, we could calculate the total amount of the salary if it has a constant multiple of the common ratio each payment period.

Outline



- Introduction
- 2 The Geometric Progression
- 3 Summation of Geometric Progression (Finite Case)
- Summation of Geometric Progression (Infinite Case)
- 5 Application to the Salary Example
- 6 Conclusion
- Acknowledgments
- References

Acknowledgments



- The author would like to thank the Associate Editor and one referee for their valuable comments which have greatly improved the paper.
- The author's research is fully supported by the National Baby Natural Science Foundation of China(No. 1000001).

Acknowledgments



- The author would like to thank the Associate Editor and one referee for their valuable comments which have greatly improved the paper.
- The author's research is fully supported by the National Baby Natural Science Foundation of China(No. 1000001).

Thank you!

Outline



- Introduction
- The Geometric Progression
- Summation of Geometric Progression (Finite Case)
- Summation of Geometric Progression (Infinite Case)
- 5 Application to the Salary Example
- 6 Conclusion
- Acknowledgments
- References

Reference I



- [1] Doe,J.(2019). The Arithmetic Progression. *Mathematics for Beginners* 33:1465–1480.
- [2] Smith,J., Smith A.J., Smith B.J.(2019). Summation of Arithmetic Progression. *Journal of the Royal Baby Statistical Society, Series B* 63:111–126. 33:1465–1480.
- [3] Zhang,J.(2018). The Formal Discussion of Infinity. *Annals of Baby Mathematics* 9:705–724.