


## CHAPTER 14

# Fluid Mechanics



楊本立副教授

## Outline

1. Fluid statics
  - Pressure
  - Buoyant Forces and Archimedes's Principle
2. Fluid Dynamics
  - Bernoulli's Equation
  - Flow of viscous fluids in pipes
  - Applications of Fluid Dynamics
3. Surface tension

- 1. Fluid statics
  - Pressure
  - Buoyant Forces and Archimedes's Principle
- 2. Fluid Dynamics
  - Bernoulli's Equation
  - Flow of viscous fluids in pipes
  - Applications of Fluid Dynamics
- 3. Surface tension

## 1. Fluid statics – Pressure

535524245

6

## Fluid

- 3 A fluid is a collection of molecules that are randomly arranged and held together by **weak cohesive forces** and by forces exerted by the walls of a container.
- 3 Both **liquids** and **gases** are fluids.
- 3 **Fluids do not sustain shearing stresses or tensile stresses.**

535524245

7

## Pressure

- 3 The **force** exerted by a static fluid on an object is always **perpendicular** to the surfaces of the object.


Pressure:

$$P = \frac{F}{A_{\perp}}, \text{ a scalar}$$

SI unit: 1 Pascal (Pa)  $\equiv 1 \text{ N/m}^2$

The atmospheric pressure

$$P_0 = 1.00 \text{ atm} = 1.013 \times 10^5 \text{ Pa} = 1.013 \text{ bar}$$



© 2004 Thomson Deloche Cole

535524245

8




535524245

9

**Q:** How do you estimate the total mass of the air on the Earth? if the Earth radius is known.

$$\begin{aligned} F &= PA \\ \Rightarrow \int \rho g dV &= 1 \text{ atm} \times 4\pi R_E^2 \\ \Rightarrow m = \int \rho dV &\approx 1 \text{ atm} \times 4\pi R_E^2 / g \end{aligned}$$

☺ The gas pressure in a closed container is dependent on temperature. However, the ambient pressure is mainly caused by gravity, in addition to temperature. If without gravity, the air will escape from the Earth.




535524245

## Variation of Pressure with Depth

$$\sum \vec{F} = 0$$

$$PA - P_0 A - \rho Ahg = 0$$

$$\Rightarrow P = P_0 + \rho gh$$



p.s., The pressure variation comes from gravitation.


535524245

## Pascal's Law

- 3 The pressure in a fluid depends on depth and on the value of  $P_0$ .
- 3 An increase in pressure at the surface must be transmitted to every other point in the fluid.

535524245

<



Pascal vase

535524245

B


**Q:** You have invented a spacesuit with a straw passing through the faceplate so that you can drink from a glass while on the surface of a planet. Out on the surface of the Moon, you attempt to drink through the straw from an open glass of water. The value of  $g$  on the Moon is about one sixth of that on Earth.

Compared to the difficulty in drinking through a straw on Earth, you find **drinking through a straw on the Moon** to be (a) easier (b) equally difficult (c) harder (d) impossible.

535524245

44

**Q:**




© 2004 Thomson/Brooks Cole

$$\begin{array}{ccc} > & & > \\ P_A = P_B ? & & P_C = P_D ? \\ > & & > \end{array}$$

535524245

45

**Ans:**




$$P_C = P_0 - \rho_{Hg}gh_1$$

$$P_D = P_0 - \rho_{Hg}gh_1$$

$$\Rightarrow P_C = P_D$$

535524245

45



535524245


47

## Pressure Measurements

### Barometer

- Invented by Torricelli
- 1 atm = 760 mm (of Hg)
- 1 mm Hg = 1 torr

$$P_0 = \rho_{Hg}gh$$



$$h = \frac{P_0}{\rho_{Hg}g} = \frac{1.013 \times 10^5 \text{ Pa}}{(13.6 \times 10^3 \text{ kg/m}^3)(9.8 \text{ m/s}^2)} = 0.76 \text{ m}$$

535524245

43



535524245

43


## Manometer

3 Absolute pressure:

$$P = P_0 + \rho gh$$

3 Gauge pressure:


$$P - P_0 = \rho gh$$



535524245

4.

## Bourdon gauge



(b)




535524245

47

**Ex.** The force on a Dam

Water is filled to a height  $H$  behind a dam of width  $w$ . Determine the resultant force exerted by the water on the dam.



535524245

4c

**Ans:**

$$P = (P_0 + \rho gh) - P_0 = \rho g(H - y)$$

$$dF = PdA = \rho g(H - y)wdy$$


$$F = \int dF = \int_0^H \rho g(H - y)wdy$$

$$= \frac{1}{2} \rho g w H^2$$

535524245

53

## A hydraulic press



$$(1) P_1 = P_2$$

(assuming  $A_1$  and  $A_2$  at same height)

$$\Rightarrow \frac{F_1}{A_1} = \frac{F_2}{A_2} \Rightarrow F_1 = \left( \frac{A_1}{A_2} \right) F_2$$

$$(2) A_1 \Delta x_1 = A_2 \Delta x_2$$

Work done by force:

$$\Rightarrow F_1 \Delta x_1 = F_2 \Delta x_2$$

Energy is conserved.

- 1. Fluid statics
  - Pressure
  - Buoyant Forces and Archimedes's Principle
- 2. Fluid Dynamics
  - Bernoulli's Equation
  - Flow of viscous fluids in pipes
  - Applications of Fluid Dynamics
- 3. Surface tension


## 1. Fluid statics

### – Buoyant force & Archimedes's principle

## Buoyant Force



(a)




(b)

$$\vec{B} = \vec{F}_g$$

## Archimedes's Principle

- 3 The magnitude of the buoyant force always equals the weight of the fluid displaced by the object,  $B = M_{\text{fluid}}g$ .



$$\begin{aligned} B &= (P_b - P_t)A \\ &= (\rho_{\text{fluid}}gh)A \\ &= \rho_{\text{fluid}}gV \\ &= M_{\text{fluid}}g \end{aligned}$$



535524245


58

**Q:** An apple is held completely submerged just below the surface of a container of water. The apple is then moved to a deeper point in the water. Compared to the force needed to hold the apple just below the surface, the force needed to hold it at a deeper point is  
(a) larger (b) the same (c) smaller (d) impossible to determine.

535524245

59

**Q:**



What will be the movement of the person and the balloon when the bus stops suddenly?

Person -> forward

Balloon -> backward

535524245

5:





535524245

[https://physics.nyu.edu/~physlab/Demos/updatedEquipment  
/kinematicsForcesEnergy/forceParadox.html](https://physics.nyu.edu/~physlab/Demos/updatedEquipment/kinematicsForcesEnergy/forceParadox.html)

5;

## Totally submerged object



$$B = \rho_{\text{fluid}} g V = \rho_{\text{fluid}} g V_{\text{obj}}$$

The net force:

$$F_g = mg = \rho_{\text{obj}} g V_{\text{obj}}$$

$$F_g - B = (\rho_{\text{obj}} - \rho_{\text{fluid}}) g V_{\text{obj}}$$

⇒ The motion of an object in a fluid is determined by the **densities** of the fluid and the object

535524245

5<



535524245

63


## Galileo thermometer




53.5524245

64

**Ex.**



(a)



(b)

$$B + T_2 - F_g = 0$$

$$\rho_w g V_w + T_2 - \rho_c g V_c = 0$$

$$V_w = V_c$$


$$\Rightarrow \rho_c = \rho_w + \frac{T_2}{g V_w}$$

Use  $\rho_c$  to determine whether  
the crown is real or not.

53.5524245

65

## Floating Object



$$B = \rho_{\text{fluid}} g V_{\text{fluid}}$$

$$F_g = mg = \rho_{\text{obj}} g V_{\text{obj}}$$


$$B = F_g \Rightarrow \frac{\rho_{\text{obj}}}{\rho_{\text{fluid}}} = \frac{V_{\text{fluid}}}{V_{\text{obj}}}$$

535524245

66

## Hydrometer


(b) Using a hydrometer to measure the density of battery acid or antifreeze



535524245

67

Q:



$$\frac{V_{\text{water}}}{V_{\text{ice}}} = ?$$

$$\frac{V_{\text{water}}}{V_{\text{ice}}} = \frac{\rho_{\text{ice}}}{\rho_{\text{water}}} = \frac{917 \text{ kg/m}^3}{1030 \text{ kg/m}^3} = 89\%$$

535524245


68

Q: A glass of water contains a single floating ice cube. When the ice melts, does the water level (a) go up (b) go down (c) remain the same?

$$m_{\text{ice}} = \rho_{\text{water}} V_{\text{melted water}}$$

$$F_g = B$$

$$\Rightarrow V'_{\text{ice in water}} = \frac{m_{\text{ice}}g}{\rho_{\text{water}}g} = \frac{\rho_{\text{water}} V_{\text{melted water}}}{\rho_{\text{water}}} = V_{\text{melted water}}$$



535524245

⌚ Why would the sea level rise if iceberg melts?

69




<https://www.facebook.com/theactionlabofficial/>

535524245

6:

**Ex.**



$\rho_0$  : liquid density

$\rho_b$  : block density

$m_b$  : block mass

$m_0$  : liquid mass

535524245

6;


**Ans:**

$$(1) \quad f_1 = ?$$

$$f_1 = T$$

System =  $m_b$

$$= m_b g - B$$



$$= m_b g - \rho_0 V_b g$$

$$= m_b g - \rho_0 \frac{m_b}{\rho_b} g$$


$$\Rightarrow f_1 = m_b g - \frac{\rho_0}{\rho_b} m_b g$$

Note that,  $\rho_b = \rho_0 \Rightarrow f_1 = 0$

535524245

6<

$$(2) \quad f_2 = ?$$



System =  $m_b + m_0$

Buoyant force is an internal force here.

$$T + N - m_b g - m_0 g = 0$$

$$\Rightarrow f_2 = N = m_b g + m_0 g - (1 - \frac{\rho_0}{\rho_b}) m_b g$$

$$= m_0 g + \underbrace{\frac{\rho_0}{\rho_b} m_b g}_{\text{Buoyant}}$$

535524245

73

1. Fluid statics
  - Pressure
  - Buoyant Forces and Archimedes's Principle
2. Fluid Dynamics
  - Bernoulli's Equation
  - Flow of viscous fluids in pipes
  - Applications of Fluid Dynamics
3. Surface tension

## 2. Fluid dynamics

### – Bernoullie's equation



535524245

74

## Types of Fluid Flow

### 3 Laminar flow

- Steady flow.
- In a smooth path called a **streamline**.
- The paths of the different particles never cross each other.



535524245

75

### 3 Turbulent flow

- An irregular flow characterized by small whirlpool like regions.
- Turbulent flow occurs when the particles go above some **critical speed**.



535524245

76



<https://fb.watch/8RNkssEGun/>

535524245

77



535524245

78

## Viscosity

- ③ The degree of **internal friction** in the fluid.
- ③ This internal friction, ***viscous force***, is associated with the resistance that **two adjacent layers of fluid** have to moving relative to each other.
- ③ It causes part of the kinetic energy of a fluid to be converted to internal energy.

535524245

79

## Superfluid - ${}^4\text{He}$



Liquid helium enters  
superfluid phase  
transition at 2.17K.

Zero viscosity

535524245

<https://www.youtube.com/watch?v=-7PNacL4n8g>

7:



Glowing filaments in liquid He. Left: normal liquid. Right: superfluid.

No bubbling in superfluid liquid helium!

( Prof. Kimitoshi Kono's lab)

535524245

7:


## Ideal Fluid Flow

- (1) **The fluid is nonviscous** – internal friction is neglected.
- (2) **The flow is steady** – the velocity of each point remains constant.
- (3) **The fluid is incompressible** – the density remains constant.
- (4) **The flow is irrotational** – the fluid has no angular momentum about any point.

535524245

7<

## Equation of Continuity



$$\text{Mass flow rate: } \frac{dm}{dt} = \frac{\rho A dx}{dt} = \rho A v$$

$$\text{Volume flow rate: } \frac{dV}{dt} = \frac{Adx}{dt} = A v$$

$$(1) \text{ Mass conservation} \Rightarrow \rho_1 A_1 v_1 = \rho_2 A_2 v_2$$

$$(2) \text{ Incompressibility} \Rightarrow \rho_1 = \rho_2 = \rho$$

$$\Rightarrow A_1 v_1 = A_2 v_2 = \text{const.}$$

Equation of continuity (incompressible fluid)

For compressible fluid,  $\rho_1 A_1 v_1 = \rho_2 A_2 v_2$

535524245

83



© Cengage Learning/George Semple

535524245

84


Why is the diameter of the water stream different?



535524245

85

## Bernoulli's Equation



$$W_1 = F_1 \Delta x_1 = P_1 A_1 \Delta x_1 = P_1 V$$

$$W_2 = F_2 \Delta x_2 = -P_2 A_2 \Delta x_2 = -P_2 V$$

$$\Rightarrow W = W_1 + W_2 = (P_1 - P_2)V$$

$$\Delta K = \frac{1}{2} m v_2^2 - \frac{1}{2} m v_1^2$$

$$\Delta U = m g y_2 - m g y_1$$

535524245

86

### Mechanical energy conservation:

$$W = \Delta K + \Delta U$$

$$(P_1 - P_2)V = \left(\frac{1}{2}m v_2^2 - \frac{1}{2}m v_1^2\right) + (m g y_2 - m g y_1)$$

$$\rho = m / V$$

$$\Rightarrow P_1 - P_2 = \left(\frac{1}{2}\rho v_2^2 - \frac{1}{2}\rho v_1^2\right) + (\rho g y_2 - \rho g y_1)$$

$$\Rightarrow P_1 + \frac{1}{2}\rho v_1^2 + \rho g y_1 = P_2 + \frac{1}{2}\rho v_2^2 + \rho g y_2$$

$$\text{Bernoulli's equation: } P + \frac{1}{2}\rho v^2 + \rho g y = \text{const.}$$

535524245

87

$$\text{Bernoulli's equation: } P + \frac{1}{2} \rho v^2 + \rho g y = \text{const.}$$

- ⌚ It is a result of energy conservation.
- ⌚ The constant depends on the initial conditions of  $P$ ,  $\rho$ , and  $y$ .
- ⌚ Bernoulli's equation reduces to Pascal's law when  $v = 0$ .

535524245

88

The **Venturi tube** can be used to measure the flow speed of an incompressible fluid.




535524245

89

**Q:** Determine the flow speed at point 2 if the height  $h$  is known.

Difference in height results from reduced pressure in throat (point 2).



535524245

8:

**Ans:**

$$\text{Bernoulli's equation: } P + \frac{1}{2} \rho v^2 + \rho g y = \text{const.}$$

$$y_1 = y_2$$

$$\text{Bernoulli's eq.} \Rightarrow P_1 + \frac{1}{2} \rho v_1^2 = P_2 + \frac{1}{2} \rho v_2^2$$


$$\text{Continuity eq.} \Rightarrow A_1 v_1 = A_2 v_2$$

$$\Rightarrow v_2 = A_1 \sqrt{\frac{2(P_1 - P_2)}{\rho(A_1^2 - A_2^2)}} = A_1 \sqrt{\frac{2\rho g h}{\rho(A_1^2 - A_2^2)}}$$

535524245

8:

**Ex.** An enclosed tank containing a liquid of density  $\rho$  has a hole in its side at a distance  $y_1$  from the tank's bottom. The hole is open to the atmosphere, and its diameter is much smaller than the diameter of the tank. The air above the liquid is maintained at a pressure  $P$ . Determine the speed of the liquid as it leaves the hole when the liquid's level is a distance  $h$  above the hole.



535524245

8<


**Ans:**

$$(1) \quad A_1 \ll A_2 \quad \& \quad A_1 v_1 = A_2 v_2 \\ \Rightarrow v_2 \ll v_1$$

$$(2) \quad P_0 + \frac{1}{2} \rho v_1^2 + \rho g y_1 = P + \frac{1}{2} \rho v_2^2 + \rho g y_2$$

$$y_2 - y_1 = h \quad \Rightarrow \quad v_1 = \sqrt{\frac{2(P - P_0)}{\rho} + 2gh}$$

$$\text{Torricelli's law: } P = P_0 \Rightarrow v_1 = \sqrt{2gh}$$



535524245

93


☺ If the tank is open to the atmosphere and sitting on a table, what position of the hole would cause the water to land on the table at the farthest distance from the tank?

$$0 = y_1 + 0 - \frac{1}{2}gt^2 \Rightarrow t = \sqrt{\frac{2y_1}{g}}$$

$$\begin{aligned} x_f &= x_i + v_{xi}t = 0 + \sqrt{2g(y_2 - y_1)} \sqrt{\frac{2y_1}{g}} \\ &= 2\sqrt{(y_2 y_1 - y_1^2)} \end{aligned}$$

$$\Rightarrow \frac{dx_f}{dy_1} = \frac{1}{2}(2)(y_2 y_1 - y_1^2)^{-1/2} (y_2 - 2y_1) = 0$$

$$\Rightarrow y_1 = \frac{1}{2}y_2$$



535524245

94




535524245

95



535524245

96

**Ex.**

(Ignore the water vapor pressure)

$$(1) \quad P_0 + 0 + 0 = P_0 - \rho gh + \frac{1}{2} \rho v^2 \quad \Rightarrow \quad v = \sqrt{2gh} \quad \text{indep. of } y.$$

$$(2) \quad P_0 + 0 + 0 = P + \rho gy + \frac{1}{2} \rho v^2 \quad \Rightarrow \quad y = \frac{P_0 - P - \frac{1}{2} \rho v^2}{\rho g}$$

$$P \geq 0, \quad v \geq 0 \quad \Rightarrow \quad y_{\max} \leq \frac{P_0}{\rho g}$$

535524245

97

1. Fluid statics
  - Pressure
  - Buoyant Forces and Archimedes's Principle
2. Fluid Dynamics
  - Bernoulli's Equation
  - Flow of viscous fluids in pipes
  - Applications of Fluid Dynamics
3. Surface tension

## 2. Fluid dynamics

### – Flow of viscous fluids in pipes

535524245

98

## Viscosity

In reality, the **viscosity** of fluids is not negligible.

The pressure differential to keep fluid moving at a fixed speed in a horizontal pipe is

$$\Delta P = I_v R,$$



$$\Delta P = P_1 - P_2 \neq 0$$


$I_v$  : the volume flow rate ( $= Av$ )

$R$  : the measure of resistance


- It is a typical transport equation, similar to the electrical current flow  $\Delta V = IR$ , and the diffusion due to concentration difference.

535524245

99



Cross section of a cylindrical pipe



The velocity profile for viscous fluid flowing in the pipe has a parabolic shape.

The resistance  $R$  can be shown as

$$R = \frac{8\eta L}{\pi r^4} \quad \eta : \text{viscosity (unit: poise)}$$

$$\Delta P = I_V R, \quad \Rightarrow \quad \Delta P = \frac{8\eta L}{\pi r^4} I_V$$

$$\Rightarrow \quad \Delta P \propto \frac{1}{r^4} !$$

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Addison Wesley

9:

1. Fluid statics
  - Pressure
  - Buoyant Forces and Archimedes's Principle
2. Fluid Dynamics
  - Bernoulli's Equation
  - Flow of viscous fluids in pipes
  - Applications of Fluid Dynamics
3. Surface tension

## 2. Fluid dynamics


– Applications of fluid dynamics

535524245

9;

## Applications of Fluid Dynamics

- 3 Streamline flow around a moving airplane wing.
- 3 Lift is the upward force on the wing from the air.
- 3 Drag is the resistance.
- 3 The lift depends on the speed of the airplane, the area of the wing, its curvature, and the angle between the wing and the horizontal.




535524245

9<

(a) Flow lines around an airplane wing

The flowlines of air moving over the top of the wing are crowded together, so the flow speed is higher and the pressure consequently lower.



An equivalent explanation: The wing's shape imparts a net downward momentum to the air, so the reaction force on the airplane is upward.

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Addison-Wesley.

535524245

(b) Computer simulation of airflow around an airplane wing






Image of air parcels flowing around a wing, showing that the air goes much faster over the top than over the bottom (and that air parcels which are together at the leading edge of the wing do *not* meet up at the trailing edge!)

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Addison-Wesley.

:3






535524245

:6




535524245

:7



Golf ball



Atomizer

535524245


:8



535524245


:9

A clockwise spinning ball experiences a clockwise deflection.



535524245

..



535524245

<https://kknews.cc/news/qjmnklr.html>

..

1. Fluid statics
  - Pressure
  - Buoyant Forces and Archimedes's Principle
2. Fluid Dynamics
  - Bernoulli's Equation
  - Flow of viscous fluids in pipes
  - Applications of Fluid Dynamics
3. Surface tension

### 3. Surface tension

535524245

<

#### Surface tension:

- A molecule at the surface of a liquid is attracted into the bulk liquid, which tends to reduce the liquid's surface area.
- Surface tension can be changed with **temperature** or by adding certain **chemical** into the liquid.




535524245

;3

Molecules in a liquid are attracted by neighboring molecules.

At the surface, the unbalanced attractions cause the surface to resist being stretched.




535524245

;4

### Capillary force vs. surface tension


535524245

;5



<https://www.youtube.com/watch?v=0l6bBB3zuGc>

### Rheological Behavior of Fluids



535524245

;6

### Fun with Non-Newtonian Fluid



535524245

<https://www.youtube.com/watch?v=RIUEZ3AhrVE>

;7