


## Chapter 12

# Gravitation



楊本立副教授

## Outline

1. Newton's law of universal gravitation
2. Gravitational field
3. Gravitational potential energy
4. Kepler's laws of planetary motion


1. Newton's law of universal gravitation
2. Gravitational field
3. Gravitational potential energy
4. Kepler's laws of planetary motion

## 1. Newton's law of universal gravitation

2022/11/7

3

Galileo



Faster or slower than the  
stone alone?


2022/11/7

4

## Newton's law of universal gravitation

$$F_g = G \frac{m_1 m_2}{r^2}$$

$$\vec{F}_{12} = (-)G \frac{m_1 m_2}{r^2} \hat{r}_{12}, \quad \hat{r}_{12} = \hat{r}_2 - \hat{r}_1$$




- Attractive, field force
- inverse-square law
- $G$ : universal gravitational constant
- $G=6.673\times10^{-11}\text{N}\cdot\text{m}^2/\text{kg}^2$

2022/11/7

5

- o The gravitation force exerted by a finite-size, spherically symmetric mass distribution on a particle outside the distribution is the same as if the entire mass of the distribution were concentrated at the center.




2022/11/7

6

Experimental proof of Newton's law:

(1) Newton's inverse square law



$$\left\{ \begin{array}{l} a_M = \frac{F_M}{m_M} = G \frac{M_E}{r_M^2} \\ g = \frac{F_g}{m_A} = G \frac{M_E}{R_E^2} \end{array} \right. \Rightarrow a_M = \left( \frac{R_E}{r_M} \right)^2 g$$

(2) Centripetal acceleration


$$a_M = \frac{v^2}{r_M} = \frac{(2\pi r_M / T)^2}{r_M} = \frac{4\pi^2 r_M}{T^2}$$

Newton verified (1)=(2).

2022/11/7

7

Cavendish apparatus for measuring  $G$




2022/11/7

8

## Free-Fall Acceleration

$$\begin{aligned} m\vec{a} &= \vec{F}_g \\ \Rightarrow mg &= \frac{GM_E m}{(R_E + h)^2} \\ \Rightarrow g &= \frac{GM_E}{(R_E + h)^2} \end{aligned}$$



$g \searrow$  as  $h \nearrow$

2022/11/7

9

1. Newton's law of universal gravitation
2. Gravitational field
3. Gravitational potential energy
4. Kepler's laws of planetary motion


## 2. Gravitational field

2022/11/7

10

## The Gravitational Field

The inverse-square law can be understood by the “field” concept.



$$\vec{g} \equiv \frac{\vec{F}_g}{m} \quad \vec{g}: \text{gravitational field}$$

$$\vec{g} = -\frac{GM}{r^2} \hat{r}$$

$|\vec{g}| \propto$  the density of field lines  
(number of lines per unit area).

The number of field lines  $N \propto M$ .


$$\Rightarrow |\vec{g}| \propto \frac{N}{4\pi r^2} \propto \frac{M}{r^2}$$

2022/11/7


11

- The Gravitational field  $g$  inside a Spherical Shell:

$$\begin{cases} \vec{g} = -\frac{GM}{r^2} \hat{r}, & r > R \\ \vec{g} = 0, & r < R \end{cases}$$



Proof:



$$\vec{g}_1 = -\frac{Gm_1}{r_1^2} \cos \theta \hat{r}_1 = -\frac{G\sigma A_1 \cos \theta}{r_1^2} \hat{r}_1 = -G\sigma \Omega \hat{r}_1$$

$$\vec{g}_2 = -\frac{Gm_2}{r_2^2} \cos \theta \hat{r}_2 = -\frac{G\sigma A_2 \cos \theta}{r_2^2} \hat{r}_2 = -G\sigma \Omega \hat{r}_2$$

$$\hat{r}_1 = -\hat{r}_2 \Rightarrow \vec{g}_1 = -\vec{g}_2$$

$$\Omega \equiv \frac{A_\perp}{r^2} : \text{solid angle}$$

$$\sigma : \text{surface density}$$

$$\text{So } \vec{g} = \sum_i \vec{g}_i = 0, \text{ for } r < R$$

2022/11/7



12

- The Gravitational field  $g$  inside a **solid sphere**:

$$g(r < R) = \int_0^r \frac{\rho(r')}{(r-r')^2} dV' + \underbrace{\int_r^R \frac{\rho(r')}{(r-r')^2} dV'}_{=0} \Rightarrow g(r) = \frac{GM'}{r^2} = \frac{GM}{r^2} \frac{r^3}{R^3}$$

$$M' = M \frac{\frac{4}{3}\pi r^3}{\frac{4}{3}\pi R^3} = M \frac{r^3}{R^3}$$

$$\Rightarrow \bar{g}(r) = -\frac{GM}{R^3} r \hat{r}, \text{ for } r < R.$$



2022/11/7

13

☺ The above results can also be derived from the gravitational potential energy approach. Please refer to the textbook (Young, p. 401).

2022/11/7

14

1. Newton's law of universal gravitation
2. Gravitational field
3. **Gravitational potential energy**
4. Kepler's laws of planetary motion

### 3. Gravitational potential energy

2022/11/7

15

The gravitational force is "conservative".

$$\vec{F}(r) = F(r)\hat{r}$$

$$dW = \vec{F} \cdot d\vec{\ell}$$


$$= (\vec{F}\hat{r}) \cdot (dr\hat{r} + rd\theta\hat{\theta} + r\sin\theta d\phi\hat{\phi})$$

$$= F(r)dr$$

$\because W = \int_{r_i}^{r_f} F(r)dr = -\Delta U(r)$  depends

only on the end points (indep. of  $\theta$  and  $\phi$ ),

$\Rightarrow F(r)$  is called "**conservative force**".




Any central force  $\vec{F}(r) = F(r)\hat{r}$  is a conservative force.

2022/11/7

16

## Gravitational Potential Energy



$$\Delta U = U_f - U_i = - \int_{r_i}^{r_f} \vec{F}_g(r) \cdot d\vec{r}$$

$$\vec{F}_g(r) = -\frac{GM_E m}{r^2} \hat{r}$$

$$\Rightarrow \Delta U = \int_{r_i}^{r_f} \frac{GM_E m}{r^2} dr$$


$$= -GM_E m \left( \frac{1}{r_f} - \frac{1}{r_i} \right)$$

$$\text{Set } r_i = \infty, \quad U_i = 0 \quad \Rightarrow \quad U(r) = -\frac{GM_E m}{r}$$

2022/11/7

17

**Ex.**



$$U_{total} = U_{12} + U_{13} + U_{23}$$


$$= -G \left( \frac{m_1 m_2}{r_{12}} + \frac{m_1 m_3}{r_{13}} + \frac{m_2 m_3}{r_{23}} \right)$$

2022/11/7

18

For a small vertical distance  $\Delta y$  near the earth's surface,

$$\begin{aligned}\Delta U &= -GM_E m \left( \frac{1}{r_f} - \frac{1}{r_i} \right) \\ &= GM_E m \left( \frac{r_f - r_i}{r_i r_f} \right) \\ &\approx \left( \frac{GM_E m}{R_E^2} \right) \Delta y\end{aligned}$$



$$\Rightarrow \Delta U = mg\Delta y$$


2022/11/7

19

## Total Energy in Planetary Motion

$$E = K + U = \frac{1}{2}mv^2 - \frac{GM_E m}{r}$$

$$\frac{GM_E m}{r^2} = m \frac{v^2}{r} \quad \Rightarrow \quad \frac{1}{2}mv^2 = \frac{GM_E m}{2r}$$




$$\Rightarrow \text{Circular orbits: } E = -\frac{GM_E m}{2r}$$

$$\left( \text{Elliptical orbits: } E = -\frac{GM_E m}{2a}, \quad a: \text{semi-major axis} \right)$$

2022/11/7

20




$$K \geq 0 \Rightarrow \begin{cases} E < 0, \text{ bound} \\ E \geq 0, \text{ unbound} \end{cases}$$

2022/11/7

21

### Gravitational Potential Energy for the Earth



Binding energy:

$$U(\infty) - U(r) = 0 - \left(-\frac{GM_E m}{r}\right) = \frac{GM_E m}{r}$$

2022/11/7

22

## Escape Speed


$$\frac{1}{2}mv_i^2 - \frac{GM_E m}{R_E} = \frac{1}{2}mv_f^2 - \frac{GM_E m}{r_{\max}}$$

$$\geq -\frac{GM_E m}{r_{\max}}$$

Escape,  $r_{\max} \rightarrow \infty \Rightarrow v_i \geq \sqrt{\frac{2GM_E}{R_E}}$

Escape speed  $\Rightarrow v_{esc} = \sqrt{\frac{2GM_E}{R_E}} = \sqrt{2gR_E}$

indep. of object's mass.



2022/11/7

23


氦气不断从地球上逃逸。

$$\because v_{rms} = \sqrt{\frac{3k_B T}{m}} > v_{esc} = \sqrt{\frac{2GM_E}{R_E}} = 1.12 \times 10^4 \text{ m/s}$$

2022/11/7

24

## Black hole



Light can't escape from black holes if  $v_{esc} < c$ .

$$\text{When } v_{esc} = \sqrt{\frac{2GM}{R_s}} = c,$$

$$\Rightarrow R_s = \frac{2GM}{c^2} \quad \text{Schwarzschild radius}$$

2022/11/7

25

💡 Black hole doesn't have to be very massive, it is the **density** that determines the radius  $R_s$  to be  $> R_{\text{black hole}}$ .

$$\text{Schwarzschild radius } R_s = \frac{2GM}{c^2} = \frac{8\pi G}{3c^2} \rho R_{\text{black hole}}^3$$

$$\Rightarrow \rho \text{ has to be large so that } R_s > R_{\text{black hole}}.$$

2022/11/7


26

## The formation of black holes

- o “**Supernova**” – the catastrophic explosion of a heavy massive star.
- o The core mass after explosion is  $m$  and the Sun mass is  $m_s$ 
  - $m < 1.4 m_s$ , gradually cool down, **white dwarf star**.
  - $m > 1.4 m_s$ , collapse further, **neutron star** (radius  $\sim 10$  km).
  - $m > 3 m_s$ , collapse into a **black hole**.

2022/11/7

27





### Properties of black holes

- “X-rays” are characteristic of a black hole.
- Tidal force.
- Gravitational red shift, time dilation.

2022/11/7

28




$$m_X = \frac{4\pi^2 a^3}{GT^2}$$

Supermassive black holes may exist at the center of galaxies.

2022/11/7

29

On April 10, 2019, the Event Horizon Telescope collaboration ([eventhorizontelescope.org](http://eventhorizontelescope.org)) released the first direct image of a black hole.






[<https://www.aps.org/publications/apsnews/201906/black-hole.cfm>]

2022/11/7

30

## Dark matter

Luminous matters (stars) contribute major mass of the universe.  
Non-luminous matters (planets, dust) are only a small fraction.




2022/11/7

31

## Dark energy


Dark energy is proposed to explain an increasing acceleration in the expansion of the Universe.



2022/11/7

32


## The General Theory of Relativity



- Einstein proposed that *no* experiment, mechanical or otherwise, could distinguish between the two situations. **Principle of equivalence: gravitation and acceleration are equivalent.**

2022/11/7

33





- Einstein assumes that all the laws of nature (including light) have the same form for observers in any reference frame, whether accelerated or not.

2022/11/7

34

## Gravitational lens



Courtesy National Radio Astronomy Observatory

2022/11/7

35




- Two postulates in Einstein's **general theory of relativity**
  - All the laws of nature have the same form for observers in any frame of reference, whether accelerated or not.
  - In the vicinity of any point, a gravitational field is equivalent to an accelerated frame of reference in gravity-free space (**the principle of equivalence**).



2022/11/7

36

## Gravitation $\Rightarrow$ Curvature of space and time



2022/11/7

37

1. Newton's law of universal gravitation
2. Gravitational field
3. Gravitational potential energy
4. **Kepler's laws of planetary motion**


## 4. Kepler's laws of planetary motion

2022/11/7

38


## Kepler's 1st Law

- o All planets move in **elliptical orbits** with the Sun at one focus.
  - This is a direct result of the inverse-square of the Gravitational force.





2022/11/7

39




2022/11/7

40



- Sun is located at focus.
- Pluto has the highest eccentricity  $e_{\text{Pluto}} = 0.25$  among the 8 planets in the solar system.
- Halley's comet has an orbit with high eccentricity  $e_{\text{Halley's comet}} = 0.97$ .



2022/11/7

43

(Solar) planets 1930–2006									
1	2	3	4	5	6	7	8	9	
Mercury ☿	Venus ♀	Earth ⊕	Mars ♂	Jupiter ♃	Saturn ♄	Uranus ♅	Neptune ♆	Pluto ♇	

My Very Eager Mother Just Serves Us Nine Pizza!

2022/11/7

44

## Kepler's 2nd Law


- o The radius vector drawn from the Sun to a planet sweeps out equal areas in equal time intervals.

$$\Rightarrow \frac{dA}{dt} = \frac{|\vec{L}|}{2M_p} = \text{const.}$$

- This is a consequence of **angular momentum conservation**.
- This law applies to any situation that involves a central force, whether inverse-square or not.
- The orbit lies in the same plane.


2022/11/7

45



(a)

$$\begin{aligned}\vec{\tau} &= \vec{r} \times \vec{F} = \vec{r} \times F_g(r)\hat{r} = 0 \\ \Rightarrow \frac{d\vec{L}}{dt} &= 0 \\ \vec{L} &= \vec{r} \times \vec{p} = \vec{r} \times M_p \vec{v} = \text{const.}\end{aligned}$$



(b)

$$\begin{aligned}dA &= \frac{1}{2} |\vec{r} \times d\vec{r}| = \frac{1}{2} |\vec{r} \times \vec{v}| dt \\ &= \frac{|\vec{L}|}{2M_p} dt \\ \Rightarrow \frac{dA}{dt} &= \frac{|\vec{L}|}{2M_p} = \text{const.}\end{aligned}$$

2022/11/7

46

## Kepler's 3rd Law

- The square of the orbital period of any planet is proportional to the cube of the semi-major axis of the elliptical orbit.

$$T^2 = \left( \frac{4\pi^2}{GM_s} \right) a^3, \text{ for elliptical orbits.}$$

2022/11/7


47

For circular orbit,  $\vec{F}_g = m\vec{a}_c$

$$\frac{GM_s M_p}{r^2} = \frac{M_p v^2}{r}$$

$$\Rightarrow \frac{GM_s}{r^2} = \frac{(2\pi r/T)^2}{r}$$

$$\Rightarrow T^2 = \left( \frac{4\pi^2}{GM_s} \right) r^3 \equiv K_s r^3$$



In general,

$$T^2 = \left( \frac{4\pi^2}{GM_s} \right) a^3, \text{ for elliptical orbits.}$$


indep. of planet's mass and eccentricity.

2022/11/7

48

### Ex. Geosynchronous Satellite

- (1)  $v = ?$  at  $h$ .
- (2) Geosynchronous satellite  $r = ?$   $v = ?$



2022/11/7

49

**Ans:**

$$(1) \vec{F} = m\vec{a}_c$$

$$\frac{GM_E m}{r^2} = m \frac{v^2}{r} \Rightarrow v = \sqrt{\frac{GM_E}{R_E + h}}$$

$$(2) T^2 = \left( \frac{4\pi^2}{GM_E} \right) r^3 = (24 \text{ hrs})^2$$


$$\Rightarrow r = \dots$$

$$v = \frac{2\pi r}{T} = \frac{2\pi r}{24 \text{ hrs}} \Rightarrow v = \dots$$

2022/11/7

50

**Ex:** Astronomers detect a distant meteoroid moving along a straight line that, if extended, would pass at a distance  $3R_E$  from the center of the Earth, where  $R_E$  is the radius of the Earth. What minimum speed must the meteoroid have if the Earth's gravitation is not to deflect the meteoroid to make it strike the Earth?



2022/11/7

51

$$\text{Not to strike} \Rightarrow R \geq R_E$$

$$(1) \ L_i = L_f \Rightarrow mv_i(3R_E) = mv_f R \\ \Rightarrow v_f = (3R_E / R)v_i \Rightarrow v_f \leq 3v_i$$

$$(2) \ E_i = E_f \Rightarrow \frac{1}{2}mv_i^2 = \frac{1}{2}mv_f^2 - \frac{GM_E m}{R} \\ \Rightarrow v_f^2 = v_i^2 + \frac{2GM_E}{R} \\ \text{So } v_i \geq \frac{1}{2} \sqrt{\frac{GM_E}{R_E}}$$

2022/11/7

52

**5 Ways to Find a Planet**

- WATCHING FOR WOBBLE**  
Radial Velocity  
536 planets discovered
- SEARCHING FOR SHADOWS**  
Transit  
306 planets discovered
- TAKING PICTURES**  
Direct Imaging  
38 planets discovered
- LIGHT IN A GRAVITY LENS**  
Gravitational Microlensing  
18 planets discovered
- MINISCULE MOVEMENTS**  
Astrometry  
2 planets discovered

<https://exoplanets.nasa.gov/interactive/11/>

2022/11/7 53

**Planetary transit**

2009 Feb 10 18:15:44 UT  
Location: Center of Earth  
Face: Jupiter  
Field: 35.1" x 36.4"

The diagram illustrates a planetary transit. On the left, there is an image of the planet Jupiter. To its right, a yellow circle represents the star, with three blue dots labeled 1, 2, and 3 representing the planet's position at different times during its orbit. A dashed horizontal line extends from the star through the planet. Below this, a graph shows the "light curve" with "Brightness" on the vertical axis and "Time" on the horizontal axis. The curve remains constant at a baseline level until point 1, then drops to a lower level at point 2 (the transit), and returns to the baseline at point 3.

2022/11/7 54

## Negative mass?

$$\vec{F}_g = (-m)\vec{g} = (-m)\vec{a}$$

$\Rightarrow \vec{F}_g$  is opposite to  $\vec{a}$  direction!

[“Negative mass can be positively amusing” by Richard H. Price,  
American Journal of Physics 61, 216 (1993);  
View online: <https://doi.org/10.1119/1.17293>]

2022/11/7

55

⌚ What would happen to the earth if the sun suddenly disappeared?




<https://fb.watch/8RNhBCxTgE/>

2022/11/7

56

## Significant Solar Flare Erupts From Sun



[https://blogs.nasa.gov/solarcycle25/wp-content/uploads/sites/304/2022/03/FlareX13\\_20220330\\_171A\\_2048p30\\_Sq.gif](https://blogs.nasa.gov/solarcycle25/wp-content/uploads/sites/304/2022/03/FlareX13_20220330_171A_2048p30_Sq.gif)

2022/11/7

57