

Coordinate Geometry (9 th std)

Choose the Appropriate Answer

1.If the y-coordinate of a point is zero, then the point always lies

- (a) In the I quadrant (b) In the II quadrant
(c) On x - axis (d) On y -axis

Ans: (c) On x - axis

2.The points (-5, 2) and (2, -5) lie in the

- (a) Same quadrant (b) II and III quadrant respectively
(c) II and IV quadrant respectively (d) IV and II quadrant respectively

Ans: (c) II and IV quadrant respectively

3.On plotting the points O (0, 0), A (3, -4), B (3, 4) and C (0, 4) and joining OA, AB, BC and CO, which of the following figure is obtained?

- (a) Square (b) Rectangle (c) Trapezium (d) Rhombus

Ans: (c) Trapezium

4.If P (-1, 1), Q (3, -4), R (1, -1), S (-2, -3) and T (- 4, 4) are plotted on a graph paper, then the points in the fourth quadrant are

- (a) P and T (b) Q and R (c) Only S (d) P and Q

Ans: (b) Q and R

5.The point whose ordinate is 4 and which lies on the Y - axis is

- (a) (4, 0) (b) (0, 4) (c) (1, 4) (d) (4, 2)

Ans: (b) (0, 4)

6.The distance between the two points (2, 3) and (1, 4) is

- (a) 2 (b) $\sqrt{56}$ (c) $\sqrt{10}$ (d) $\sqrt{2}$

Ans: (d) $\sqrt{2}$

$$\sqrt{(1-2)^2 + (4-3)^2} = \sqrt{(-1)^2 + 1^2} = \sqrt{1+1} = \sqrt{2}$$

7.If the points A (2, 0), B (-6, 0), C (3, a – 3) lie on the X - axis then the value of 'a' is

- (a) 0 (b) 2 (c) 3 (d) -6

Ans: (c) 3

8.If $(x + 2, 4) = (5, y - 2)$, then the coordinates (x, y) are

- (a) (7, 12) (b) (6, 3) (c) (3, 6) (d) (2, 1)

Ans: (c) (3, 6)

$$x + 2 = 5 \Rightarrow x = 5 - 2 = 3, \quad 4 = y - 2 \Rightarrow y = 4 + 2 = 6$$

9.If Q₁, Q₂, Q₃, Q₄ are the quadrants in a Cartesian plane then Q₂ ∩ Q₃ is

- (a) Q₁ ∪ Q₂ (b) Q₂ ∪ Q₃ (c) Null set (d) Negative x-axis

Ans: (c) Null set

10. The distance between the point (5, -1) and the origin is

- (a) $\sqrt{24}$ (b) $\sqrt{37}$ (c) $\sqrt{26}$ (d) $\sqrt{17}$

Ans: (c) $\sqrt{26}$

$$d = \sqrt{(5-0)^2 + (-1-0)^2} = \sqrt{5^2 + (-1)^2} = \sqrt{25+1} = \sqrt{26}$$

11. The coordinates of the point C dividing the line segment joining the points P (2, 4) and Q (5, 7) internally in the ratio 2 : 1 is

- (a) $(7/2, 11/2)$ (b) (3, 5) (c) (4, 4) (d) (4, 6)

Ans: (d) (4, 6)

A line divides internally in the ratio m : n $\Rightarrow$ m = 2, n = 1

$$\text{The point P} = \left(\frac{mx_2 + nx_1}{m+n}, \frac{my_2 + ny_1}{m+n} \right) = \left(\frac{10+2}{3}, \frac{14+4}{3} \right) = \left(\frac{12}{3}, \frac{18}{3} \right) = (4, 6)$$

12. If P (a / 3, b / 2) is the mid - point of the line segment joining A (-4, 3) and B (-2, 4) then (a, b) is

- (a) (-9, 7) (b) (-3, 7 / 2) (c) (9, -7) (d) (3, -7 / 2)

Ans: (a) (-9, 7)

$$\text{Mid point of a line} = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)$$

$$\left(\frac{a}{3}, \frac{b}{2} \right) = \left(\frac{-4-2}{2}, \frac{3+4}{2} \right) = \left(-3, \frac{7}{2} \right)$$

$$\frac{a}{3} = -3 \Rightarrow a = -9 \quad \frac{b}{2} = \frac{7}{2} \Rightarrow b = 7 \quad (a, b) \text{ is } (-9, 7)$$

13. In what ratio does the point Q (1, 6) divide the line segment joining the points P (2, 7) and R (-2, 3)

- (a) 1 : 2 (b) 2 : 1 (c) 1 : 3 (d) 3 : 1

Ans: (c) 1 : 3

A line divides internally in the ratio m : n the point P =

$$\left(\frac{mx_2 + nx_1}{m+n}, \frac{my_2 + ny_1}{m+n} \right) = (1, 6) = \left(\frac{-2m+2n}{m+n}, \frac{3m+7n}{m+n} \right)$$

$$\frac{-2m+2n}{m+n} = 1 \Rightarrow -2m + 2n = m + nx \Rightarrow 3m = n \Rightarrow \frac{m}{n} = \frac{1}{3}$$

14. If the coordinates of one end of a diameter of a circle is (3, 4) and the co - ordinates of its centre is (-3, 2), then the coordinate of the other end of the diameter is

- (a) (0, -3) (b) (0, 9) (c) (3, 0) (d) (-9, 0)

Ans: (d) (-9, 0)

$$\text{Mid point of a line} = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right) \Rightarrow (-3, 2) = \left(\frac{3+a}{2}, \frac{4+b}{2} \right)$$

$$\frac{3+a}{2} = -3 \Rightarrow 3 + a = -6 \Rightarrow a = -9$$

$$\frac{4+b}{2} = 2 \Rightarrow 4 + b = 4 \Rightarrow b = 0$$

15. The ratio in which the x-axis divides the line segment joining the points A (a_1 , b_1) and B (a_2 , b_2) is

(a) $b_1 : b_2$ (b) $-b_1 : b_2$ (c) $a_1 : a_2$ (d) $-a_1 : a_2$

Ans: (b) - $b_1 : b_2$

$$\begin{aligned} \text{The point P is } (a, 0) &= \left(\frac{ma_2 + na_1}{m+n}, \frac{mb_2 + nb_1}{m+n} \right) \\ \therefore \frac{mb_2 + nb_1}{m+n} &= 0 \Rightarrow mb_2 + nb_1 = 0 \Rightarrow mb_2 = -nb_1 \\ \therefore m : n &= -b_1 : b_2 \end{aligned}$$

16. The ratio in which the x-axis divides the line segment joining the points (6, 4) and (1, -7) is

(a) 2 : 3 (b) 3 : 4 (c) 4 : 7 (d) 4 : 3

Ans: (c) 4 : 7

$$\begin{aligned} P &= \left(\frac{m+6n}{m+n}, \frac{-7m+4n}{m+n} \right) \Rightarrow (a, 0) = \left(\frac{m+6n}{m+n}, \frac{-7m+4n}{m+n} \right) \\ 0 &= \frac{-7m+4n}{m+n} \Rightarrow -7m+4n=0 \Rightarrow 4n=7m \Rightarrow \text{The ratio is } 4 : 7 \end{aligned}$$

17. The mid-point of the line joining (-a, 2b) and (-3a, -4b) is

(a) (2a, 3b) (b) (-2a, -b) (c) (2a, b) (d) (-2a, -3b)

Ans: (b) (-2a, -b)

$$\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right) = \left(\frac{-a - 3a}{2}, \frac{2b - 4b}{2} \right) = \left(\frac{-4a}{2}, \frac{-2b}{2} \right) = (-2a, -b)$$

18. In what ratio does the y-axis divide the line joining the points (-5, 1) and (2, 3) internally

(a) 1 : 3 (b) 2 : 5 (c) 3 : 1 (d) 5 : 2

Ans: (d) 5 : 2

$$\begin{aligned} P &= \left(\frac{mx_2 + nx_1}{m+n}, \frac{my_2 + ny_1}{m+n} \right) \Rightarrow (0, a) = \left(\frac{2m-5n}{m+n}, \frac{3m+n}{m+n} \right) \\ \frac{2m-5n}{m+n} &= 0 \quad 2m-5n=0 \Rightarrow 2m=5n \Rightarrow m : n = 5 : 2 \end{aligned}$$

19. If (1, -2), (3, 6), (x, 10) and (3, 2) are the vertices of the parallelogram taken in order, then the value of x is

(a) 6 (b) 5 (c) 4 (d) 3

Ans: (b) 5

$$\begin{aligned} \text{Mid-point of AC} &= \text{Mid-point of BD} \\ \left(\frac{1+x}{2}, \frac{-2+10}{2} \right) &= \left(\frac{3+3}{2}, \frac{6+2}{2} \right) \\ \frac{1+x}{2} &= \frac{6}{2} \Rightarrow 1+x=6 \Rightarrow x=6-1=5 \Rightarrow x=5 \end{aligned}$$

20. On which quadrant does the point $(-4, 3)$ lie?
 (a) I (b) II (c) III (d) IV
Ans: (b) II
21. The point whose abscissa is 5 and lies on the x-axis is
 (a) $(-5, 0)$ (b) $(5, 5)$ (c) $(0, 5)$ (d) $(5, 0)$
Ans: (d) $(5, 0)$
22. A point which lies in the III quadrant is
 (a) $(5, 4)$ (b) $(5, -4)$ (c) $(-5, -4)$ (d) $(-5, 4)$
Ans: (c) $(-5, -4)$
23. A point on the y-axis is
 (a) $(1, 1)$ (b) $(6, 0)$ (c) $(0, 6)$ (d) $(-1, -1)$
Ans: (c) $(0, 6)$
24. The distance between the points $(4, -1)$ and the origin is
 (a) $\sqrt{24}$ (b) $\sqrt{37}$ (c) $\sqrt{26}$ (d) $\sqrt{17}$
Ans: (d) $\sqrt{17}$
25. The distance between the points $(-1, 2)$ and $(3, 2)$ is
 (a) $\sqrt{14}$ (b) $\sqrt{15}$ (c) 4 (d) 0
Ans: (c) 4
26. The centre of a circle is $(0, 0)$. One end point of a diameter is $(5, -1)$, then the radius is
 (a) $\sqrt{24}$ (b) $\sqrt{37}$ (c) $\sqrt{26}$ (d) $\sqrt{17}$
Ans: (c) $\sqrt{26}$
27. The point $(0, -3)$ lies on
 (a) +ve x - axis (b) +ve y - axis (c) -ve x - axis (d) -ve y - axis
Ans: (d) -ve y - axis
28. The point which is on y - axis with ordinate -5 is
 (a) $(0, -5)$ (b) $(-5, 0)$ (c) $(5, 0)$ (d) $(0, 5)$
Ans: (a) $(0, -5)$
29. The diagonal of a square formed by the points $(1, 0)$, $(0, 1)$, $(-1, 0)$ and $(0, -1)$ is
 (a) 2 (b) 4 (c) $\sqrt{2}$ (d) 8
Ans: (a) 2
30. The distance between the points $(-2, 2)$ and $(3, 2)$ is
 (a) 10 units (b) 5 units (c) $5\sqrt{3}$ units (d) 20 units
Ans: (b) 5 units
31. The midpoint of the line joining the points $(1, -1)$ and $(-5, 3)$ is
 (a) $(2, 1)$ (b) $(2, -1)$ (c) $(-2, -1)$ (d) $(-2, 1)$
Ans: (d) $(-2, 1)$

32.If the centroid of a triangle is at (1, 3) and two of its vertices are (-7, 6) and (8, 5) then the third vertex is

- (a) (-2, 2) (b) (2, -2) (c) (-2, -2) (d) (2, 2)

Ans: (b) (2, -2)

33.The ratio in which the X-axis divides the line segment joining the points (6, 4) and (1, -7) is

- (a) 1 : 2 (b) 2 : 3 (c) 4 : 7 (d) 7 : 4

Ans: (c) 4 : 7

34.The centroid of a triangle (3, -5), (-7, 4) and (10, -2) is

- (a) (2, -1) (b) (2, 1) (c) (-2, 1) (d) (1, -2)

Ans: (a) (2, -1)

35.The name of the horizontal line in the Cartesian plane which determines the position of a point is called:

- a. Origin b. X - axis c. Y- axis d. Quadrants

Ans: b. X - axis

36.The name of the vertical line in the Cartesian plane which determines the position of a point is called:

- a. Origin b. X – axis c. Y – axis d. Quadrants

Ans: c. Y – axis

37.The section formed by horizontal and vertical lines determining the position of the point in a Cartesian plane is called:

- (a) Origin (b) X – axis (c) Y – axis (d) Quadrants

Ans: (d) Quadrants

38.The point of intersection of horizontal and vertical lines determining the position of a point in a Cartesian plane is called:

- (a) Origin (b) X – axis (c) Y – axis (d) Quadrants

Ans: (a) Origin

39.If the coordinates of a point are (0, -4), then it lies in:

- (a) X – axis (b) Y – axis (c) At origin (d) Between x – axis & y – axis

Ans: Y – axis

40.If the coordinates of a point are (3, 0), then it lies in:

- (a) X – axis (b) Y – axis (c) At origin (d) Between x – axis & y – axis

Ans: (a) X – axis

41.If the coordinates of a point are (-3, 4), then it lies in:

- (a) First quadrant (b) Second quadrant
(c) Third quadrant (d) Fourth quadrant

Ans: (b) Second quadrant

42.If the coordinates of a point are (-3, -4), then it lies in_____ quadrant.

- (a) First (b) Second (c) Third (d) Fourth

Ans: (c) Third

43. Points (1, 2), (-2, -3), (2, -3);

- (a) First quadrant (b) Do not lie in the same quadrant
(c) Third quadrant (d) Fourth quadrant

Ans: (b) Do not lie in the same quadrant

44. If x coordinate of a point is zero, then the point lies on:

- (a) First quadrant (b) Second quadrant (c) X – axis (d) Y – axis

Ans: (d) Y – axis

45. Signs of the abscissa and ordinate of a point in the second quadrant are respectively

- (a) +, + (b) +, – (c) –, + (d) –, –

Ans: (c) –, +

46. The point (-10, 0) lies in

- (a) Third quadrant (b) On the negative direction of the x – axis
(c) Fourth quadrant (d) On the negative direction of the y – axis

Ans: (b) On the negative direction of the x – axis

47. A quadrant in which both x and y values are negative is

- (a) First quadrant (b) Second quadrant
(c) Third quadrant (d) Fourth quadrant

Ans: (c) Third quadrant

48. Abscissa of all the points on the x – axis is

- (a) 0 (b) 1 (c) 2 (d) Any number

Ans: (d) Any number

49. Ordinate of all points on the x-axis is

- (a) -1 (b) 0 (c) 1 (d) Any number

Ans: (b) 0

50. Abscissa of a point is positive in

- (a) I quadrant (b) I and II quadrants
(c) II quadrant only (d) I and IV quadrants

Ans: (d) I and IV quadrants

51. Points (1, -1), (2, -2), (4, -5), (-3, -4)

- (a) lie in II quadrant (b) lie in III quadrant
(c) lie in IV quadrant (d) Does not lie in the same quadrant

Ans: (d) Doesn't lie in the same quadrant

52. Abscissa of all the points on the y – axis is

- (a) 0 (b) 1 (c) -1 (d) Any number

Ans: (a) 0

53. Ordinate of all the points on the y – axis is

- (a) 0 (b) 1 (c) -1 (d) Any number

Ans: (d) Any number

54. The point which lies on the y – axis at a distance of 5 units in the negative direction of the y – axis is

- (a) (5, 0) (b) (0, 5) (c) (-5, 0) (d) (0, -5)

Ans: (d) (0, -5)

55. The linear equation $3x - 11y = 10$ has:

- (a) Unique solution (b) Two solutions
(c) Infinitely many solutions (d) No solutions

Ans: (c) Infinitely many solutions

56. $3x + 10 = 0$ will have:

- (a) Unique solution (b) Two solutions
(c) Infinitely many solutions (d) No solutions

Ans: (a) Unique solution

57. The solution of equation $x - 2y = 4$ is:

- (a) (0, 2) (b) (2, 0) (c) (4, 0) (d) (1, 1)

Ans: (c) (4, 0)

58. Find the value of k , if $x = 1$, $y = 2$ is a solution of the equation $2x + 3y = k$.

- (a) 5 (b) 6 (c) 7 (d) 8

Ans: (d) 8

59. Point (3, 4) lies on the graph of the equation $3y = kx + 7$. The value of k is:

- (a) $4/3$ (b) $5/3$ (c) 3 (d) $7/3$

Ans: (b) $5/3$

$$3y = kx + 7 \Rightarrow (3 \times 4) = (k \times 3) + 7 \Rightarrow 12 = 3k + 7 \Rightarrow 3k = 12 - 7 \Rightarrow k = 5/3$$

60. The graph of linear equation $x + 2y = 2$, cuts the y – axis at:

- (a) (2, 0) (b) (0, 2) (c) (0, 1) (d) (1, 1)

Ans: (c) (0, 1)

$$x + 2y = 2 \Rightarrow y = (2 - x) / 2 \Rightarrow y = (2 - 0) / 2 = 2 / 2 = 1$$

61. Any point on line $x = y$ is of the form:

- (a) (k, -k) (b) (0, k) (c) (k, 0) (d) (k, k)

Ans: (d) (k, k)

62. The graph of $x = 3$ is a line:

- (a) Parallel to the x -axis at a distance of 3 units from the origin
(b) Parallel to the y -axis at a distance of 3 units from the origin
(c) Makes an intercept 3 on the x -axis
(d) Makes an intercept 3 on the y -axis

Ans: (b) Parallel to the y -axis at a distance of 3 units from the origin

63. In equation, $y = mx + c$, m is:

- (a) Intercept (b) Slope (c) Solution of the equation (d) None of the above

Ans: (b) Slope

64.If x and y are both positive solutions of equation $ax + by + c = 0$, always lie in the:

- (a) First quadrant
- (b) Second quadrant
- (c) Third quadrant
- (d) Fourth quadrant

Ans: (a) First quadrant

65.A linear equation in two variables is of the form $ax + by + c = 0$, where

- (a) $a = 0, c = 0$
- (b) $a \neq 0, b = 0$
- (c) $a = 0, b \neq 0$
- (d) $a \neq 0, b \neq 0$

Ans: (d) $a \neq 0, b \neq 0$

66.Any point on the x -axis is of the form

- (a) (x, y)
- (b) $(0, y)$
- (c) $(x, 0)$
- (d) (x, x)

Ans: (c) $(x, 0)$

67.Any point on the y -axis is of the form

- (a) (y, y)
- (b) $(0, y)$
- (c) (x, y)
- (d) $(x, 0)$

Ans: (b) $(0, y)$

68.The linear equation $2x - 5y = 7$ has

- (a) No solution
- (b) Unique solution
- (c) Two solutions
- (d) Infinitely many solutions

Ans: (d) Infinitely many solutions

69.The linear equation $3x - y = x - 1$ has

- (a) No solution
- (b) Unique solution
- (c) Two solutions
- (d) Infinitely many solutions

Ans: (d) Infinitely many solutions

70.The graph of the linear equation $2x + 3y = 6$ cuts the y – axis at the point

- (a) $(2, 0)$
- (b) $(0, 2)$
- (c) $(3, 0)$
- (d) $(0, 3)$

Ans: (b) $(0, 2)$

71.The equation $2x + 5y = 7$ has a unique solution, if x, y are:

- (a) Rational numbers
- (b) Real numbers
- (c) Natural numbers
- (d) Positive real numbers

Ans: (c) Natural numbers

72.The point of the form (a, a) always lies on:

- (a) On the line $x + y = 0$
- (b) On the line $y = x$
- (c) x – axis
- (d) y – axis

Ans: (b) On the line $y = x$

73.If we multiply or divide both sides of a linear equation with the same non-zero number, then the solution of the linear equation:

- (a) Remains the same
- (b) Changes
- (c) Changes in case of multiplication only
- (d) Changes in case of division only

Ans: (a) Remains the same

74.If (2, 0) is a solution of the linear equation $2x + 3y = k$, then the value of k is
(a) 2 (b) 4 (c) 5 (d) 6

Ans: (b) 4

75.Which graph is parallel to x – axis?

(a) $y = x + 1$ (b) $y = 2$ (c) $x = 3$ (d) $x = 2y$

Ans: (b) $y = 2$

76.Which point lies on x – axis?

(a) (3, 2) (b) (-3, 2) (c) (2, 0) (d) (-1,-2)

Ans: (c) (2, 0)

77.Which point lies on y – axis?

(a) (1, 3) (b) (0, 3) (c) (5, 2) (d) (-2,-3)

Ans: (b) (0, 3)

78.Which point lies to the right of y – axis?

(a) (0, 3) (b) (-2,-1) (c) (3, 5) (d) (-3,-2)

Ans: (c) (3, 5)

79.Which line is parallel to $y = x - 2$?

(a) $y = 2x + 1$ (b) $y = 2x - 6$ (c) $2y = x + 7$ (d) $y = 3x + 1$

Ans: (b) $y = 2x - 6$

80.Which point lies on the left of y – axis?

(a) (2, 0) (b) (-2,-4) (c) (5, 2) (d) (3, 6)

Ans: (b) (-2, -4)

81.Which point lies in IV quadrant?

(a) (-3,-4) (b) (2,-4) (c) (-2, 3) (d) (0, 1)

Ans: (b) (2, -4)

82.Which point lies above x – axis?

(a) (-1, 2) (b) (2, 0) (c) (-1,-5) (d) (0,-3)

Ans: (a) (-1, 2)

83.The distance between the point (0,5) and (-5,0) is:

(A) 5 (B) $5\sqrt{2}$ (C) $2\sqrt{3}$ (D) 2

Ans: (B) $5\sqrt{2}$

Using distance formula $d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$

$$d = \sqrt{(-5 - 0)^2 + (0 - 5)^2} = \sqrt{(-5)^2 + (-5)^2} = \sqrt{50} = 5\sqrt{2}$$

84.The distance between the point P(3,2) and Q(-2,-3) is:

(A) 7.09 (B) 7.07 (C) 7.21 (D) 1.41

Ans: (B) 7.07

$$d = \sqrt{(-2 - 3)^2 + (-3 - 2)^2} = \sqrt{(-5)^2 + (-5)^2} = \sqrt{50} \approx 7.07$$

85. The distance between the points (2,3) and (4,1) is:
 (A) $\sqrt{8}$ (B) $3\sqrt{2}$ (C) $2\sqrt{2}$ (D) $5\sqrt{3}$

Ans: (C) $2\sqrt{2}$

$$d = \sqrt{(4-2)^2 + (1-3)^2} = \sqrt{(2)^2 + (-2)^2} = \sqrt{8} = 2\sqrt{2}$$

86. Point's (-5,-4) is situated in which quadrant:
 (A) I (B) II (C) III (D) IV

Ans: (C) III

87. Distance between points (0, 4) and (3, 0) is _____.
 (a) 6 (b) 5 (c) 4 (d) 7

Ans: (b) 5

$$d = \sqrt{(3-0)^2 + (0-4)^2} = \sqrt{9+16} = \sqrt{25} = 5$$

88. Distance of any point from x-axis is called _____.
 (a) Origin (b) Abscissa (c) Ordinate (d) None of these

Ans: (c) Ordinate

89. Distance between (0, 0) and (x, y) is _____.
 (a) $x^2 + y^2$ (b) $x^2 - y^2$ (c) $\sqrt{x^2 + y^2}$ (d) $\sqrt{x^2 - y^2}$

Ans: (c) $\sqrt{x^2 + y^2}$

90. If the perpendicular distance of a point P from the x-axis is 7 units and the foot of the perpendicular lies on the negative direction of the x-axis, then the point p will have
 (a) Y-coordinate = 7 or -7 only (b) Y- coordinate = 7 only
 (c) Y – coordinate = -7 only (d) X- coordinate = -7

Ans. (a) Y-coordinate = 7 or -7 only

91. Given the coordinates of two points P and Q as (-7, 5) and (-6, 9) respectively. What will be the value of (abscissa of P) – (abscissa of Q)?
 (a) -3 (b) 1 (c) -2 (d) -1

Ans. (d) -1

92. In which quadrant will the points Q (1, 2), R (-2, -3), and S (2, -3) lie?
 (a) I quadrant (b) IV quadrant
 (c) III quadrant (d) Don't lie in the same quadrant

Ans. (d) Don't lie in the same quadrant

93. Name the sections formed by the intersection of horizontal and vertical lines determining the location of the point in a Cartesian plane:
 (a) Origin (b) X – axis (c) Y – axis (d) Quadrants

Ans. (d) Quadrants

94. Determine in which quadrant the given points (-5,2) and (2,-5) will lie in:
 (a) II and III quadrants respectively (b) I and IV quadrants respectively
 (c) II and IV quadrants respectively (d) In the same quadrants

Ans. (c) II and IV quadrants respectively

95. A point M lies in the IV quadrant. What will be the coordinate of point M?

- (a) (a, b) (b) (-a, -b) (c) (-a, b) (d) (a, -b)

Ans: (d) (a, -b)

96. Perpendicular distance of the point P(-3, 8) from y-axis is

- (a) -3 (b) 8 (c) 3 (d) -8

Ans: (c) 3

97. If points P(5, 1), Q(0, 2), R(3, 0), S(-1, 0) are plotted on a graph paper, then the points on y-axis are

- (a) only P (b) only Q (c) Q and R (d) R and S

Ans: (b) Only Q

98. If P(-1, 1), Q(3, -4), R(1, -1), S(-2, -3) and T(-4, 4) are plotted on the graph paper, then the point(s) in the fourth quadrant are

- (a) P and T (b) Q and R (c) Only S (d) P and R

Ans: (b) Q and R

99. If the coordinates of the two points are P(-5, 3) and Q(8, -9), then (abscissa of Q) - (abscissa of P) is

- (a) 4 (b) -12 (c) 13 (d) -13

Ans: (c) 13

$(\text{abscissa of Q}) - (\text{abscissa of P}) = 8 - (-5) = 8 + 5 = 13$

100. Perpendicular distance of the point P(-3, 8) from y-axis is

- (a) -3 (b) 8 (c) 3 (d) -8

Ans: (c) 3