

Statistics (9 th std)

Choose the Appropriate Answer

1. Let 'm' be the mid-point and b be the upper limit of a class in a continuous frequency distribution. The lower limit of the class is

- (a) $2m - b$ (b) $2m + b$ (c) $m - b$ (d) $m - 2b$

Ans: (a) $2m - b$

2. The mean of a set of seven numbers is 81. If one of the numbers is discarded, the mean of the remaining numbers is 78. The value of discarded number is _____.

- (a) 101 (b) 100 (c) 99 (d) 98

Ans: (c) 99

Total of 8 numbers = $81 \times 7 = 567$ $\Rightarrow$ Total of 7 numbers = $78 \times 6 = 468$ $\Rightarrow$ Number = $567 - 468 = 99$

3. A particular observation which occurs maximum number of times in a given data is called its

- (a) Frequency (b) Range (c) Mode (d) Median

Ans: (c) Mode

4. For which set of numbers do the mean, median and mode all have the same values?

- (a) 2, 2, 2, 4 (b) 1, 3, 3, 3, 5 (c) 1, 1, 2, 5, 6 (d) 1, 1, 2, 1, 5

Ans: (b) 1, 3, 3, 3, 5

5. The algebraic sum of the deviations of a set of n values from their mean is ____.

- (a) 0 (b) $n - 1$ (c) n (d) $n + 1$

Ans: (a) 0

6. The mean of a, b, c, d and e is 28. If the mean of a, c and e is 24, then mean of b and d is

- (a) 24 (b) 36 (c) 26 (d) 34

Ans: (d) 34

Mean = 28 $\Rightarrow a + b + c + d + e = 28 \times 5 = 140$ (a + c + e) / 3 = 24 $\Rightarrow a + c + e = 72 \Rightarrow b + d + 72 = 140$ $b + d = 140 - 72 = 68 \Rightarrow$ Mean = $68 / 2 = 34$

7. If the mean of five observations x, x + 2, x + 4, x + 6, x + 8, is 11, then the mean of first three observations is

- (a) 9 (b) 11 (c) 13 (d) 15

Ans: (a) 9

Mean = $x + x + 2 + x + 4 + x + 6 + x + 8$ $\Rightarrow 11 = (5x + 20) / 5 \Rightarrow 5x + 20 = 55 \Rightarrow x = 7$ Mean of first 3 observation is $= (7 + 9 + 11) / 3 = 9$

8. The mean of 5, 9, x, 17 and 21 is 13, then find the value of x.

- (a) 9 (b) 13 (c) 17 (d) 21

Ans: (b) 13

$$\text{Mean} = (5 + 9 + x + 17 + 21) / 5 \Rightarrow 13 = (52 + x) / 5 \Rightarrow 65 = 52 + x \Rightarrow x = 13$$

9. The mean of the square of first 11 natural numbers is

- (a) 26 (b) 46 (c) 48 (d) 52

Ans: (b) 46

$$\begin{aligned} \text{Mean} &= \frac{1^2 + 2^2 + 3^2 + 4^2 + 5^2 + 6^2 + 7^2 + 8^2 + 9^2 + 10^2 + 11^2}{11} \\ &= \frac{1 + 4 + 9 + 16 + 25 + 36 + 49 + 64 + 81 + 100 + 121}{11} = \frac{506}{11} = 46 \end{aligned}$$

10. The mean of a set of numbers is $\bar{X}$. If each number is multiplied by z, the mean is _____

- (a) $\bar{X} + z$ (b) $\bar{X} - z$ (c) $z\bar{X}$ (d) $\bar{X}$

Ans: (c) $z\bar{X}$

11. The Arithmetic mean of all the factors of 10 is

- (a) 4.5 (b) 5.5 (c) 10 (d) 55

Ans: (a) 4.5

12. The mean of five numbers is 27, if one number is excluded, then mean is 25. Then the excluded number is

- (a) 0 (b) 15 (c) 25 (d) 35

Ans: (d) 35

13. The mean of 8 numbers is 15. If each number is multiplied by 2, then the new mean will be

- (a) 7.5 (b) 30 (c) 10 (d) 25

Ans: (b) 30

14. The median of 11, 8, 4, 9, 7, 5, 2, 4, & 10 is

- (a) 1 (b) 8 (c) 4 (d) 11

Ans: (a) 1

15. Median is

- (a) The most frequent value (b) The least frequent value
(c) Middle most value (d) Mean of first and last values

Ans: (c) Middle most value

16. The mode of the distribution is

x	1	2	3	4	5	6
f	2	3	7	14	8	10

- (a) 3 (b) 4 (c) 6 (d) 14

Ans: (b) 4

17. Mode is

- (a) The middle value (b) Extreme value
(c) Minimum value (d) The most repeated value

Ans: (d) The most repeated value

18. The mode of the data 72, 33, 44, 72, 81, 72, 15 is

- (a) 12 (b) 33 (c) 81 (d) 15

Ans: (a) 12

19. The Arithmetic mean of 10 number is -7. If 5 is added to every number, then the new arithmetic mean is

- (a) 17 (b) 12 (c) -2 (d) -7

Ans: (c) -2

20. The Arithmetic mean of integers from -5 to 5 is

- (a) 25 (b) 10 (c) 3 (d) 0

Ans: (d) 0

21. The ratio of the sum of observations and the total number of observations is called:

- (a) Mean (b) Median (c) Mode (d) Central tendency

Ans: (a) Mean

22. The mean of $x + 2$, $x + 3$, $x + 4$ and $x - 2$ is:

- (a) $(x + 7) / 4$ (b) $(2x + 7) / 4$ (c) $(3x + 7) / 4$ (d) $(4x + 7) / 4$

Ans: (d) $(4x + 7) / 4$

$$\text{Mean} = (x + 2 + x + 3 + x + 4 + x - 2) / 4 = (4x + 7) / 4$$

23. The median of the data: 4, 6, 8, 9, 11 is

- (a) 6 (b) 8 (c) 9 (d) 11

Ans: (b) 8

24. The median of the data: 155, 160, 145, 149, 150, 147, 152, 144, 148 is

- (a) 149 (b) 150 (c) 147 (d) 144

Ans: (a) 149

Arrange the data in ascending order. 144 145 147 148 149 150 152 155 160
Median = $(n + 1) / 2$ th = $(9 + 1) / 2 = 10 / 2 = 5$ th number = 149

25. Median of the data: 17, 2, 7, 27, 15, 5, 14, 8, 10, 24, 48, 10, 8, 7, 18, 28 is:

- (a) 10 (b) 24 (c) 12 (d) 8

Ans: (c) 12

Arrange the given data in ascending order:
2, 5, 7, 7, 8, 8, 10, 10, 14, 15, 17, 18, 24, 27, 28, 48
 $n / 2$ th = $16 / 2 = 8$ th term $(n / 2 + 1)$ th = $(16 / 2 + 1)$ th = 9th term
Median = $(10 + 14) / 2 = 12$

26. The value which appears very frequently in a data is called:
(a) Mean (b) Median (c) Mode (d) Central tendency

Ans: (c) Mode

27. The collection of information, collected for a purpose is called:
(a) Mean (b) Median (c) Mode (d) Data

Ans: (d) Data

28. The mean of the data 2, 3, 4, 5, 0, 1, 3, 3, 4, 3 is
(a) 2 (b) 2.2 (c) 2.4 (d) 2.8

Ans: (d) 2.8

$$\text{Mean} = (2+3+4+5+0+1+3+3+4+3) / 10 = 28 / 10 = 2.8$$

29. Which of the following is not a measure of central tendency?
(a) Standard deviation (b) Mean (c) Median (d) Mode

Ans: (a) Standard deviation

30. Find the range of the following data: 25, 18, 20, 22, 16, 6, 17, 15, 12, 30, 32, 10, 19, 8, 11, & 20.
(a) 10 (b) 15 (c) 18 (d) 26

Ans: (d) 26

$$\text{Range} = \text{Maximum value} - \text{Minimum value} \Rightarrow \text{Range} = 32 - 6 = 26.$$

31. What is the class mark of the class interval 90 - 120?
(a) 90 (b) 105 (c) 115 (d) 120

Ans: (b) 105

$$\text{Class mark} = (\text{Upper limit} + \text{Lower limit}) / 2 = (120+90) / 2 = 105$$

32. In the class intervals 10 - 20, 20 - 30, 20 is included in which interval?
(a) 10 - 20 (b) 20 - 30 (c) Both the intervals (d) None of these

Ans: (b) 20 - 30

33. Find the class width for the grouped frequency distribution of the class intervals 1 - 20, 21 - 40, 41 - 60
(a) 10 (b) 15 (c) 17 (d) 20

Ans: (d) 20

34. The arithmetic mean of the first 5 natural numbers is
(a) 3 (b) 4 (c) 5 (d) 6

Ans: (a) 3

$$\text{Arithmetic mean} = (1+2+3+4+5) / 5 = 15 / 5 = 3$$

35. Find the value of x, if the arithmetic mean of 4, 5, 6, 7, 8 and x is 7.
(a) 4 (b) 6 (c) 8 (d) 12

Ans: (d) 12

$$(4+5+6+7+8+x) / 6 = 7 \Rightarrow 4+5+6+7+8+x = 42 \Rightarrow 30+x = 42 \Rightarrow x = 42-30 = 12$$

36. Find the mode of the following data: 15, 14, 19, 20, 14, 15, 16, 14, 15, 18, 14, 19, 15, 17, & 15.

- (a) 14 (b) 15 (c) 16 (d) 17

Ans: (b) 15

37. If each data in the observation is increased by 10, then the mean

- (a) Remains the same (b) Increased by 10
(c) Decreased by 5 (d) None of the above

Ans: (b) Increased by 10

38. The difference between the maximum and minimum values of the given observation is called

- (a) Class (b) Class interval (c) Class mark (d) Range

Ans: (d) Range

39. Find the maximum value if the range is 38 and the minimum value is 82.

- (a) 60 (b) 76 (c) 120 (d) 82

Ans: (c) 120

$$\text{Range} = \text{Maximum value} - \text{Minimum value.} \\ 38 = x - 82 \Rightarrow x = 38 + 82 \Rightarrow x = 120.$$

40. From the table given below, the marks of which the highest frequency exists is _____

Marks	3	4	5	6	7	8	9	10
Students	1	2	2	4	5	3	2	1

- (a) 5 (b) 7 (c) 10 (d) 20

Ans: (b) 7

41. Consider the runs scored by 11 players in a match which are given below.

What is the range of this given data? 8 22 64 16 32 29 16 2 7 15 1

- (a) 63 (b) 64 (c) 1 (d) 22

Ans: (a) 63

$$\text{Range} = 64 - 1 = 63.$$

42. From the table given below, what is the upper limit of class 50-55?

Mass	40-45	45-50	50-55	55-60	60-65	65-70
Students	4	12	3	2	3	1

- (a) 25 (b) 52.5 (c) 55 (d) 60

Ans: (c) 55

43. From the table given below, what is the lower limit of class 65 – 70?

Mass	40-45	45-50	50-55	55-60	60-65	65-70
Students	4	12	3	2	3	1

- (a) 25 (b) 67.5 (c) 70 (d) 65

Ans: (d) 65

44. If the mean of frequency distribution is 6.5 and $\sum f_i x_i = 120 + 2k$, $\sum f_i = 20$, then k is equal to:

- (a) 10 (b) 15 (c) 5 (d) 25

Ans: (c) 5

$$X_{\text{mean}} = \sum f_i x_i / \sum f_i \Rightarrow 6.5 = (120 + 2k) / 20 \Rightarrow 130 = 120 + 2k \Rightarrow 2k = 10 \Rightarrow k = 5$$

45. If $x_1, x_2, x_3, \dots, x_n$ are the observations of a given data. Then the mean of the observations will be:

- (a) Total number of observations / Sum of observations
 (b) Sum of observations / Total number of observations
 (c) Sum of observations + Total number of observations
 (d) None of the above

Ans: (b) Sum of observations / Total number of observations

46. The mode and mean are given by 5 and 6, respectively. Then the median is:

- (a) 17 / 3 (b) 13 / 3 (c) 23 / 3 (d) 33

Ans: (a) 17 / 3

$$\text{Mode} = 3 \text{ Median} - 2 \text{ Mean} \Rightarrow \text{Median} = (\text{Mode} + 2 \text{ Mean}) / 3 \\ \text{Median} = (5 + 2(6)) / 3 = (5 + 12) / 3 = 17 / 3$$

47. The mean of the data: 4, 11, 5, 8, 12 is;

- (a) 16 (b) 8 (c) 9 (d) 15

Ans: (b) 8

$$\text{Mean} = (4 + 11 + 5 + 8 + 12) / 5 = 40 / 5 = 8$$

48. The median of the data 11, 12, 14, 17, 20, 27 is:

- (a) 30 / 2 (b) 31 / 2 (c) 33 / 2 (d) 35 / 2

Ans: (b) 31 / 2

$$\text{Median} = (14 + 17) / 2 = 31 / 2$$

49. Find the mean of frequency distribution, if $\sum f_i x_i = 110$, and $\sum f_i = 10$, then k is equal to:

- (a) 5 (b) 23 (c) 15 (d) 11

Ans: (d) 11

$$X_{\text{mean}} = \sum f_i x_i / \sum f_i = (110) / 10 = 11$$

50. The mean and median are given by 4 and 12, respectively. Then the mode is:
 (a) 28 (b) 32 (c) 23 (d) 45

Ans: (a) 28

$$\text{Mode} = 3 \text{ Median} - 2 \text{ Mean} = (3 \times 12) - 2 \times 4 = 36 - 8 = 28$$

51. If the mean of first n natural numbers is $4n / 6$, then the value of n is:
 (a) 4 (b) 2 (c) 6 (d) 3

Ans: (d) 3

$$\begin{aligned} \text{Sum of natural numbers} &= n(n+1)/2 \Rightarrow \text{Mean} = 4n/6 \\ \text{Mean} &= \text{Sum of natural numbers} / n \\ 4n/6 &= n(n+1)/2n \Rightarrow 4n/6 = (n+1)/2 \Rightarrow 8n = 6n + 6 \Rightarrow n = 3 \end{aligned}$$

52. If the mean of a , $a+2$, $a+4$, $a+8$ and $a+11$ is 10, then a is equal to;
 (a) 1 (b) 2 (c) 5 (d) 4

Ans: (c) 5

$$(a + a + 2 + a + 4 + a + 8 + a + 11) / 5 = 10 \Rightarrow 5a + 25 = 50 \Rightarrow a = 5$$

53. The class interval of a given observation is 5 to 10, then the class mark for this interval will be:

(a) 7.5 (b) 13.5 (c) 8 (d) 14.5

Ans: (a) 7.5

$$\text{Class mark} = (\text{Upper limit} + \text{Lower limit}) / 2 = (10 + 5) / 2 = 15 / 2 = 7.5$$

54. If the sum of frequencies is 30, then the value of x in the observation: x , 5, 6, 1, 2, will be;

(a) 4 (b) 16 (c) 9 (d) 10

Ans: (b) 16

$$\sum f_i = x + 5 + 6 + 1 + 2 = 14 + x \Rightarrow 30 = 14 + x \Rightarrow x = 30 - 14 = 16$$

55. The class marks of a frequency distribution are given as follows: 15, 20, 25...
 The class corresponding to the class mark 20 is

(a) 12.5 – 17.5 (b) 17.5 – 22.5 (c) 18.5 – 21.5 (d) 19.5 – 20.5

Ans: (b) 17.5 – 22.5

56. In the class intervals 10–20 and 20–30, the number 20 is included in
 (a) 10 – 20 (b) 20 – 30 (c) Both the intervals (d) None of these

Ans: (b) 20 – 30

57. The class mark of the class 90 – 130 is

(a) 90 (b) 110 (c) 115 (d) 120

Ans: (b) 110

$$\text{Class mark} = (90 + 130) / 2 = 110$$

58. In a frequency distribution, the mid value of a class is 10 and the width of the class is 6. The lower limit of the class is
 (a) 6 (b) 7 (c) 8 (d) 12

Ans: (b) 7

$$\text{The lower limit} = \text{Mid value} - (\text{Class size} / 2) = 10 - (6 / 2) = 10 - 3 = 7.$$

59. To draw a histogram to represent the following frequency distribution:

Class Interval	5 - 10	10 - 15	15 - 25	25 - 45	45 - 75
Frequency	6	12	10	8	15

The adjusted frequency for the class 25–45 is

- (a) 6 (b) 5 (c) 3 (d) 2

Ans: (d) 2

$$\begin{aligned} \text{Class size of CI} &= 25 - 45 = 20 \\ \text{Adjusted frequency for class interval for } 25 - 45 &= (5 / 20) \times 8 = 2. \end{aligned}$$

60. For drawing a frequency polygon of a continuous frequency distribution, we plot the points whose ordinates are the frequencies of the respective classes and abscissa are respectively:

- (a) Upper limits of the classes (b) Lower limits of the classes
 (c) Class marks of the classes (d) Upper limits of preceding classes

Ans: (c) Class marks of the classes

61. In a frequency distribution, the mid value of a class is 10 and the width of the class is 6. The upper limit of the class is:

- (a) 10 (b) 7 (c) 8 (d) 13

Ans: (d) 13

$$\begin{aligned} \text{Lower limit} &= \text{Mid value} - (\text{Class width} / 2) = 10 - (6 / 2) = 10 - 3 = 7 \\ \text{Class width} &= \text{Upper limit} - \text{Lower limit} \Rightarrow 6 = \text{Upper limit} - 7 \\ \text{Upper limit} &= 6 + 7 = 13 \end{aligned}$$

62. The width of each of five continuous classes in a frequency distribution is 5 and the lower class-limit of the lowest class is 10. The upper class - limit of the highest class is:

- (a) 15 (b) 25 (c) 35 (d) 40

Ans: (c) 35

$$\text{The upper class limit of the highest class} = 10 + 25 = 35$$

63. Find the mean of $x + 77$, $x + 7$, $x + 5$, $x + 3$ and $x - 2$?

- (a) $x + 8$ (b) $x + 18$ (c) $x - 8$ (d) $x - 18$

Ans: (b) $x + 18$

64. If the mean of n observations $x_1, x_2, x_3, \dots, x_n$ is $\bar{x}$ then what is $\sum_{i=1}^n (x_i - \bar{x})$?

- (a) 0 (b) 1 (c) 2 (d) 3

Ans: (a) 0

65.If each observation of the data is increased by 5, then what happens to its mean?

- (a) Is increased by 4 (b) Is increased by 5
(c) Is decreased by 4 (d) Is decreased by 5

Ans: (b) Is increased by 5

66.In the class intervals 40 – 50, 50 – 60, the number 50 is included in which of the following?

- (a) 40 – 50 (b) 30 – 40 (c) 50 – 60 (d) 60 – 70

Ans: (c) 50 – 60

67.If the mean of x , $x + 3$, $x + 5$, $x + 7$ and $x + 10$ is 9 then what is the means of last three observations?

- (a) $9\frac{4}{5}$ (b) $10\frac{1}{2}$ (c) $11\frac{1}{3}$ (d) 12

Ans: (c) $11\frac{1}{3}$

68.If x be the mid-point and 1 be the upper class limit of a class in a continuous frequency distribution. What is the lower limit of the class?

- (a) $x - 1$ (b) $3x + 8$ (c) $2x + 2$ (d) $2x - 1$

Ans: (d) $2x - 1$

69.The mean of five numbers is 30. If one number is excluded, their mean becomes 28. What is the excluded number?

- (a) 38 (b) 35 (c) 33 (d) 36

Ans: (a) 38

70.What is the median of the data 78, 56, 22, 34, 45, 54, 39, 68, 54, 84?

- (a) 54 (b) 53 (c) 55 (d) 51

Ans: (a) 54

71.What is the median of the data set 5, 8, 10, 12, 15?

- (a) 8 (b) 10 (c) 12 (d) 15

Ans: (b) 10

72.If the mean of the numbers 2, 4, 6, 8, and 10 is increased by 2, what is the new mean?

- (a) 5 (b) 7 (c) 8 (d) 6

Ans: (c) 8

73.The mode of the data set 3, 5, 7, 5, 9, 5, 11 is:

- (a) 3 (b) 5 (c) 7 (d) 9

Ans: (b) 5

74.The range of the data set 15, 20, 35, 40, 50 is:

- (a) 30 (b) 35 (c) 40 (d) 50

Ans: (b) 35

75.Which of the following is affected the most by extreme values?

(a) Mean (b) Median (c) Mode (d) Range

Ans: (a) Mean

76. The sum of observations = 180 and number of observations = 9. Find the mean.

(a) 18 (b) 20 (c) 15 (d) 25

Ans: (b) 20

77. If the median of a data set is 15, this means:

(a) 15 is the most frequent value
(b) 15 is the middle value when data is arranged in order
(c) The average of all values is 15
(d) None of these

Ans: (b) 15

78. Which measure of central tendency can be used for qualitative data?

(a) Mean (b) Median (c) Mode (d) All of these

Ans: (c) Mode

79. If the mean of 5 numbers is 12, what is their total sum?

(a) 60 (b) 65 (c) 70 (d) 55

Ans: (a) 60

80. The mean of the observations 4, 8, 12, 16, 20 is:

(a) 10 (b) 12 (c) 14 (d) 16

Ans: (b) 12

81. Which of the following is not a measure of central tendency?

(a) Standard deviation (b) Mean (c) Median (d) Mode

Ans: (a) Standard deviation

82. The mean of the data 2, 3, 4, 5, 0, 1, 3, 3, 4, 3 is

(a) 2 (b) 2.2 (c) 2.4 (d) 2.8

Ans: (d) 2.8

83. The collection of information, collected for a purpose is called:

(a) Mean (b) Median (c) Mode (d) Data

Ans: (d) Data

84. The number of times a particular item occurs in a class interval is called its:

(a) Mean (b) Frequency (c) Cumulative frequency (d) Range

Ans: (b) Frequency

85. Find the maximum value if the range is 38 and the minimum value is 82.

(a) 60 (b) 76 (c) 120 (d) 82

Ans: (c) 120

Range = Maximum value – Minimum value.

$$38 = x - 82 \Rightarrow x = 38 + 82 = 120.$$

86. The difference between the maximum and minimum values of the given observation is called

- (a) Class (b) Class interval (c) Class mark (d) Range

Ans: (d) Range

87. If each data in the observation is decreased by 5, then the mean

- (a) Remains the same (b) Increased by 5
(c) Decreased by 5 (d) None of the above

Ans: (c) Decreased by 5

88. Find the mode of the following data: 15, 14, 19, 20, 14, 15, 16, 14, 15, 18, 14, 19, 15, 17, & 15.

- (a) 14 (b) 15 (c) 16 (d) 17

Ans: (b) 15

89. Find the value of x, if the arithmetic mean of 4, 5, 6, 7, 8 and x is 7.

- (a) 4 (b) 6 (c) 8 (d) 12

Ans: (d) 12

90. The arithmetic mean of the first 5 natural numbers is

- (a) 3 (b) 4 (c) 5 (d) 6

Ans: (a) 3

91. In the class intervals 10 - 20, 20 - 30, the number 20 is included in:

- (a) 10 - 20 (b) 20 - 30 (c) Both the intervals (d) None of these

Ans: (b) 20 - 30

92. One of the sides of a frequency polygon is

- (a) Either of the coordinate axes (b) The x - axis
(c) Neither of the coordinate axes (d) The y - axis

Ans: (b) The x - axis

93. Find the class width for the grouped frequency distribution of the class intervals 1 - 20, 21 - 40, 41 - 60 ...

- (a) 10 (b) 15 (c) 17 (d) 20

Ans: (d) 20

94. What is the class mark of the class interval 80 - 120?

- (a) 90 (b) 105 (c) 115 (d) 100

Ans: (d) 100

95. Find the range of the following data: 25, 18, 20, 22, 16, 6, 17, 15, 12, 30, 32, 10, 19, 8, 11, & 20.

- (a) 10 (b) 15 (c) 18 (d) 26

Ans: (d) 26

96. The value which appears very frequently in a data is called:

- (a) Mean (b) Median (c) Mode (d) Central tendency

Ans: (c) Mode

97. The mode of the given data: 4, 6, 5, 9, 3, 2, 7, 7, 6, 5, 4, 9, 10, 10, 3, 4, 7, 6, 9, 9 is;
(a) 7 (b) 9 (c) 10 (d) 6

Ans: (b) 9

98. Given the class intervals 1 - 10, 11 - 20, 21 - 30,, then 20 is considered in class
(a) 11 - 30 (b) 11 - 20 (c) 21 - 30 (d) 15 - 25

Ans: (b) 11 - 20

99. The median of the data: 17, 2, 7, 27, 15, 5, 14, 8, 10, 24, 48, 10, 8, 7, 18, 28 is:
(a) 10 (b) 24 (c) 12 (d) 8

Ans: (c) 12

100. A student collects information about the number of school going children in a locality consisting of a hundred households. The data collected by him is
(a) Arrayed data (b) Primary data
(c) Secondary data (d) Grouped data

Ans: (b) Primary data