

ANALYTIC PLANETARY TRANSIT LIGHT CURVES AND DERIVATIVES FOR STARS WITH POLYNOMIAL LIMB DARKENING

ERIC AGOL,^{1,2,*} RODRIGO LUGER,^{3,2} AND DANIEL FOREMAN-MACKEY³

¹*Department of Astronomy, University of Washington, Seattle, WA*

²*Virtual Planetary Laboratory, University of Washington, Seattle, WA*

³*Center for Computational Astrophysics, Flatiron Institute, New York, NY*

ABSTRACT

We derive analytic, closed-form solutions for the light curve of a planet transiting a star with a limb darkening profile which is a polynomial function of the stellar elevation, up to arbitrary integer order. We provide improved analytic expressions for the uniform, linear, and quadratic limb-darkened cases, as well as novel expressions for higher order integer powers of limb darkening. The formulae are crafted to be numerically stable over the expected range of usage. We additionally present analytic formulae for the partial derivatives of instantaneous flux with respect to the radius ratio, impact parameter, and limb darkening coefficients. These expressions are rapid to evaluate, and compare quite favorably in speed and accuracy to existing transit light curve codes. We also use these expressions to numerically compute the first partial derivatives of exposure-time averaged transit light curves with respect to all model parameters. An additional application is modeling eclipsing binary or eclipsing multiple star systems in cases where the stars may be treated as spherically symmetric. We provide code which implements these formulae in C++, Python, and Julia, with tests and examples of usage.

Keywords: methods: analytic — techniques: photometric

* Guggenheim Fellow

1. INTRODUCTION

The precise measurement of the transits of an exoplanet offers a host of information about the planet’s properties. To start with, the times of transit give the planet’s orbital ephemeris. The depth of transit, corrected for stellar limb darkening, gives the planet’s radius relative to that of the star. The shape of the transit, especially the duration of ingress and egress relative to the full transit duration, yields the orbital impact parameter of the planet. Beyond these basic properties, if the transit depth is seen to vary with wavelength, the presence of spectral features may be used to constrain the chemical composition of the planet’s atmosphere. If the transit times are seen to vary, a dynamical model can constrain the masses of the planet companions (and vice versa). If the planet is seen in eclipse, its temperature, emission spectrum, and atmospheric circulation pattern can be constrained. When combined with radial velocity measurements, the bulk density of a planet can be inferred, yielding constraints on its bulk composition.

And yet, all of these inferences are predicated on the precise computation of models of the planetary transit which may be used to infer the model parameters. Stars are non-uniform in brightness, with the general trend of growing dimmer towards the limb, and so limb darkening must be accounted for to accurately infer the planetary parameters (Csizmadia 2018). Indeed, fast and accurate computation of limb-darkened transit light curve models has enabled the detection and characterization of thousands of transiting exoplanets (Mandel & Agol 2002). The most important ingredient to these models has been a description of the limb darkening model which is flexible enough and accurate enough to describe the emission from a stellar photosphere. Linear and quadratic limb darkening laws were sufficient for lower-precision measurements; however, the measurement of transit light curves has steadily improved in precision. Higher order terms or non-linear laws have become necessary to describe higher precision measurements (Kopal 1950; Claret 2000; Giménez 2006), which tend to involve more computational burden.

In addition to computing transit light curves, the derivatives of these light curves with respect to the model parameters are also beneficial for accurate characterization of exoplanets. The derivatives enable fast and stable optimization of the transit light curve parameters, which is critical for obtaining initial estimates for a Markov Chain Monte Carlo simulation (MCMC; e.g. Ford 2005, 2006), for looking for multi-modal solutions, for initializing the multi-nest algorithm (Feroz & Hobson 2008), or for computing the Fisher information matrix. In some cases, MCMC can be slow to converge, and derivatives can accelerate convergence by adding an artificial momentum term to the log likelihood, and then allowing the sampler to follow contours of constant “energy.” This so-called “hybrid” or “Hamiltonian” MCMC approach holds great promise, but its application has been hampered by the lack of models with derivatives, as derivatives are in general more difficult to compute (Girolami & Calderhead 2011; Betancourt 2017).

Finally, the analytic computation of transit light curves with quadratic limb darkening is limited by numerical round-off error for limiting values near some special cases. In particular, when the radius equals the impact-parameter, which corresponds to the edge of the planet crossing the center of the star, the computation of the elliptic integrals becomes unstable. At the second and third points of contact, when the radius of the planet plus the impact parameter equals the radius of the star, the elliptic integrals diverge logarithmically. In the limit that the impact parameter approaches zero, the equations can also diverge. All of these special cases are in principle encountered rarely, but in practice with thousand of planets with tens to thousands of transits each, along with hundreds to hundreds of thousands of light curves with time sub-sampled for each exposure, these rare cases can be encountered with some frequency.

Based on these considerations, the primary goals of the current paper are threefold:

1. To extend the analytic quadratic limb-darkened transit model to higher order.
2. To compute the derivatives of the model analytically.
3. To stabilize the analytic light curve computation (and its derivatives) in all limits near special cases.

Secondary goals include modeling eclipsing binaries, for which the same considerations apply, and integrating the light curve model, and its derivatives, quickly and accurately over time to account for finite exposure times.

Some progress has been made already towards these goals. To describe this progress, we pause first to introduce some notation. Limb darkening models of spherical stars are parameterized with the cosine of the angle measured from the sub-stellar point, $\mu = \cos \theta$, where θ is the polar angle on the photosphere, with $\theta = 0$ at the center of the observed stellar disk, and $\theta = \pi/2$ at the limb. In a coordinate system in which the projected disk of the star lies in the $x - y$ plane, and the z coordinate points towards the observer, then $\mu = z$, where z is measured in units of the stellar radius. In terms of b , the normalized radius projected onto the sky, this parameter is given by $\mu = \cos \theta = \sqrt{1 - b^2}$, where $0 \leq b \leq 1$ within the stellar disk. We also introduce the radius ratio, r , which is the radius of the occulter divided by the radius of the source. In general, we will closely follow the notation introduced by [Luger et al. \(2018\)](#) for the `starry` code package.

Uniform limb darkening scales as $I(\mu) \propto \mu^0$, first-order limb darkening as $I(\mu) \propto \mu^1$, and second-order limb darkening as $I(\mu) \propto \mu^2$; these are the three most commonly used terms that can be integrated analytically, which we describe in detail below in sections 2, 3, and 4. These are typically combined to yield the quadratic limb darkening law,

$$I(\mu) = I(1)[1 - u_1(1 - \mu) - u_2(1 - \mu)^2]. \quad (1)$$

In this paper, we will show that higher order powers of μ^n with integer n can be integrated analytically when expressed as recursion relations. Linear combinations

of these laws can be constructed, with various parameterizations, to describe stellar limb darkening more precisely.

The first goal, of modelling higher-order limb darkening, was accomplished by Giménez (2006), who derived transit light curves for a limb darkening function

$$I(\mu) = I(1) \left[1 - \sum_{n=1}^N a_n (1 - \mu^n) \right], \quad (2)$$

where a_n is a limb darkening coefficient. Giménez (2006) found an infinite series expansion for computing the limb-darkened light curve for each a_n term. Here we present closed-form expressions for these terms which can be easily computed with recursion relations, although for purposes of numerical stability we need to revert to series solutions in some limits.

The second goal, of computing derivatives of the light curve with respect to the model parameters, was accomplished by Pál (2008) for the quadratic limb darkening case. Pál derived the partial derivatives of the quadratic limb darkening model with respect to b , r , and the two quadratic limb darkening coefficients. In this work, we give modified expressions for the quadratic limb-darkened flux and its derivatives which are more numerically stable, as well as extend the computation of derivatives to higher order limb darkening.

The third goal, of numerical stability, has yet to be addressed in the literature. Although some numerical approaches are numerically stable, such as Giménez (2006), Kreidberg (2015), and Parviainen & Aigrain (2015), these approaches tend to be slower, they have precisions which may depend upon the tolerance of the computation, and, in addition, they do not yield derivatives of the light curves.

A disadvantage of our approach is that it requires integer powers of the limb darkening expansion. Claret (2000) has shown that a non-linear limb darkening law, with half-integer powers of μ , gives an accurate description of stellar limb darkening models. More recently, the power-law model, $I(\mu) = 1 - c_\alpha(1 - \mu^\alpha)$ (Hestroffer 1997) was shown to be an accurate limb darkening law despite only using two parameters (Morello et al. 2017; Maxted 2018). We were unable to find an analytic solution for these limb darkening laws, but we will compare with these models below in §10.

We turn now towards presenting formulae for more accurate limb darkening transit, occultation, and eclipse models, starting with the simplest case: a source of uniform surface brightness.

Eric: Before we begin defining the \mathfrak{s}_0 , \mathfrak{s}_1 , ... functions, we need to explain how they relate to the total flux. I know we do this later on, but I think we'll really confuse the reader if we make them wait that long. We should dedicate a paragraph to explaining how we're doing the same thing as in starry: expressing the flux as the dot product between a polynomial basis and a vector of integral solutions.

2. UNIFORM BRIGHTNESS

Evaluation of the transit light curve of a uniformly bright star amounts to computing the area of overlap of two disks (Mandel & Agol 2002). This has a well-known analytic solution (e.g. Weisstein 2018); however, we find that the standard formula leads to round-off error that is larger than necessary or desirable. In this section we present a new formula which we demonstrate yields machine precision for the area of overlap, along with its derivatives.

Figure 1 shows how the area of overlap can be computed for two circles. The sums of the areas of the sectors of each circle which span the area of overlap, minus the area of a kite-shaped region which connects the centers of the circles with their points of intersection gives the area of the lens-shaped region of overlap of the two circles.

Taking the radius of the larger circle to be unity, the standard formula for the lens-shaped overlap area is given by

$$A_{lens} = \pi \Lambda^e(r, b) = \begin{cases} 0 & 1 + r \leq b, \\ r^2 \kappa_0 + \kappa_1 - \sqrt{\frac{4b^2 - (1 + b^2 - r^2)^2}{4}} & |1 - r| < b \leq 1 + r, \\ \pi r^2 & b \leq 1 - r, \\ \pi & b \leq r - 1, \end{cases} \quad \text{Eq. (3)}$$

(e.g. Mandel & Agol 2002), where

$$\begin{aligned} \cos \kappa_0 &= \left(\frac{(r - 1)(r + 1) + b^2}{2br} \right), \\ \cos \kappa_1 &= \left(\frac{(1 - r)(1 + r) + b^2}{2b} \right), \end{aligned} \quad (4)$$

where κ_0 and κ_1 are the angles defined in Figure 1. The second term is the same as the standard formula for the area of overlap of two partially overlapping circles, with one of the circles scaled to a radius of unity (Weisstein 2018). This term corresponds to ingress (and egress), it is the most expensive to compute, and it is most subject to numerical inaccuracy; we focus on this term in what follows.

We find that numerical round-off error limits the precision of the ingress formula when $b \approx 0$, $b + r \approx 1$, or $b \approx 1 + r$; these are the cases in which the kite-shaped region becomes thin, in which the sum of two sides becomes similar in length to the spine of the kite. The square root term in this formula (Equation 3) computes the area of the kite-shaped region, which in this form causes round-off error when the kite is flattened. The same issue occurs when computing the area of a triangle in which two of the sides are of similar length; the kite has an area that is twice the area of

the triangle connecting the centers of both circles and one of the intersection points. [Goldberg \(1991\)](#) gives a formula for precisely computing the area of a triangle, based on a method developed by [Kahan \(2000\)](#), which we use to compute the area of the kite-shaped region,

$$A_{kite} = \frac{1}{2} \sqrt{(A + (B + C))(C - (A - B))(C + (A - B))(A + (B - C))}, \quad (5)$$

for $A \geq B \geq C$, where the tuple $\{A, B, C\}$ equals $\{1, r, b\}$ sorted from greatest to least. **Eric:** The previous sentence states that the 1991 formula is based on a method published in 2000; please check.

Next, the inverse cosine formulae are also imprecise when $\cos \kappa_0 = x_0 \approx 1$ or $\cos \kappa_1 = x_1 \approx 1$. The approximate solutions in this limit are $\kappa_0 \approx [2(1 - x_0)]^{1/2}$ and $\kappa_1 \approx [2(1 - x_1)]^{1/2}$, and so round-off error can occur both in taking the difference of two numbers close to unity, and in taking the square root.

Instead, we use the function $\theta = \text{atan2}(y, x)$ with $y = \sin \theta$ and $x = \cos \theta$ to compute κ_0 and κ_1 , which avoids the quadrant and division-by-zero problems of the $\theta = \tan^{-1}(y/x)$ function. In addition to the cosine values above, we require the sine terms, which are given by

$$\begin{aligned} \sin \kappa_0 &= \frac{A_{kite}}{br}, \\ \sin \kappa_1 &= \frac{A_{kite}}{b}, \end{aligned} \quad (6)$$

which can be derived from the area of the triangles formed by the centers of the circles and one intersection point. Note that both $\sin \kappa_0$ and $\cos \kappa_0$ are divided by br , and $\sin \kappa_1$ and $\cos \kappa_1$ are divided by b , so that in the arctangent formula these denominator terms cancel, which can improve numerical stability for small values of b or r ; this cancellation does not happen in the arccosine case given in Equation (4).

This results in the following equations for the overlap area, A_{lens} , of two partially overlapping circles:

$$\begin{aligned} A_{lens} &= \kappa_1 + r^2 \kappa_0 - A_{kite}, \\ \kappa_0 &= \text{atan2}(2A_{kite}, (r - 1)(r + 1) + b^2), \\ \kappa_1 &= \text{atan2}(2A_{kite}, (1 - r)(1 + r) + b^2). \end{aligned} \quad \text{🔗 (7)}$$

The performance of this formula relative to the standard formula is profiled in Figure 2 for $r = 0.1$, a typical value for transiting exoplanets. We have carried out the computation in the `Julia` language, both in double-precision (`Float64`), and 256-bit precision (`BigFloat`), and subtracted the results to measure the numerical errors of the computation.

We find that the standard formula (Equation 3) approaches errors of 10^{-8} in the limit of $b \rightarrow 1 - r$. This error exceeds the value of the area of the smaller circle minus the area of overlap for values of $1 - r < b < 1 - r + 10^{-8}$. Thus, even though this calculation is carried out in double precision, the precision achieved is of order single precision. Likewise, for $b \rightarrow 1 + r$, the error of the standard formula approaches 10^{-8} , with the error exceeding the value of the area of overlap for $1 + r - 10^{-8} < b < 1 + r$.

In contrast, Equation (7) gives a precision that is double-precision in both limits. Figure 2 shows that Equation (7) gives a precision of $\approx 10^{-17}$ in the limit $b \rightarrow 1 - r$ for $r = 0.1$; this limit is due to the limiting precision of representing πr^2 in double-precision, which in this case is $\pi r^2 / 2^{53} = 10^{-17.4}$, indicated with a dash-dot grey line in the left hand panels of Figure 2. At the beginning of ingress/end of egress when $b \approx 1 + r$, even higher precision is achieved since the area of overlap approaches zero, as shown in the right hand panels of Figure 2.

The final modification we make when computing the uniform light curve is that our computations make use of the visible area of the disk, \mathfrak{s}_0 , defined as

$$\begin{aligned}\mathfrak{s}_0(r, b) &= \pi - A_{lens} \\ &= \pi - \kappa_1 - r^2 \kappa_0 + A_{kite},\end{aligned}\tag{8}$$

rather than the obscured area, A_{lens} . So, instead of computing κ_1 , we compute $\pi - \kappa_1 = -\text{atan2}(\sin \kappa_1, \cos \kappa_1)$. We find that this leads to machine-precision as well. Note that \mathfrak{s}_0 is identical to the first basis function (s_0) in the **starry** implementation from Luger et al. (2018).

2.1. Derivatives

The partial derivatives of this formula with respect to the radius ratio, r , and impact parameter, b , turn out to be straightforward:

$$\begin{aligned}\frac{\partial \mathfrak{s}_0(r, b)}{\partial r} &= -2r \kappa_0, \\ \frac{\partial \mathfrak{s}_0(r, b)}{\partial b} &= \frac{2A_{kite}}{b},\end{aligned}\tag{9}$$

which can be computed from the quantities already used in calculating \mathfrak{s}_0 . At the contact points, when $b = |1 \pm r|$, the derivatives are undefined. In practice this can be a problem when taking finite-differences across the discontinuous boundary, but with the analytic formulae, these points are a set of measure zero, and so we simply set the derivatives to zero at these points.

In the remainder of this paper we will need to use these formulae in computing the higher order limb-darkened light curves. In the next section, we revisit the formulae for linear limb darkening.




Figure 1. The area of overlap of two circles can be computed as the sum of the area of the sectors formed by the centers of each circle and the boundary between the points of intersection, minus the area of the kite-shaped region formed by the centers of the circles and the intersection points. [↗](#)




Figure 2. Precision of formulae for the area of overlap of two circles with radius ratio r . Plotted are the regions near $b = 1 - r$ (second and third points of contact) and $b = 1 + r$ (first and fourth points of contact) for the standard formula (Equation 3, blue) and our new formula (Equation 7, orange dots). The high-precision calculation is shown in green dashed for comparison; this is limited by the conversion of the result to double-precision. The solid and empty circles (top) indicate the positions of the circles at the left hand side of the axes. In the left panels the horizontal dash-dotted grey line indicates the limiting precision for representing πr^2 . [↗](#)

3. LINEAR LIMB DARKENING

In this section we turn to the case of linear limb darkening, $I(\mu)/I(1) = 1 - u_1(1 - \mu)$ (Russell & Shapley 1912a,b). In this section we set $I(1) = u_1 = 1$, so that $I(\mu) = \mu$; the general case can be computed as a linear combination with the uniform case. Note that since $\mu = \sqrt{1 - x^2 - y^2}$, this problem is equivalent to computing the volume of intersection between a sphere and a cylinder, which was solved in terms of elliptic integrals by Lamarche & Leroy (1990). A similar solution was found by Mandel & Agol (2002), who show that the total flux visible during the occultation of a body whose surface map is given by $I(x, y) = \sqrt{1 - x^2 - y^2}$ may be computed as

$$\mathfrak{s}_1 = \frac{2\pi}{3} \left(1 - \frac{3\Lambda(r, b)}{2} - \Theta(r - b) \right) \quad (10)$$

where $\Theta(\cdot)$ is the Heaviside step function and

$$\Lambda(r, b) = \begin{cases} \frac{1}{9\pi\sqrt{br}} \left[\frac{(r+b)^2 - 1}{r+b} \left(-2r(2(r+b)^2 + (r+b)(r-b) - 3)K(k^2) \right. \right. \\ \left. \left. + 3(b-r)\Pi(k^2(b+r)^2, k^2) \right) - 4br(4 - 7r^2 - b^2)E(k^2) \right] & k^2 < 1 \\ \frac{2}{9\pi} \left[(1 - (r+b)^2) \left(\sqrt{1 - (b-r)^2} K\left(\frac{1}{k^2}\right) + 3 \left(\frac{b-r}{(b+r)\sqrt{1 - (b-r)^2}} \right) \right. \right. \\ \left. \left. \times \Pi\left(\frac{1}{k^2(b+r)^2}, \frac{1}{k^2}\right) \right) - \sqrt{1 - (b-r)^2}(4 - 7r^2 - b^2)E\left(\frac{1}{k^2}\right) \right] & k^2 \geq 1 \end{cases} \quad (11)$$

with

$$k^2 = \frac{1 - r^2 - b^2 + 2br}{4br}. \quad (12)$$

Note that $\mathfrak{s}_1(r, b) = s_2(r, b)$ in the spherical harmonic expansion used in **starry** as described in Luger et al. (2018). For the cases $b = r$, $b = 1 - r$, $b = 0$, $r = 0$, or $|r - b| \geq 1$, there are special expressions for $\Lambda(r, b)$ given below. In the expressions above, $K(\cdot)$, $E(\cdot)$, and $\Pi(\cdot, \cdot)$ are the complete elliptic integrals of the first, second

kind, and third kind, respectively, defined as

$$\begin{aligned}
 K(k^2) &\equiv \int_0^{\frac{\pi}{2}} \frac{d\varphi}{\sqrt{1 - k^2 \sin^2 \varphi}} \\
 E(k^2) &\equiv \int_0^{\frac{\pi}{2}} \sqrt{1 - k^2 \sin^2 \varphi} d\varphi \\
 \Pi(n, k^2) &\equiv \int_0^{\frac{\pi}{2}} \frac{d\varphi}{(1 - n \sin^2 \varphi) \sqrt{1 - k^2 \sin^2 \varphi}}.
 \end{aligned} \tag{13}$$

In Equation (11) we have transformed the formulae from [Mandel & Agol \(2002\)](#) using Equation (17.7.17) from [Abramowitz & Stegun \(1970\)](#) which yields equations that are better behaved in the vicinity of $b = r$.¹ However, these elliptic integrals are still subject to numerical instability as $r \rightarrow 1 - b$ and $r \gg 1$. The main issue is the logarithmic divergence of K and Π as $k \rightarrow 1$, as well as numerical cancellations leading to round-off errors which occur in the limit $k \rightarrow 0$.

Through trial and error, we have found that these instabilities can be removed by combining elliptic integrals into a general complete elliptic integral defined by [Bulirsch \(1969\)](#) as

$$\text{cel}(k_c, p, a, b) = \int_0^{\pi/2} \frac{a \cos^2 \phi + b \sin^2 \phi}{\cos^2 \phi + p \sin^2 \phi} \frac{d\phi}{\sqrt{\cos^2 \phi + k_c^2 \sin^2 \phi}}, \tag{14}$$

where $k_c = \sqrt{1 - m_k}$, and for $b + r \geq 1$, $m_k = k^2$, while for $b + r \leq 1$, $m_k = 1/k^2$. Although k_c can be computed from m_k , we have found better numerical stability in computing k_c analytically from b and r :

$$k_c = \begin{cases} \sqrt{\frac{(b+r)^2 - 1}{4br}} & k^2 \leq 1 \\ \sqrt{\frac{1 - (b+r)^2}{1 - (b-r)^2}} & k^2 > 1. \end{cases} \tag{15}$$

In practice, we let the subroutine that computes cel accept both m_k and k_c as input for numerical precision.

¹ Note that we corrected several typos in [Mandel & Agol \(2002\)](#), which are listed in the Appendix.

To transform the elliptic integrals in Equation (11), we used the following relations from Bulirsch (1969):

$$\lambda K(m_k) + qE(m_k) = \text{cel}(k_c, 1, \lambda + q, \lambda + qk_c^2) \quad (16)$$

$$\lambda K(m_k) + q\Pi(n, m_k) = \text{cel}(k_c, 1 - n, \lambda + q, \lambda + qk_c^2) \quad (17)$$

$$E(m_k) = \text{cel}(k_c, 1, 1, 1 - m_k) \quad (18)$$

$$E(m_k) - (1 - m_k)K(m_k) = m_k \text{cel}(k_c, 1, 1, 0) \quad (19)$$

$$\Pi(n, m_k) - K(m_k) = n \text{cel}(k_c, 1 - n, 0, 1), \quad (20)$$

noting that Bulirsch (1969) uses a different sign convention for $\Pi(n, m_k)$. In particular, the expressions for $\Pi(n, m_k) - K(m_k)$ and $E(m_k) - (1 - m_k)K(m_k)$ are useful for eliminating the singularities and cancellations which occur at $m_k = 1$ when $b + r = 1$ and $m_k = 0$ when $r \rightarrow \infty$. The general complete elliptic integral is evaluated with the approach of Bartky (1938), which uses recursion to approximate the integral to a specified precision.

These elliptic integral transformations lead to the following numerically-stable expression for the linear limb darkening flux, $\mathfrak{s}_1(r, b)$, in which

$$\Lambda = \begin{cases} 0 & r = 0 \\ 0 & |r - b| \geq 1 \\ -\frac{2}{3}(1 - r^2)^{3/2} & b = 0 \\ \frac{1}{3} - \frac{4}{9\pi} & b = r = \frac{1}{2} \\ \frac{1}{3} + \frac{2}{9\pi} \text{cel}(k_c, 1, m_k - 3, (1 - m_k)(2m_k - 3)) & b = r < \frac{1}{2} \\ \frac{1}{3} + \frac{4r}{9\pi} \text{cel}(k_c, 1, 1 - 3m_k, m_k - 1) & b = r > \frac{1}{2} \\ \frac{2}{9\pi} \left[3 \cos^{-1}(1 - 2r) - 2(3 + 2r - 8r^2)\sqrt{rb} - 3\pi\Theta(r - \frac{1}{2}) \right] & b + r = 1 \\ \frac{1 - (b - r)^2}{9\pi\sqrt{br}} \left[\frac{(b + r)^2 - 1}{4br} (b^2 - r^2) \text{cel}(k_c, (b - r)^2(1 - m_k), 0, 3) \right. \\ \quad \left. - (3 - 6r^2 - 2br) \text{cel}(k_c, 1, 1, 0) - 4brE(m_k) \right] & k^2 < 1 \\ \frac{2\sqrt{1 - (b - r)^2}}{9\pi} \left[(1 - (r + b)^2) \text{cel}(k_c, p, 1 + q, p + q) \right. \\ \quad \left. - (4 - 7r^2 - b^2)E(m_k) \right] & k^2 > 1 \end{cases} \quad (21)$$

where

$$\begin{aligned} q &= 3 \frac{b-r}{(b+r)(1-(b-r)^2)} \\ p &= \left(\frac{b-r}{b+r} \right)^2 \frac{1-(b+r)^2}{1-(b-r)^2} \end{aligned} \quad (22)$$

in the $k^2 > 1$ case. Note that in this equation the conditions should be evaluated in the order they appear.

The $\mathfrak{s}_1(r, b)$ function is plotted in Figure 3. The function varies smoothly from the lower right where the disk is unocculted to the upper left where it is completely occulted. There are several points which need to be handled separately as the Equation (11) expressions become singular or are no longer valid; the solid lines in Figure 3 show these points. When $b = 0$, the integral over the center of the disk simplifies greatly. When $b = r = 1/2$, at the intersection of $b = r$ and $b = 1 - r$, another simplification occurs. For $b = r$, the disk of the occulter crosses the center of the source; this needs to be computed separately in the $r < 1/2$, $r = 1/2$, and $r > 1/2$ limits. The first and fourth contacts occur at $b = 1 + r$, where $\mathfrak{s}_1 = 1$; this is the upper bound to the $k^2 < 1$ region for $b + r > 1$. For $r \geq 1$, the second and third contacts (at the start and end of complete occultation) occur when $b = 1 - r$, which is the lower bound to the $k^2 < 1$ region when $b + r > 1$. For $r < 1$, the second and third contacts occur when $r = 1 - b$.

Near these boundaries, the standard Mandel & Agol (2002) expressions can become singular, and so we paid particular care to the accuracy of these new expressions in these regions. Figure 4 shows that Equation (21) is accurate to machine precision in all of these regimes. We tested the accuracy by computing the equations with 256 bit arithmetic, which is much less subject to round-off error, and hence gives more precise expressions than double precision. We implemented the pseudocode from Bulirsch (1969) to compute $\text{cel}(k_c, p, a, b)$, which has a termination test that scales as the square root of the machine precision. We find that the transformed expressions are accurate to $\lesssim \times 10^{-14}$ when computed in double precision within $\epsilon = 10^{-8}$ of the vicinity of $b = r$ and $b = 1 - r$.

Finally, in Figures 5 and 6 we plot the relative numerical error in the flux of a linearly limb-darkened source when using the equations in Mandel & Agol (2002) and in this paper, respectively, over a portion of the $b - r$ plane. The former method (Figure 5) yields errors on the order of 10^{-7} over most of the domain, although the error approaches unity near the singular regions discussed above. In contrast, the method introduced in this paper (Figure 6; note the change in the color scale) yields errors close to machine precision everywhere, including the vicinity of the singular points.

3.1. Derivatives




Figure 3. The intensity of a linearly limb-darkened star ($u_1 = 1$) being eclipsed, $\mathfrak{s}_1(r, b)$. In the limit $b > r + 1$, no eclipse occurs, so $\mathfrak{s}_1 = 1$. For $b < r - 1$, the star is completely eclipsed and $\mathfrak{s}_1 = 0$. In the limits $b = r$ and $b = 1 - r$, special expressions must be used. [☐](#)




Figure 4. The accuracy of $\mathfrak{s}_1(r, b)$ near $b = r$ (left panel) and $b = 1 - r$ (right panel) for $\epsilon = 10^{-8}$. The x -axes are impact parameter b , while the y axes in the top panels show $\mathfrak{s}_1(r, b)$, with r given in the legend of each panel. The middle panels plot the difference $(\mathfrak{s}_1(b, b \pm \epsilon) - \mathfrak{s}_1(b, b))/\epsilon$ and $(\mathfrak{s}_1(b, 1 - b \pm \epsilon) - \mathfrak{s}_1(b, 1 - b))/\epsilon$. The bottom panels show the numerical precision by the comparing double precision computation with **BigFloat** precision (256-bit). [☐](#)




Figure 5. The numerical error in computing the flux of an eclipsed, linearly limb-darkened star ($u_1 = 1$) using the equations in Mandel & Agol (2002). [↗](#)




Figure 6. The numerical error in computing the flux of an eclipsed, linearly limb-darkened star ($u_1 = 1$) using the $s_1(r, b)$ formalism introduced in this paper. Compare to Figure 5, noting the change in the color scale. The new method is eight orders of magnitude more precise on average, approaching machine epsilon everywhere in the domain. [↗](#)

The derivatives of Λ with respect to r and b are:

$$\frac{\partial \Lambda}{\partial r} = \begin{cases} 0 & r = 0 \\ 0 & |r - b| \geq 1 \\ 2r\sqrt{1 - r^2} & b = 0 \\ \frac{2}{\pi} & b = r = \frac{1}{2} \\ \frac{4r}{\pi} E(4r^2) & b = r < \frac{1}{2} \\ \frac{2}{\pi} \text{cel}(k_c, 1, 1, 0) & b = r > \frac{1}{2} \\ \frac{8r}{\pi} \sqrt{r(1 - r)} & b + r = 1 \\ \frac{8br^2 E(k^2) + 2r(1 - (b + r)^2) K(k^2)}{\pi \sqrt{br}} & \\ = \frac{1}{\pi \sqrt{br}} \text{cel}(k_c, 1, 2r(1 - (b - r)^2), 0) & k^2 < 1 \\ \frac{4r}{\pi} \sqrt{1 - (b - r)^2} E(k^{-2}) & \\ = \frac{4r}{\pi} \sqrt{1 - (b - r)^2} \text{cel}(k_c, 1, 1, k_c^2) & k^2 > 1 \end{cases} \quad (23)$$

and

$$\frac{\partial \Lambda}{\partial b} = \begin{cases} 0 & r = 0 \\ 0 & |r - b| \geq 1 \\ 0 & b = 0 \\ -\frac{2}{3\pi} & b = r = \frac{1}{2} \\ \frac{4r}{3\pi} \text{cel}(k_c, 1, -1, k_c^2) & b = r < \frac{1}{2} \\ -\frac{2}{3\pi} \text{cel}(k_c, 1, 1, 2k_c^2) & b = r > \frac{1}{2} \\ -\frac{8r}{3\pi} \sqrt{r(1 - r)} & b + r = 1 \\ \frac{4r(r^2 + b^2 - 1)E(k^2) + 2r(1 - (b + r)^2)K(k^2)}{3\pi \sqrt{br}} & \\ = \frac{1 - (b - r)^2}{3\pi \sqrt{br}} \text{cel}(k_c, 1, -2r, (1 - (b + r)^2)/b) & k^2 < 1 \\ \frac{2}{3b\pi} \sqrt{1 - (b - r)^2} [(r^2 + b^2 - 1)E(k^{-2}) + (1 - (b + r)^2)K(k^{-2})] & \\ = \frac{4r}{3\pi} \sqrt{1 - (b - r)^2} \text{cel}(k_c, 1, -1, k_c^2) & k^2 > 1, \end{cases} \quad (24)$$

where we have given some of the expressions in terms of both the standard elliptic integrals and the general elliptic integral.

Eric: We need an equation here explicitly stating the derivatives of the \mathfrak{s}_2 term.

We have tested these formulae with finite-difference derivatives evaluated at 256-bit precision, and, as with the total flux term, we find that these are accurate to $\lesssim 2 \times 10^{-15}$, close to machine precision.

We next increase the power of limb darkening by one, μ^2 .

4. QUADRATIC LIMB DARKENING

The next order of limb darkening has been widely studied due to its accurate description of stellar atmospheres (Claret 2000; Mandel & Agol 2002; Pál 2008). We summarize here the formulae for quadratic limb darkening with $I(1) = 1$, $u_1 = 2$ and $u_2 = -1$, so that $I(\mu) = \mu^2$, along with the derivatives, using the transformed expressions described above. The general quadratic case may be computed as a linear combination with the foregoing uniform and linear cases.

We first give the formula for the function $\eta(r, b)$, which is used in Mandel & Agol (2002), in terms of quantities we have defined above for the uniform case:

$$\eta(r, b) = \begin{cases} \frac{1}{2\pi} \left[\kappa_1 + r^2(r^2 + 2b^2)\kappa_0 - \frac{1}{2}(1 + 5r^2 + b^2)A_{kite} \right] & k^2 \leq 1 \\ \frac{r^2}{2}(r^2 + 2b^2) & k^2 > 1 \end{cases} \quad \text{\textcolor{blue}{\(\square\)}} \quad (25)$$

As κ_0 , κ_1 , and A_{kite} were already computed in the uniform case, these quantities are reused in the quadratic computation.

With this definition, the quadratic term, $\mathfrak{s}_2(r, b)$ is given simply by

$$\mathfrak{s}_2 = 2\mathfrak{s}_0 + 4\pi\eta - 2\pi, \quad (26)$$

$$(27)$$

where \mathfrak{s}_0 is defined in Equation (8).

4.1. Derivatives

The derivatives of η are given by:

$$\frac{\partial \eta}{\partial r} = \begin{cases} \frac{2r}{\pi} [(r^2 + b^2)\kappa_0 - 2A_{kite}] & k^2 \leq 1 \\ 2r(r^2 + b^2) & k^2 > 1 \end{cases} \quad \text{\textcolor{blue}{\(\square\)}} \quad (28)$$

and

$$\frac{\partial \eta}{\partial b} = \begin{cases} \frac{1}{2b\pi} [4r^2 b^2 \kappa_0 - 2(1 + b^2 + r^2) A_{kite}] & k^2 \leq 1 \\ 2br^2 & k^2 > 1 \end{cases} \quad (29)$$

where the derivatives of \mathfrak{s}_0 are defined in Equation (9). The derivatives of the \mathfrak{s}_2 term are thus

$$\begin{aligned} \frac{\partial \mathfrak{s}_2}{\partial r} &= 2 \frac{\partial \mathfrak{s}_0}{\partial r} + 4\pi \frac{\partial \eta}{\partial r}, \\ \frac{\partial \mathfrak{s}_2}{\partial b} &= 2 \frac{\partial \mathfrak{s}_0}{\partial b} + 4\pi \frac{\partial \eta}{\partial b}. \end{aligned} \quad (30)$$

In the following section we turn our attention to the general polynomial limb darkening case, μ^n with $n > 2$. As discussed in Luger et al. (2018), these terms may be expressed exactly as the sum of spherical harmonics with $m = 0$. However, it is possible to exploit the azimuthal symmetry of the limb darkening problem to derive far more efficient and accurate formulae, which we describe in the following section.

5. HIGHER ORDER LIMB DARKENING

In analogy with the linear and quadratic limb darkening laws, let us define the polynomial limb darkening law of order N as

$$\begin{aligned} \frac{I(\mu)}{I_0} &= 1 - u_1(1 - \mu) - u_2(1 - \mu)^2 - \dots - u_N(1 - \mu)^N \\ &= - \sum_{i=0}^N u_i (1 - \mu)^i \end{aligned} \quad (31)$$

where $I_0 \equiv I(1)$, $u_0 \equiv -1$, and μ is the cosine of the angle between the line of sight and the vector normal to the body's surface. In a right-handed Cartesian coordinate system centered on the body, with the z -axis pointing to the observer,

$$\mu(x, y) = z(x, y) = \sqrt{1 - x^2 - y^2}. \quad (32)$$

If we let \mathbf{u} be the column vector of limb darkening coefficients $\mathbf{u} \equiv (u_0 \ u_1 \ u_2 \ \dots \ u_N)^\top$ and $\tilde{\mathbf{u}}$ be the *limb darkening basis*

$$\tilde{\mathbf{u}} = - \begin{pmatrix} 1 & (1 - z) & (1 - z)^2 & \dots & (1 - z)^N \end{pmatrix}^\top, \quad (33)$$

we may express Equation (31) more compactly as the dot product

$$\frac{I(z)}{I_0} = \tilde{\mathbf{u}}^\top \mathbf{u}. \quad (34)$$

Our task now is to compute the flux, F , observed during an occultation by integrating this function over the visible area of the disk:

$$F = \oiint I(z) \, dS. \quad (35)$$

In general, the surface integral in Equation (35) is difficult—if not impossible—to solve directly with $I(z)$ given by Equation (34). However, as in Luger et al. (2018), we note that the problem is made significantly more tractable if we first perform a couple change of basis operations.

5.1. Change of basis

We wish to find a basis in which to express the limb darkening profile that makes evaluating Equation (35) easy. This section follows closely the discussion in Luger et al. (2018), in which the authors first transform to a *polynomial basis*, whose terms are simple powers of the coordinates, and then to a *Green's basis*, whose terms make application of Green's theorem convenient in reducing the surface integral to a one-dimensional line integral.

Let us define the transformation to the polynomial basis by the linear equation

$$\mathbf{p} = \mathcal{A}_1 \mathbf{u} \quad (36)$$

where \mathbf{p} is the vector of limb darkening coefficients in the polynomial basis $\tilde{\mathbf{p}}$ and \mathbf{A}_1 is a change of basis matrix. We define the polynomial basis to be the power series in z ,

$$\tilde{\mathbf{p}} = \begin{pmatrix} 1 & z & z^2 & z^3 & \dots & z^N \end{pmatrix}^\top. \quad (37)$$

The transformation between vectors in $\tilde{\mathbf{u}}$ and vectors in $\tilde{\mathbf{p}}$ is straightforward. By the binomial theorem, we may write the i^{th} coefficient of \mathbf{p} as

$$\mathbf{p}_i = \sum_{j=0}^N \binom{i}{j} (-1)^{j+1} u_j. \quad (38)$$

The elements of the matrix \mathbf{A}_1 are thus given by

$$\mathcal{A}_{1_{i,j}} = \binom{i}{j} (-1)^{j+1}. \quad (39)$$

Note that, as with the limb darkening basis, the specific intensity at a point may be written

$$\begin{aligned} \frac{I(z)}{I_0} &= \tilde{\mathbf{p}}^\top \mathbf{p} \\ &= \tilde{\mathbf{p}}^\top \mathcal{A}_1 \mathbf{u}. \end{aligned} \quad (40)$$

Next, we transform to the Green's basis via the equation

$$\mathbf{g} = \mathcal{A}_2 \mathbf{p} \quad (41)$$

where \mathbf{g} is the vector of limb darkening coefficients in the Green's basis $\tilde{\mathbf{g}}$ and \mathcal{A}_2 is another change of basis matrix. For reasons that will become clear in the following section, we define the Green's basis to be

$$\begin{aligned} \tilde{\mathbf{g}}_n &= \begin{cases} 1 & n = 0 \\ z & n = 1 \\ (n+2)z^n - nz^{n-2} & n \geq 2 \end{cases} \\ \tilde{\mathbf{g}} &= \begin{pmatrix} 1 & z & 4z^2 - 2 & 5z^3 - 3z & \dots \end{pmatrix}^\top, \end{aligned} \quad (42)$$

Given this definition, the columns of the change of basis matrix \mathcal{A}_2 are the Green's basis vectors corresponding to each of the polynomial terms in Equation (37). Note that in practice, it is more efficient to transform vectors in the $\tilde{\mathbf{p}}$ basis to vectors in the $\tilde{\mathbf{g}}$ basis via the downward recursion relation

$$\mathbf{g}_n = \frac{\mathbf{p}_n}{n+2} + \mathbf{g}_{n+2}, \quad (43)$$

starting with $n = N$ and $\mathbf{g}_{N+1} = \mathbf{g}_{N+2} = 0$.

As before, the specific intensity at a point may be written

$$\begin{aligned}
 \frac{I(z)}{I_0} &= \tilde{\mathbf{g}}^\top \mathbf{g} \\
 &= \tilde{\mathbf{g}}^\top \mathcal{A}_2 \mathbf{p} \\
 &= \tilde{\mathbf{g}}^\top \mathcal{A} \mathbf{u},
 \end{aligned} \tag{44}$$

where we define the complete change of basis matrix from limb darkening coefficients to Green's coefficients

$$\mathcal{A} \equiv \mathcal{A}_2 \mathcal{A}_1. \tag{45}$$

5.2. Computing the surface integral

Given our reparametrization in terms of Green's polynomials, we may re-write Equation (35) as

$$\begin{aligned}
 F &= \oint\!\!\!\oint I(z) \, dS \\
 &= I_0 \oint\!\!\!\oint \tilde{\mathbf{g}}^\top \mathcal{A} \mathbf{u} \, dS \\
 &= I_0 \left(\oint\!\!\!\oint \tilde{\mathbf{g}}(z) \, dS \right)^\top \mathcal{A} \mathbf{u} \\
 &= I_0 \mathbf{s}^\top \mathcal{A} \mathbf{u},
 \end{aligned} \tag{46}$$

where

$$\mathbf{s} \equiv \oint\!\!\!\oint \tilde{\mathbf{g}}(z) \, dS \tag{47}$$

is the *solution vector* first introduced in §2. If we can find the general solution to the integral in Equation (47), we can compute the occultation flux for arbitrary order limb darkening.

As in Luger et al. (2018), we use Green's theorem to re-express the surface integral as a line integral over the boundary of the visible portion of the occulted body's disk, $\mathbf{r} = x\hat{\mathbf{x}} + y\hat{\mathbf{y}}$:

$$\mathbf{s} = \oint \mathbf{G}(z) \cdot d\mathbf{r}, \tag{48}$$

where \mathbf{G} is a matrix whose n^{th} row is the vector

$$\mathbf{G}_n(z) = G_{n,x}(z) \hat{\mathbf{x}} + G_{n,y}(z) \hat{\mathbf{y}}. \quad (49)$$

The components $G_{n,x}$ and $G_{n,y}$ are chosen such that

$$\begin{aligned} \mathbf{D} \wedge \mathbf{G}_n &\equiv \frac{dG_{n,y}}{dx} - \frac{dG_{n,x}}{dy} \\ &= \tilde{\mathbf{g}}_n(z). \end{aligned} \quad (50)$$

As in Pál (2012) and Luger et al. (2018), the operation $\mathbf{D} \wedge \mathbf{G}_n$ denotes the *exterior derivative* of \mathbf{G}_n . Following Luger et al. (2018), if we choose the following form for Equation (49),

$$\mathbf{G}_n(z) = z^n(-y\hat{\mathbf{x}} + x\hat{\mathbf{y}}), \quad (51)$$

we arrive at the expression presented in Equation (42) for the components of the Green's basis:

$$\begin{aligned} \tilde{\mathbf{g}}_n(z) &= \frac{dG_{ny}}{dx} - \frac{dG_{nx}}{dy} \\ &= (n+2)z^n - nz^{n-2} \end{aligned} \quad (52)$$

for $2 \leq n \leq N$. Note that we already introduced the first two terms, $\tilde{\mathbf{g}}_0 = 1$ (uniform limb darkening) and $\tilde{\mathbf{g}}_1 = z$ (linear limb darkening). Since we already know how to integrate them (§2 and §3), we treat them separately from the higher order terms.

Returning to Equation (48), we note that the line integral consists of two arcs: an arc \mathcal{P} along the boundary of the occulting body and an arc \mathcal{Q} along the boundary of the occulted body. We may therefore write the n^{th} component of the solution vector as

$$\mathbf{s}_n = \mathcal{Q}(\mathbf{G}_n) - \mathcal{P}(\mathbf{G}_n), \quad (53)$$

where, as in Pál (2012) and Luger et al. (2018), we define the *primitive integrals*

$$\mathcal{P}(\mathbf{G}_n) \equiv \int_{\pi-\phi}^{2\pi+\phi} [G_{n,y}(rc_\varphi, b + rs_\varphi)c_\varphi - G_{n,x}(rc_\varphi, b + rs_\varphi)s_\varphi] r d\varphi, \quad (54)$$

taken along the boundary of the occulting body of radius r , and

$$\mathcal{Q}(\mathbf{G}_n) \equiv \int_{\pi-\lambda}^{2\pi+\lambda} [G_{n,y}(c_\varphi, s_\varphi)c_\varphi - G_{n,x}(c_\varphi, s_\varphi)s_\varphi] d\varphi, \quad (55)$$

taken along the boundary of the occulted body of radius unity. For convenience, we defined $c_\varphi \equiv \cos \varphi$ and $s_\varphi \equiv \sin \varphi$ and we used the fact that along the arc of a circle,

$$d\mathbf{r} = -rs_\varphi d\varphi \hat{\mathbf{x}} + rc_\varphi d\varphi \hat{\mathbf{y}}. \quad (56)$$

The angles ϕ and λ are the same as those used in [Luger et al. \(2018\)](#) (see their Figure 2) and are given by $\phi = \kappa_0 - \pi/2$ and $\lambda = \pi/2 - \kappa_1$ (c.f. Equation 7 and Figure 1).

Inserting our expression for \mathbf{G}_n into Equations (54) and (55), we arrive at a fairly simple form for the primitive integrals:

$$\mathcal{P}(\mathbf{G}_n) = \int_{\pi-\phi}^{2\pi+\phi} z^n (r + b \sin \varphi) r d\varphi \quad (57)$$

and

$$\mathcal{Q}(\mathbf{G}_n) = \int_{\pi-\lambda}^{2\pi+\lambda} z^n d\varphi. \quad (58)$$

Conveniently, the primitive integral $\mathcal{Q}(\mathbf{G}_n) = 0$ for all $n > 0$, since $z = 0$ at the boundary of the star. Since we need not compute the integral for $n = 0$, as we already found a solution for uniform limb darkening in §2, our final task is to find the solution to Equation (57).

5.3. Solving the \mathcal{P} integral

The primitive integral $\mathcal{P}(\mathbf{G}_n)$ can be rewritten as

$$\mathcal{P}(\mathbf{G}_n) = \int_{\pi-\phi}^{2\pi+\phi} (1 - r^2 - b^2 - 2brs_\varphi)^{\frac{n}{2}} (r + bs_\varphi) r d\varphi, \quad (59)$$

where $s_\varphi = \sin \varphi$. We make the transformation $\xi = \frac{1}{2}(\varphi - \frac{3\pi}{2})$, yielding

$$\mathcal{P}(\mathbf{G}_n) = 2r(4br)^{\frac{n}{2}} \int_{-\frac{\kappa_0}{2}}^{\frac{\kappa_0}{2}} (k^2 - \sin^2 \xi)^{\frac{n}{2}} (r - b + 2b \sin^2 \xi) d\xi, \quad (60)$$

for $2 \leq n \leq N$, where $\kappa_0 = 2 \sin^{-1} k$ for $k^2 \leq 1$ and $\kappa_0 = \pi$ for $k^2 > 1$. We reuse the value of κ_0 which was computed in the uniform limb darkening case (§2).

We can express $\mathcal{P}(\mathbf{G}_n)$ in terms of a sequence of integrals, $\mathcal{M}_n(r, b)$, given by:

$$\mathcal{M}_n(r, b) = (4br)^{n/2} \int_{-\kappa_0/2}^{\kappa_0/2} (k^2 - \sin^2 \xi)^{\frac{n}{2}} d\xi, \quad (61)$$

in terms of which the primitive integral takes the particular simple form

$$\mathcal{P}(\mathbf{G}_n) = (1 + r^2 - b^2)\mathcal{M}_n - \mathcal{M}_{n+2}. \quad \textcolor{blue}{\square} (62)$$

The integrals \mathcal{M}_n obey straightforward recursion relations

$$\begin{aligned} \mathcal{M}_n = & \frac{1}{n} [2(n-1)(1-r^2-b^2)\mathcal{M}_{n-2} \\ & + (n-2)(1-(b-r)^2)((b+r)^2-1)\mathcal{M}_{n-4}], \end{aligned} \quad \textcolor{blue}{\square} (63)$$

$$\mathcal{M}_n = \frac{(n+4)\mathcal{M}_{n+4} - 2(n+3)(1-r^2-b^2)\mathcal{M}_{n+2}}{(n+2)(1-(b-r)^2)((b+r)^2-1)}, \quad \textcolor{blue}{\square} (64)$$

where the first relation may be used for upwards recursion in n for $k^2 > \frac{1}{2}$, and the second for downward recursion in n otherwise. In practice we replace \mathcal{M}_{n+2} in Equation (62) with the recursion relation to obtain a more stable expression for $\mathcal{P}(\mathbf{G}_n)$:

$$\mathcal{P}(\mathbf{G}_n) = 2r^2\mathcal{M}_n - \frac{n}{n+2} ((1-r^2-b^2)\mathcal{M}_n + (1-(b-r)^2)((b+r)^2-1)\mathcal{M}_{n-2}). \quad \textcolor{blue}{\square} (65)$$

Note that these recursion relations involve every fourth term, so we need to compute the first four terms analytically. These are given by:

$$\begin{aligned}
\mathcal{M}_0 &= \kappa_0, \\
\mathcal{M}_1 &= 2(4br)^{1/2} [E(k^2) - (1 - k^2)K(k^2)], \\
\mathcal{M}_2 &= 4br \left[(k^2 - \tfrac{1}{2})\kappa_0 + k\sqrt{1 - k^2} \right], \\
\mathcal{M}_3 &= \tfrac{2}{3}(4br)^{3/2} [(4k^2 - 2)E(k^2) + (3k^2 - 2)(k^2 - 1)K(k^2)], \tag{66}
\end{aligned}$$

for $k^2 \leq 1$, while for $k^2 > 1$,

$$\begin{aligned}
\mathcal{M}_0 &= \pi, \\
\mathcal{M}_1 &= 2(1 - (r - b)^2)^{1/2} E(k^{-2}), \\
\mathcal{M}_2 &= \pi(1 - b^2 - r^2), \\
\mathcal{M}_3 &= \tfrac{2}{3}(4br)^{3/2} k^3 [2(2 - k^{-2})E(k^{-2}) - (1 - k^{-2})K(k^{-2})]. \tag{67}
\end{aligned}$$

We re-express the elliptic integrals for $n = 1$ and $n = 3$ in terms of the cel integrals which were already computed for the linear limb darkening case,

$$\begin{aligned}
\mathcal{M}_1 &= 2(4br)^{1/2} k^2 \text{cel}(k_c, 1, 1, 0), \\
\mathcal{M}_3 &= \tfrac{2}{3}(4br)^{3/2} k^2 [\text{cel}(k_c, 1, 1, k_c^2) + (3k^2 - 2)\text{cel}(k_c, 1, 1, 0)], \tag{68}
\end{aligned}$$

for $k^2 \leq 1$, while for $k^2 > 1$,

$$\begin{aligned}
\mathcal{M}_1 &= 2(1 - (r - b)^2)^{1/2} \text{cel}(k_c, 1, 1, k_c^2), \\
\mathcal{M}_3 &= \tfrac{2}{3}(1 - (b - r)^2)^{3/2} [(3 - 2k^{-2})\text{cel}(k_c, 1, 1, k_c^2) + k^{-2}\text{cel}(k_c, 1, 1, 0)]. \tag{69}
\end{aligned}$$

where, as before, $k_c = \sqrt{1 - k^2}$ for $k^2 \leq 1$, and $k_c = \sqrt{1 - k^{-2}}$ for $k^2 > 1$.

For downward recursion, we compute the top four \mathcal{M}_n expressions, $N - 3 \leq n \leq N$, in terms of series expansions. When $k^2 \leq 1$, the integrals may be expressed in terms

of the following Hypergeometric functions and infinite series,

$$\begin{aligned}
\mathcal{M}_n &= (4br)^{n/2} k^{n+1} \pi^{1/2} \frac{\Gamma(1 + \frac{n}{2})}{\Gamma(\frac{3}{2} + \frac{n}{2})} {}_2F_1(\frac{1}{2}, \frac{1}{2}; \frac{3}{2} + \frac{n}{2}; k^2), \\
&= (1 - (r - b)^2)^{n/2} k \sum_{j=0}^{j_{max}} \alpha_j k^{2j}, \\
\alpha_0 &= \sqrt{\pi} \frac{\Gamma(1 + \frac{n}{2})}{\Gamma(\frac{3}{2} + \frac{n}{2})}, \\
\alpha_j &= \alpha_{j-1} \frac{(2j - 1)^2}{2j(1 + n + 2j)}. \tag{70}
\end{aligned}$$

Although $j_{max} = \infty$, in practice we set $j_{max} = 100$, and the series is truncated when a term goes below a tolerance specified by the numerical precision.

We find that upward recursion in n is more stable for $k^2 > \frac{1}{2}$, while downward recursion is more stable for $k^2 < \frac{1}{2}$. Note that this differs from [Luger et al. \(2018\)](#), for which downward recursion was also required for $k^2 > 2$.

5.4. Analytic derivatives

The derivatives of $\mathcal{P}(\mathbf{G}_n)$ may be expressed simply as functions of \mathcal{M}_n :

$$\frac{\partial \mathcal{P}}{\partial r} = 2r [(n + 2)\mathcal{M}_n - n\mathcal{M}_{n-2}], \tag{71}$$

$$\frac{\partial \mathcal{P}}{\partial b} = \frac{n}{b} [(r^2 + b^2)(\mathcal{M}_n - \mathcal{M}_{n-2}) + (r^2 - b^2)^2 \mathcal{M}_{n-2}]. \tag{72}$$

We find that the derivative with respect to impact parameter becomes numerically unstable for small values of b due to cancellation between the two terms, followed by division by b . To avoid this problem for small b , we have derived an alternative expression which we utilize when $b < b_c$, where b_c is a (small) cutoff value, which avoids division by b :

$$\frac{\partial \mathcal{P}}{\partial b} = n [b\mathcal{M}_n + (2r^3 + b^3 - 3r^2b - b - 3)\mathcal{M}_{n-2} - 4r^3\mathcal{N}_{n-2}], \tag{73}$$

where we have defined a new integral, \mathcal{N}_n ,

$$\mathcal{N}_n(r, b) = (4br)^{n/2} \int_{-\kappa_0/2}^{\kappa_0/2} (k^2 - \sin^2 \xi)^{\frac{n}{2}} \sin^2 \xi d\xi, \tag{74}$$

which obeys the recursion relation

$$\mathcal{N}_n = \frac{1}{n+2} [\mathcal{M}_n + n(1 - (b+r)^2)\mathcal{N}_{n-2}]. \quad (75)$$

Since this recursion relation involves every other term, we only need the two lowest terms, which are given by:

$$\begin{aligned} \mathcal{N}_0 &= \frac{1}{2}\kappa_0 - kk_c, \\ \mathcal{N}_1 &= \frac{2}{3}(4br)^{1/2}k^2 [-\text{cel}(k_c, 1, 1, k_c^2) + 2\text{cel}(k_c, 1, 1, 0)], \end{aligned} \quad (76)$$

for $k^2 \leq 1$ and

$$\begin{aligned} \mathcal{N}_0 &= \frac{\pi}{2}, \\ \mathcal{N}_1 &= \frac{2}{3}(4br)^{1/2}k [2\text{cel}(k_c, 1, 1, k_c^2) - \text{cel}(k_c, 1, 1, 0)], \end{aligned} \quad (77)$$

for $k^2 > 1$.

In the $k^2 < \frac{1}{2}$ limit, we find the upward recursion to be unstable, and so we evaluate the expressions for \mathcal{N}_N and \mathcal{N}_{N-1} with a series solution (as we did for $\mathcal{M}_{N-3}, \dots, \mathcal{M}_N$):

$$\begin{aligned} \mathcal{N}_n &= (4br)^{n/2}k^{n+3} \frac{\pi^{1/2}}{2} \frac{\Gamma(1 + \frac{n}{2})}{\Gamma(\frac{5}{2} + \frac{n}{2})} {}_2F_1(\frac{1}{2}, \frac{3}{2}; \frac{5}{2} + \frac{n}{2}; k^2) \\ &= (1 - (r-b)^2)^{n/2}k^3 \sum_{j=0}^{j_{max}} \gamma_j k^{2j}, \\ \gamma_0 &= \frac{\sqrt{\pi}}{2} \frac{\Gamma(1 + \frac{n}{2})}{\Gamma(\frac{5}{2} + \frac{n}{2})}, \\ \gamma_j &= \gamma_{j-1} \frac{(4j^2 - 1)}{2j(3 + n + 2j)}. \end{aligned} \quad (78)$$

We then use downward recursion with the relation

$$\mathcal{N}_n = \frac{(n+4)\mathcal{N}_{n+2} - \mathcal{M}_{n+2}}{(n+2)(1 - (b+r)^2)} \quad (79)$$

to iterate down to $n = 3$, while finally computing $n = 1$ and $n = 2$ exactly.

Evaluating this additional integral adds further computational expense, but in practice we only need to compute it for $b < b_c = 10^{-3}$ to obtain similar accuracy to

the other expressions. This is encountered rarely as it only applies when the occulter is nearly aligned with the source.

With the computation of $\mathfrak{s}_n = -\mathcal{P}(\mathbf{G}_n)$, we then compute the derivatives of the solution vector as

$$\frac{\partial \mathfrak{s}_n}{\partial r} = -\frac{\partial \mathcal{P}(\mathbf{G}_n)}{\partial r}, \quad (80)$$

$$\frac{\partial \mathfrak{s}_n}{\partial b} = -\frac{\partial \mathcal{P}(\mathbf{G}_n)}{\partial b}, \quad (81)$$

for $2 \leq n \leq N$, while the $n = 0$ and $n = 1$ terms are handled separately as in section §3. The derivatives of the normalized flux, F , with respect to r and b are then computed as

$$\frac{\partial F}{\partial r} = I_0 \sum_{n=0}^N \mathfrak{g}_n \frac{\partial \mathfrak{s}_n}{\partial r}, \quad (82)$$

$$\frac{\partial F}{\partial b} = I_0 \sum_{n=0}^N \mathfrak{g}_n \frac{\partial \mathfrak{s}_n}{\partial b}. \quad (83)$$

We will see in the following section that the normalization constant, I_0 , is independent of \mathfrak{g} for $n \geq 2$, so the derivative of F with respect to \mathfrak{g} is trivial:

$$\frac{\partial F}{\partial \mathfrak{g}_n} = I_0 \mathfrak{s}_n \quad (84)$$

for $n \geq 2$. For the first two terms, we differentiate Equation (94) in the following section to obtain

$$\frac{\partial F}{\partial \mathfrak{g}_0} = I_0 \mathfrak{s}_0 - \pi \frac{F}{I_0^2}, \quad (85)$$

$$\frac{\partial F}{\partial \mathfrak{g}_1} = I_0 \mathfrak{s}_1 - \frac{2\pi}{3} \frac{F}{I_0^2}. \quad (86)$$

The derivatives of the coefficients, $\frac{\partial \mathfrak{g}_j}{\partial u_i}$, are computed by differentiating each term in Equation (38) to obtain $\frac{\partial \mathfrak{p}_n}{\partial u_i}$, and differentiating each term in Equation (43) to obtain $\frac{\partial \mathfrak{g}_j}{\partial \mathfrak{p}_n}$, and propagating the derivatives via the chain rule. Then, the light curve derivatives are given by

$$\frac{\partial F}{\partial u_i} = \sum_j \frac{\partial \mathfrak{g}_j}{\partial u_i} \frac{\partial F}{\partial \mathfrak{g}_j}. \quad (87)$$

5.5. Normalization

We now know how to compute the flux and all its derivatives during an occultation, but only up to an unknown normalization constant I_0 . It is convenient to choose a normalization such that the total unocculted flux is unity, regardless of the value of the limb darkening coefficients. We therefore require that

$$\begin{aligned} F &= \oint\!\!\!\oint I(z) \, dS \\ &= I_0 \mathfrak{s}^\top \mathcal{A} \mathbf{u} \\ &= 1 \end{aligned} \tag{88}$$

when the integral is taken over the entire disk of the body. We must thus have

$$\begin{aligned} I_0 &= \frac{1}{\mathfrak{s}^\top(r=0) \mathcal{A} \mathbf{u}} \\ &= \frac{1}{\mathfrak{s}^\top(r=0) \mathfrak{g}}, \end{aligned} \tag{89}$$

where $\mathfrak{s}^\top(r=0)$ is the solution vector when there is no occulter (i.e., $r=0$). We could compute I_0 using the formalism from the previous section, but it is easier to evaluate the integral directly. When there is no occulter, the n^{th} term of \mathfrak{s} corresponds to the double integral in polar coordinates

$$\begin{aligned} \mathfrak{s}_n(r=0) &= \int_0^{2\pi} \int_0^1 \tilde{\mathfrak{g}}_n(z) r \, dr \, d\theta \\ &= 2\pi \int_0^1 \tilde{\mathfrak{g}}_n(z) r \, dr, \end{aligned} \tag{90}$$

From Equation (42), we may write

$$\mathfrak{s}_n(r=0) = 2\pi \begin{cases} \int_0^1 r \, dr & n=0 \\ \int_0^1 z r \, dr & n=1 \\ (n+2) \int_0^1 z^n r \, dr - n \int_0^1 z^{n-2} r \, dr & n \geq 2. \end{cases} \tag{91}$$

The first case is trivial and integrates to $\mathfrak{s}_0(r=0) = \pi$. The remaining cases involve integrands of the form $z^n r$, where $z = \sqrt{1-r^2}$. We may evaluate these integrals by

substituting $u = z^2 = 1 - r^2$ and $du = -2r dr$:

$$\begin{aligned} \int_0^1 z^n r dr &= \frac{1}{2} \int_0^1 u^{\frac{n}{2}} du \\ &= \frac{1}{2+n}. \end{aligned} \quad (92)$$

The solution vector then simplifies to

$$\mathfrak{s}_n(r=0) = \begin{cases} \pi & n=0 \\ \frac{2\pi}{3} & n=1 \\ 0 & n \geq 2. \end{cases} \quad (93)$$

Interestingly, the net flux contribution for all terms in the Green's basis with $n \geq 2$ is exactly zero. We may finally evaluate our normalization constant:

$$I_0 = \frac{1}{\pi(\mathfrak{g}_0 + \frac{2}{3}\mathfrak{g}_1)}. \quad (94)$$

5.6. Summary

In this section, we showed that if we express the specific intensity distribution on the surface of a spherical body as the series

$$\frac{I(\boldsymbol{\mu})}{I_0} = 1 - u_1(1 - \boldsymbol{\mu}) - u_2(1 - \boldsymbol{\mu})^2 - \dots - u_N(1 - \boldsymbol{\mu})^N, \quad (95)$$

(see Equation 31), the total flux observed during an occultation is given by the analytic and closed form expression

$$F = I_0 \mathfrak{s}^\top \mathcal{A} \mathbf{u}, \quad (96)$$

where I_0 is a normalizing constant (Equation 94), \mathfrak{s} is the solution vector (Equation 53), a function of only the impact parameter b and radius r of the occulter, \mathcal{A} is a change of basis matrix (Equation 45), and \mathbf{u} is the vector of limb darkening coefficients $(u_0 \ u_1 \ u_2 \ \dots \ u_N)^\top$.

Rodrigo: Write a slightly more detailed summary. Emphasize how the uniform and linear terms are computed separately.

With the description of the light curve computation complete, we next discuss the integration of the light curve over a finite time step.

6. TIME INTEGRATION

Given that most observations are made over a finite exposure time, the integration of the light curve over time is necessary for representing a transit light curve with high fidelity (Kipping 2010). For constructing a light curve, usually one divides the fluence by the integration time to obtain the time-averaged flux. For optimizing and inferring the posterior of model parameters, we would like to compute the derivatives of the time-averaged flux with respect to the model parameters.

The instantaneous flux is a function of $3 + n$ parameters in the Green's basis, $\{r, b, \mathbf{g}_n\}$, or $2 + n$ parameters in the polynomial in $1 - \mu$ basis, and of these, only one varies with time, $b(t)$. Thus, we can compute the time-dependent flux with a model for $b(t) = b(\mathbf{x}, t)$, where $b(\mathbf{x}, t)$ is a model for the impact parameter as a function of time and model parameters \mathbf{x} . The set of model parameters need to be specified by a function, which, for example, might be a Keplerian orbit of the two bodies with respect to one another, or a full dynamical model of an n-body system. To compute the derivatives of the light curve with respect to \mathbf{x} , the derivatives of the dynamical model must be computed as well.

The time-averaged flux, \bar{F} , and its derivatives, are given by

$$\bar{F} = \frac{1}{\Delta t} \int_{t-\frac{1}{2}\Delta t}^{t+\frac{1}{2}\Delta t} F(t') dt', \quad (97)$$

$$\frac{\partial \bar{F}}{\partial r} = \frac{1}{\Delta t} \int_{t-\frac{1}{2}\Delta t}^{t+\frac{1}{2}\Delta t} \frac{\partial F(t')}{\partial r} dt', \quad (98)$$

$$\frac{\partial \bar{F}}{\partial \mathbf{g}_i} = \frac{1}{\Delta t} \int_{t-\frac{1}{2}\Delta t}^{t+\frac{1}{2}\Delta t} \frac{\partial F(t')}{\partial \mathbf{g}_i} dt', \quad (99)$$

$$\frac{\partial \bar{F}}{\partial \mathbf{x}} = \frac{1}{\Delta t} \int_{t-\frac{1}{2}\Delta t}^{t+\frac{1}{2}\Delta t} \frac{\partial F}{\partial b} \frac{\partial b(t')}{\partial \mathbf{x}} dt', \quad (100)$$

where t is taken to be the mid-point of the transit exposure time, and Δt is the exposure time.

As an example, we choose the approximate transit model $b(t) = (b_0^2 + v^2(t - t_0)^2)^{1/2}$, which ignores acceleration and curvature during a transit, and thus is valid in the limit of large orbital separation. We compute the time-averaged flux and derivatives with respect to $\mathbf{x} = \{t_0, v, b_0\}$ for a length of integration time Δt . **Eric: Update this description of the time-integration.** The integration is carried out with an adaptive refinement of the exposure time interval by factors of two until the midpoint in each refined interval is within $10^{-6}r^2$ of the mean of the start and end with a maximum number of refinement depths of 32. Figure 7 shows a comparison of the derivatives of the time-integrated light curve for this impact parameter model.





Figure 7. Comparison of the normalized flux and its derivatives with (solid) and without (dashed) time-integration. The integration time is indicated in the upper left panel with a horizontal blue line. The derivatives are computed with respect to $\{r, t_0, v, b_0, u_1, u_2\}$. The parameters are given by $r = 0.1$, $t_0 = 0$, $b_0 = 0.5$, $u_1 = u_2 = 0.3$. 

The time-integration smooths both the features of the light curve, as well as the features of the derivative curves. The similarity of the shape of the derivative with respect to v and b_0 makes it apparent that there may be partial degeneracies between the impact parameter and duration of a transit, which can make the impact parameter more difficult to measure, especially when the exposure time is longer than the time of ingress/egress. Likewise, the derivatives with respect to the two limb darkening parameters have a similar shape, which explains why in some cases it can be difficult to constrain both parameters.

This completes the description of the light curve computation, along with its derivatives. We now turn to discussing the implementation of the computation, followed by comparison with existing codes.

7. IMPLEMENTATION DETAILS

There are several details in our implementation of the foregoing equations which give further speedup of the computation, which we describe in this section.

When a light curve is computed, there are some computations which only need to be carried out once, and then reused at each time step in the light curve computation. We define a structure to hold the variables which are reused throughout the light curve. In addition, due to the greater computational expense of square-roots and divisions,

where possible we try to only compute a square root or division once, storing these in a variable within the structure, and then reusing these with cheaper multiplication throughout the computation when needed. For instance, for many formulae we require the inverse of an integer, so an array of integer inverses is computed once and stored, and then accessed as required rather than recomputed.

Once the number of limb darkening terms, N , is specified, then the series coefficients for \mathcal{M}_n and \mathcal{N}_n , α_j and γ_j , are a simple function of j and n , and so we compute these coefficients once, and store them in a vector for $k^2 \leq 1$, separately for $N - 3$ to N for \mathcal{M}_n , and for $N - 1$ and N for \mathcal{N}_n .

In addition, once N is specified, then the transformation matrix for the Jacobian from \mathbf{g}_i to u_j , $\frac{\partial \mathbf{g}_i}{\partial u_j}$, remains the same throughout the light curve computation, so we compute this matrix only once, and then compute the flux derivative (Equation 87) with matrix multiplication. In fact, since the Jacobian matrix for transforming the derivatives from \mathbf{g}_i to u_j can be expensive to apply, we can carry out the gradient of the likelihood function with respect to \mathbf{g}_i , and then apply the Jacobian transformation from \mathbf{g}_i to u_j only once to obtain the gradient of the likelihood with respect to the limb darkening parameterization. In practice, we are usually only concerned with optimizing a likelihood or computing gradients of a likelihood for Hamiltonian Markov Chain Monte Carlo, so the derivatives of the particular points in the light curve with respect to u_i aren't needed. This results in a significant computational savings, especially for large N . Transformation to other parameterizations, such as q_1 and q_2 defined by [Kipping \(2013\)](#), may also be accomplished after the fact by applying the Jacobian to compute the gradient of the likelihood in terms of these transformed parameters.

In computing the elliptic integrals, `cel`, we found that several terms which appear in the Barty formalism are repeated amongst all three elliptic integrals which appear in the expressions for \mathfrak{s}_1 . Consequently, we carry out a parallel computation of these elliptic integrals such that these repeated terms are only computed once; this improves the efficiency of the elliptic integral computation. Once these elliptic integrals are computed for \mathfrak{s}_1 , the elliptic integrals $\text{cel}(k_c, 1, 1, k_c^2) = E(m_k)$ and $\text{cel}(k_c, 1, 1, 0) = (E(m_k) - (1 - m_k)K(m_k))/m_k$ are stored in the structure and reused for computing \mathcal{M}_n and \mathcal{N}_n .

We find that the precision of the computation begins to degrade for $N \approx 25 - 30$. For large values of N , the computing time for propagating the derivatives from \mathbf{g}_i to u_i scales as N^2 , and thus can dominate the computation time. We utilize the BLAS linear algebra library (specifically, the routine `gemv`) for efficient multiplication of this matrix times the derivative of the light curve with respect to \mathbf{g}_i in order to obtain the derivatives with respect to u_i . However, as we note above, an even more efficient approach is to simply apply this transformation once to the gradient of the likelihood.

8. BENCHMARKING

We have measured the performance of the limb-darkened light curves with derivatives as a function of the number of computed data points and as a function of the number of limb darkening coefficients. We have computed the timing for $r = 0.1$ and for a number of impact parameters ranging from 10^2 to 10^6 , and the number of limb darkening coefficients ranging from 1 to 144. For each set of timing benchmark parameters, we carried out nine measurements of the timing, and we use the median of these for plotting purposes. The benchmarking for the `Julia` code was carried out with `v0.7` of Julia on a MacBook Pro with 2.8 GHz Intel Core i7. No sub-sampling was carried out in this computation. **Rodrigo:** Update this sentence with the Travis CPU specs.

Figure 8 shows that the time dependence is linear with the number of b values (which is equivalent to the number of data points in the light curve). The linear scaling with time holds for each value of the number of limb darkening coefficients.

Figure 9 shows that the time dependence scales approximately as $N^{0.2-1}$. As with the number of light-curve points, we have taken the median over nine measurements for each set of parameters. We then scaled the timing to the single-coefficient case, and took a second median over the number of light curve points as the cube-root scaling scales about the same with different numbers of points in the light curve.

9. NON-LINEAR LIMB DARKENING

Claret (2000) introduced a “non-linear” limb darkening model which was found to be an effective model for describing the limb darkening functions which are produced by models of stellar atmospheres. Although we can only model limb darkening which is integer powers of μ , we can use the polynomial model as an alternative limb darkening model.

We have computed an example non-linear light curve with $r = 0.1$ and $c_1 = c_2 = c_3 = c_4 = 0.2$, and then fit it with successive orders of the polynomial approximation. The non-linear light curve model we computed numerically as the analytic expressions in Mandel & Agol (2002) are in terms of hypergeometric functions which are expensive to evaluate. We compute the light curve with a “layer-cake” model in which sums of layers of surface brightness with a grid of increasing radii are added together to approximate the lightcurve; this is the approach taken in the numerical model used to compute the non-linear limb darkening light curves in the code of Mandel & Agol (2002), and it is analagous to the approach taken by Kreidberg (2015) for computing models with arbitrary limb darkening profiles.

We find that the fit improves steadily up until $N = 6$ (a sextic polynomial), while beyond sextic, the RMS improves imperceptibly. The RMS of the sextic fit for this example is $< 5 \times 10^{-7}$ relative to a depth of transit of about 1.4% (Figure 10).

10. COMPARISON WITH PRIOR WORK

In this section we compare our computations with existing code in terms of accuracy and speed.




Figure 8. Scaling of the computation time in seconds with the number of data points in the light curve for $r = 0.1$ with b ranging from 0 to 1.2, and with the number of limb darkening coefficients, N . [\[46\]](#)

10.1. Comparison with Mandel & Agol

For uniform, linear or quadratic limb darkening, the widely used computation by Mandel & Agol (2002) has been improved in speed by utilizing the Bulirsch (1965a,b) expressions for the complete elliptic integral of the third kind which is needed for the linear case, as implemented in the EXOFAST routines (Eastman et al. 2013). It also uses a series approximation for the complete elliptic integrals of the first and second kind (Hastings 1955). These three elliptic integrals, especially the third kind, are the bottleneck in the computation, and the Bulirsch version is faster than widely used Carlson implementation of elliptic integrals (Carlson 1979).

We have carried out a numerical comparison of the Mandel & Agol (2002) for the linear case ($u_1 = 1$), and find that the most severe errors occur for $b = r \pm \epsilon$. Figure 11 shows the computed models and the errors as a function of r for $b = 1 - r - \epsilon$ and $b = r - \epsilon$, with $\epsilon = 10^{-12}$ (the results look very similar with $+\epsilon$, so we have only plotted one case for clarity). In the $b \approx 1 - r$ case (near second and third contacts), the errors are larger than our new expression, reaching $\approx 10^{-10}$ for $r = 1$. However, the errors become much more severe in the $b \approx r$ case. For $b = r \pm 10^{-12}$, the errors




Figure 9. Scaling of the computation time with the number of limb darkening coefficients, N . The y -axis scales the timing with respect to the timing for a single limb darkening coefficient. [\[4\]](#)

grow to 10^{-4} , and continue to grow as b gets closer to r . No such instability occurs for our new expressions, demonstrating their utility in all regions of parameter space.

A speed comparison for a transit computed with 10^7 data points shows that the Julia implementation of these routines takes about 55% of the CPU time as the IDL EXOFAST implementation without derivatives, and about 65% of the computation time when including the derivatives. Consequently, we conclude that our new implementation is both faster (by 35-45%) and more accurate than the Mandel & Agol IDL implementation.

10.2. Derivative comparison with Pál

We have computed the quadratic limb-darkened light curve using the F77 code written by András Pál, `ntiq_fortran.f`. Figure 12 shows the results of this comparison. The light curve models agree quite well, as do the derivatives, which is a good check on both codes. However, we find that the Pál model only achieves single precision for the computation, with errors reaching as much as a few $\times 10^{-8}$ for the flux and the derivatives with respect to the limb darkening parameters. Pál (2008) uses the Carlson implementation of elliptic integrals (Carlson 1979), which in practice




Figure 10. Comparison of the non-linear limb darkening with polynomial fits of various orders. [🔗](#)

we find can be both less precise and slower to evaluate than the [Bulirsch \(1965a\)](#) code for computing elliptic integrals.

We have also compared the evaluation speed of our code with Pál’s as well. We compiled Pál’s code using `gfortran -O3`, and found that the computation of quadratic limb-darkened light curves and derivatives takes an average of 0.52 seconds to compute 10^6 models, while the `transit_poly_struct.jl` takes an average of 0.16 seconds, giving our Julia code a 70% speed advantage over the Fortran code.

10.3. Comparison to *batman*

A Python implementation of transit light curves which has been widely applied is the `batman` package by [Kreidberg \(2015\)](#). This package implements a fast C++ version of the computation, called by Python for ease of use.




Figure 11. Comparison of Mandel & Agol (2002) with Agol & Luger (2018). [📄](#)

The `batman` package computes the quadratic limb darkening model of Mandel & Agol (2002), and uses the same approach for computing the complete elliptic integrals as EXOFAST. We have made a comparison of our implementation of quadratic limb darkening with `batman`, which is shown in Figure 13. Without computing derivatives, our approach take about 60% of the time of `batman`; with derivatives, the two are comparable in speed.

Next, we ran a comparison with the non-linear limb darkening model which proves to be a better fit than the quadratic model to both simulated and observed stellar atmospheres. The `batman` code carries out a numerical integration over the surface brightness as a function of radius over the stellar disk, which requires additional computational time and limits the precision. We have carried out a fit to the non-linear limb darkening profile with $c_1 = c_2 = c_3 = c_4 = 0.2$ with a polynomial limb darkening model with $N = 15$. We then ran a timing comparison between the `batman`




Figure 12. Comparison of Pál (2008) with Agol & Luger (2018). The coefficients are $u_1 = 0.2$ and $u_2 = 0.3$. 




Figure 13. Comparison of Kreidberg (2015) with Agol & Luger (2018) for a transit across a quadratically limb-darkened star. [\[4\]](#)




Figure 14. Comparison of Kreidberg (2015) with Agol & Luger (2018) for a transit across a nonlinearly limb-darkened star. [\[4\]](#)

model and the polynomial model, and we find that the polynomial model is about 7 times as accurate and 25 times faster to evaluate compared with *batman*.




Figure 15. Comparison of Giménez (2006) with Agol & Luger (2018). [📄](#)

10.4. Comparison to *PyTransit*

Rodrigo: Describe comparison with the Gimenez computation.

11. DISCUSSION

We have presented formulae for the transit (or occultation/eclipse) of a limb-darkened body with a limb darkening profile which is given by a polynomial in μ . These formulae have multiple assumptions built in: both bodies are treated as spherical (but see Seager & Hui 2002; Hui & Seager 2002), so that their projected surfaces are assumed to be circular (but see Barnes & Fortney 2003, 2004; Barnes et al. 2009; Dobbs-Dixon et al. 2012); limb darkening is treated as azimuthally-symmetric (but see Barnes 2009); refraction and any relativistic effects are ignored (but see Sidis & Sari 2010); and the edges of both bodies are assumed to have a sharp boundary. All of these assumptions are violated in every transit event to some extent, but in the majority of cases these assumptions can yield a precise model for a given signal-to-noise ratio.

Given these assumptions, generally one next assumes a particular functional form for the limb darkening law (Csizmadia 2018). The parameterization of the limb darkening model can impact the precision of the computation of transit light curves. A common approach is to derive limb darkening coefficients for a particular limb darkening model from stellar atmosphere models, and to either fix these at the tabulated values given an observing band and an estimate of stellar parameters (Claret & Bloemen 2011; Howarth 2011), or at least to place a prior that the limb darkening parameters should nearly match these values. This approach can have several

pitfalls: the limb darkening model may not be sufficiently precise, the stellar atmosphere model may not be accurate, and the stellar parameters may not be precise. In computing limb darkening from stellar atmosphere models, the spherical nature of limb darkening can affect the transit light curve (Neilson & Lester 2013; Neilson et al. 2017), and thus the limb darkening coefficients must be fit with care (Claret 2018). Even more importantly, full three dimensional stellar atmosphere models appear to give a more accurate description of stellar limb darkening by capturing the structure of the atmosphere under the influence of granulation (Hayek et al. 2012; Magic et al. 2015). However, any physical model for a stellar atmosphere has limitations in the fidelity at which it can model actual stellar atmospheres, and any modeler can only explore a finite set of parameters (effective temperature, metallicity, surface gravity, and magnetic field strength). In practice, then, it may be most robust simply to let the limb darkening parameters be free parameters, to let the limb darkening model be as flexible as possible, and to let the limb darkening model be fit along with the radius ratio and orbital parameters (Csizmadia et al. 2012; Espinoza & Jordán 2015).

Even so, this approach still assumes azimuthal symmetry for the star, while any model for the surface brightness of a star can only be approximate: most stars are convective, rotationally-oblate, spotted, oscillating, flaring, etc. The model we have presented, then, will only resemble any given star to a precision which is limited by the lack of uniformity of the actual stellar surface. This begs the question of why a numerically precise model is required for modelling transit light curves. The answer is computational accuracy and stability: this more accurate model can be used over all of parameter space, without returning spurious results, and the high precision enables computation of derivatives which are beneficial when optimizing model parameters, computing the Fisher information matrix, or deriving parameter posteriors with MCMC.

Since we are limited in the knowledge of the properties of any given star, the discrepancies of an azimuthally-symmetric limb-darkened model can be treated as a source of noise. The deviation of the star from the model can be absorbed into noise models that account for outliers, account for correlations in the noise, or actually try to model the deviations of the star from azimuthal symmetry, such as induced by star spots (e.g. Sanchis-Ojeda & Winn 2011).

One question is what order of the limb darkening model to choose to fit the data? Here we suggest several possible solutions. The order of the limb darkening can be varied until the chi-square no longer improves (subject to a penalty for the greater freedom in the model, such as Bayesian Information Criterion). A high-order limb darkening model can be chosen, with the coefficients regularized to favor small values; should the data require a higher-order model, then the coefficients will increase to accommodate the data. The parameterization of the limb darkening with terms with $\mathbf{g}_n((n+2)\mu^n - \mu^{n-2})$ for $2 \leq n \leq N$ may be particularly convenient for this model in that these terms do not contribute to the total flux of the star. A third possibility

is to fit stellar atmosphere models with the polynomial limb darkening model until a sufficient precision is reached given that warranted by the data, and then to place priors on the limb darkening parameters, informed by the stellar limb darkening models. A limitation of our computational approach is that the precision begins to degrade for $N \approx 25 - 30$; however, we anticipate that such a high order will rarely be required.

12. APPLICATIONS

We envision that this code will be used for fits to higher precision transit data, such as gathered by the James Webb Space Telescope (JWST), which require an improved model of stellar limb darkening. Here we discuss some potential avenues for application of this model.

The derivatives of the time-integrated light curves may be used to revisit the Fisher information analysis as carried out by [Price & Rogers \(2014\)](#), as originally investigated without time-integration by [Carter et al. \(2008\)](#). Accounting for correlated noise in this analysis will give more plausible estimates for the impact of stellar variability on the determination of transit transmission spectroscopy and transit-timing variations ([Foreman-Mackey et al. 2017](#)). This limit will be encountered as more precise measurements are made by gathering more photons during a transit. For example, for some targets, one can expect to obtain $\sim 10^2$ times as many photons with JWST as collected with Kepler. With such higher precision, as well as the wavelength-dependence afforded by several JWST observing modes, one can expect that high fidelity transit models will be required for making precise measurements of transit parameters.

The detection of transit-timing variations with low-amplitude sinusoidal variations can make use of the fact that small variations in transit time can be expanded as a Taylor series to linear order in time so that perturbations in the transit time are the sum of a periodic component and a constant times the derivative of the limb-darkened light curve ([Ofir et al. 2018](#)). This approach requires derivatives of the light curve with respect to time, for which the [Mandel & Agol \(2002\)](#) computation is too imprecise near the points of contact, $b \approx r$ and $b \approx 1 - r$, within an impact parameter distance of 10^{-4} , as shown by [Ofir et al. \(2018\)](#). They extrapolated over these regions with polynomials. However, our new precise formulae, with derivatives, will be useful for the perturbative approach to the detection of transit timing variations, avoiding the numerical errors inherent in the [Mandel & Agol \(2002\)](#) model.

13. CONCLUSIONS

We have presented an analytic model for the transits, occultations, and eclipses of limb-darkened bodies with a polynomial dependence of the limb darkening on the z component of the stellar surface (or, alternatively, the cosine of the angle from the sub-stellar point, μ). The model is more precise and accurate than prior models that we have compared to, especially near special limits such as the points of contact and

the coincidence of the edge of the occulter with the center of the source. The model also compares favorably in speed of evaluation, about a factor of three faster than the code due to Pál (2008), XXX faster than Gimenez, a factor of 2-25 faster than *batman* (depending on the order of the limb darkening), and 35-45% faster than EXOFAST.

We expect that this code may be used both as a workhorse model for general fitting of transit models, as well as a tool for more specialized applications, such as photodynamical modeling of interacting planets (Carter et al. 2012), triple stars (Carter et al. 2011), and transiting circumbinary planets (Doyle et al. 2011).

The code is open source, and has two versions: one of which is a part of the *starry* package, <http://github.com/rodluger/starry/>, written in a combination of C++ and Python, and a new code written in Julia as part of the development of the equations in this paper, <http://github.com/rodluger/limbdark/>. We welcome usage of these codes, and contributions to further develop and enhance their capabilities.

We thank András Pál for sharing his Fortran code, *ntiq-fortran.f*. We thank András Pál, Kevin Stevenson, Kai Ueltzhöffer, Mario Damasso, Matthew Heising, Robert Morehead, and Laura Kreidberg for pointing out errors or inaccuracies in the Mandel & Agol paper and code, which we have hopefully rectified in this paper. EA acknowledges NSF grant AST-1615315, NASA grant NNX13AF62G, and from the NASA Astrobiology Institute’s Virtual Planetary Laboratory Lead Team, funded through the NASA Astrobiology Institute under solicitation NNH12ZDA002C and Cooperative Agreement Number NNA13AA93A.

REFERENCES

- | | |
|--|--|
| <p>Abramowitz, M., & Stegun, I. A. 1970, Handbook of Mathematical Functions: With Formulas, Graphs, and Mathematical Tables (Washington, D.C.: U.S. Dept. of Commerce, National Bureau of Standards)</p> <p>Barnes, J. W. 2009, The Astrophysical Journal, 705, 683. https://doi.org/10.1088/0004-637x/705/1/683</p> <p>Barnes, J. W., et al. 2009, The Astrophysical Journal, 706, 877. https://doi.org/10.1088/0004-637x/706/1/877</p> <p>Barnes, J. W., & Fortney, J. J. 2003, The Astrophysical Journal, 588, 545. https://doi.org/10.1086/373893</p> <p>—. 2004, The Astrophysical Journal, 616, 1193. https://doi.org/10.1086/425067</p> | <p>Bartky, W. 1938, Reviews of Modern Physics, 10, 264. https://doi.org/10.1103/revmodphys.10.264</p> <p>Betancourt, M. 2017, arXiv preprint arXiv:1701.02434</p> <p>Bulirsch, R. 1965a, Numerische Mathematik, 7, 78. https://doi.org/10.1007/bf01397975</p> <p>—. 1965b, Numerische Mathematik, 7, 353. https://doi.org/10.1007/bf01436529</p> <p>—. 1969, Numerische Mathematik, 13, 305. https://doi.org/10.1007/bf02165405</p> <p>Carlson, B. C. 1979, Numerische Mathematik, 33, 1. https://doi.org/10.1007/bf01396491</p> <p>Carter, J. A., et al. 2008, The Astrophysical Journal, 689, 499. https://doi.org/10.1086/592321</p> |
|--|--|

- . 2011, *Science*, 331, 562. <https://doi.org/10.1126/science.1201274>
- . 2012, *Science*, 337, 556. <https://doi.org/10.1126/science.1223269>
- Claret, A. 2000, *A&A*, 363, 1081
- . 2018, ArXiv e-prints, arXiv:1804.10135
- Claret, A., & Bloemen, S. 2011, *Astronomy & Astrophysics*, 529, A75. <https://doi.org/10.1051/0004-6361/201116451>
- Csizmadia, S. 2018, in *Handbook of Exoplanets* (Springer International Publishing), 1–15. https://doi.org/10.1007/978-3-319-30648-3_41-1
- Csizmadia, S., et al. 2012, *Astronomy & Astrophysics*, 549, A9. <https://doi.org/10.1051/0004-6361/201219888>
- Dobbs-Dixon, I., et al. 2012, *The Astrophysical Journal*, 751, 87. <https://doi.org/10.1088/0004-637x/751/2/87>
- Doyle, L. R., et al. 2011, *Science*, 333, 1602. <https://doi.org/10.1126/science.1210923>
- Eastman, J., et al. 2013, *Publications of the Astronomical Society of the Pacific*, 125, 83. <https://doi.org/10.1086/669497>
- Espinoza, N., & Jordán, A. 2015, *Monthly Notices of the Royal Astronomical Society*, 450, 1879. <https://doi.org/10.1093/mnras/stv744>
- Feroz, F., & Hobson, M. P. 2008, *Monthly Notices of the Royal Astronomical Society*, 384, 449. <https://doi.org/10.1111/j.1365-2966.2007.12353.x>
- Ford, E. B. 2005, *The Astronomical Journal*, 129, 1706. <https://doi.org/10.1086/427962>
- . 2006, *The Astrophysical Journal*, 642, 505. <https://doi.org/10.1086/500802>
- Foreman-Mackey, D., et al. 2017, *The Astronomical Journal*, 154, 220. <https://doi.org/10.3847/1538-3881/aa9332>
- Giménez, A. 2006, *Astronomy & Astrophysics*, 450, 1231. <https://doi.org/10.1051/0004-6361:20054445>
- Girolami, M., & Calderhead, B. 2011, *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 73, 123. <https://doi.org/10.1111/j.1467-9868.2010.00765.x>
- Goldberg, D. 1991, *ACM Computing Surveys*, 23, 5. <https://doi.org/10.1145/103162.103163>
- Hastings, C. 1955, *Approximations for digital computers*, Princeton, N.J.: Princeton University Press
- Hayek, W., et al. 2012, *Astronomy & Astrophysics*, 539, A102. <https://doi.org/10.1051/0004-6361/201117868>
- Hestroffer, D. 1997, *A&A*, 327, 199
- Howarth, I. D. 2011, *Monthly Notices of the Royal Astronomical Society*, 413, 1515. <https://doi.org/10.1111/j.1365-2966.2011.18122.x>
- Hui, L., & Seager, S. 2002, *The Astrophysical Journal*, 572, 540. <https://doi.org/10.1086/340017>
- Kahan, W. 2000, Technical report, University of California, Berkeley
- Kipping, D. M. 2010, *Monthly Notices of the Royal Astronomical Society*, 408, 1758. <https://doi.org/10.1111/j.1365-2966.2010.17242.x>
- . 2013, *Monthly Notices of the Royal Astronomical Society*, 435, 2152. <https://doi.org/10.1093/mnras/stt1435>
- Kopal, Z. 1950, *Harvard College Observatory Circular*, 454, 1
- Kreidberg, L. 2015, *PASP*, 127, 1161
- Lamarque, F., & Leroy, C. 1990, *Computer Physics Communications*, 59, 359. [https://doi.org/10.1016/0010-4655\(90\)90184-3](https://doi.org/10.1016/0010-4655(90)90184-3)
- Luger, R., et al. 2018, *ApJ*, XX, YY
- Magic, Z., et al. 2015, *Astronomy & Astrophysics*, 573, A90
- Mandel, K., & Agol, E. 2002, *ApJL*, 580, L171
- Maxted, P. F. L. 2018, ArXiv e-prints, arXiv:1804.07943
- Morello, G., et al. 2017, *The Astronomical Journal*, 154, 111. <https://doi.org/10.3847/1538-3881/aa8405>

- Neilson, H. R., & Lester, J. B. 2013, *Astronomy & Astrophysics*, 556, A86.
<https://doi.org/10.1051/0004-6361/201321888>
- Neilson, H. R., et al. 2017, *The Astrophysical Journal*, 845, 65.
<https://doi.org/10.3847/1538-4357/aa7edf>
- Ofir, A., et al. 2018, *The Astrophysical Journal Supplement Series*, 234, 9.
<https://doi.org/10.3847/1538-4365/aa9f2b>
- Pál, A. 2008, *Monthly Notices of the Royal Astronomical Society*, 390, 281.
<https://doi.org/10.1111/j.1365-2966.2008.13723.x>
- Pál, A. 2012, *MNRAS*, 420, 1630
- Parviainen, H., & Aigrain, S. 2015, *Monthly Notices of the Royal Astronomical Society*, 453, 3822.
<https://doi.org/10.1093/mnras/stv1857>
- Price, E. M., & Rogers, L. A. 2014, *The Astrophysical Journal*, 794, 92.
<https://doi.org/10.1088/0004-637x/794/1/92>
- Russell, H. N., & Shapley, H. 1912a, *The Astrophysical Journal*, 36, 239.
<https://doi.org/10.1086/141960>
- . 1912b, *The Astrophysical Journal*, 36, 385.
<https://doi.org/10.1086/141972>
- Sanchis-Ojeda, R., & Winn, J. N. 2011, *The Astrophysical Journal*, 743, 61.
<https://doi.org/10.1088/0004-637x/743/1/61>
- Seager, S., & Hui, L. 2002, *The Astrophysical Journal*, 574, 1004.
<https://doi.org/10.1086/340994>
- Sidis, O., & Sari, R. 2010, *The Astrophysical Journal*, 720, 904.
<https://doi.org/10.1088/0004-637x/720/1/904>
- Weisstein, E. W. 2018, "Circle-Circle Intersection." From *MathWorld* - A Wolfram Web Resource.,
<http://mathworld.wolfram.com/Circle-CircleIntersection.html>, , ,
 accessed 2018-07-09

APPENDIX

Here are a list of errata for [Mandel & Agol \(2002\)](#):

1. In Equation (7), λ_3 and λ_4 should have $2k \rightarrow 2p$ in arguments of the elliptic integrals.
2. In Equation (7), λ_5 should have $-\frac{2}{3}\Theta(p - 1/2)$ at the end.
3. For Case 11 in Table 1, η^d should be $1/2$, not 1 , and λ^d should be zero, not 1 . This mistake affects the code, but it is never encountered for planets that transit main-sequence stars since $p < 1$. This typo was discussed in [Eastman et al. \(2013\)](#).
4. The case $z = 1 - p$ is missing for $z < p$ (as pointed out by Pal 2008).
5. There is a π missing in the denominator of the second term on the right hand side of Equation (8).

With the exception of 3, none of these errors affected the publicly available code. Table 1 gives a list of the notation used throughout the main paper.

Table 1. Symbols used in this paper

Symbol	Definition	Reference
a_n	Giménez coefficients	Equation (2)
\mathbf{A}	Change of basis matrix: \mathbf{u} to Green's polynomials	Equation (45)
\mathbf{A}_1	Change of basis matrix: \mathbf{u} to polynomials	Equation (39)
\mathbf{A}_2	Change of basis matrix: polynomials to Green's polynomials	Equation (41)
A_{lens}	Lens-shaped area of overlap of two circles	Equation (3)
A_{kite}	Kite-shaped area b/w center of circles and points of contact	Equation (5)
b	Impact parameter in units of occulted body's radius	
b_c	Cutoff for using alternative expression for $d\mathcal{P}/db$	§5.4
$c_1 - c_4$	Non-linear limb darkening coefficients	§9
$\mathbf{D} \wedge$	Exterior derivative	Equation (50)
$\text{cel}(k_c, p, a, b)$	General complete elliptic integral (Bulirsch 1969)	Equation (14)
$E(\cdot)$	Complete elliptic integral of the second kind	Equation (13)
F	Normalized flux seen by observer	Equation (35)
\bar{F}	Time-averaged normalized flux	Equation (97)
${}_2F_1$	Generalized Hypergeometric function	Equation (70)
$\tilde{\mathbf{g}}$	Green's basis	Equation (42)
\mathbf{g}	Vector in the basis $\tilde{\mathbf{g}}$	
\mathbf{G}_n	Anti-exterior derivative of the n^{th} term in the Green's basis	Equation (49)
i	Dummy index	
I	Specific intensity, $I(x, y)$	
I_0	Intensity normalization constant	Equation (94)
j	Dummy index	
k	Elliptic parameter	Equation (12)
	Dummy index	
k_c	$\sqrt{1 - k^2}$	Equation (14)
$K(\cdot)$	Complete Elliptic integral of the first kind	Equation (13)
m_k	Elliptic integral parameter	§3
n	Order of limb darkening/Green's basis	
$\mathcal{M}_n(r, b)$	Integral computed recursively	Equation (61)
N	Highest order of limb darkening polynomial	
$\mathcal{N}_n(r, b)$	Integral computed recursively	Equation (74)
p	Coefficient of cel	Equation (14)
$\tilde{\mathbf{p}}$	Polynomial basis	Equation (37)
\mathbf{p}	Vector in the basis $\tilde{\mathbf{p}}$	
q	Term in cel identities	Equation (16)
\mathcal{P}	Primitive integral along perimeter of occulor	Equation (57)
\mathcal{Q}	Primitive integral along perimeter of occulted body	Equation (58)
r	Occulor radius in units of occulted body's radius	§1
\mathbf{r}	Vector for integration over boundary of visible disk	Equation (48)
\mathbf{s}	Occultation light curve solution vector	Equation (48)

Table 1 – continued from previous page

Symbol	Definition	Reference
t	Time variable	§6
t_0	Central time of transit	§6
u	Dummy index	
u_1, u_2	Quadratic limb darkening coefficients	Equation (1)
$\tilde{\mathbf{u}}$	Limb darkening basis	Equation (33)
\mathbf{u}	Vector of limb darkening coefficients in the basis $\tilde{\mathbf{u}}$	§5.1
\mathbf{x}	Parameters used in time integration	§6
x	Cartesian coordinate	Equation (32)
y	Cartesian coordinate	Equation (32)
z	Cartesian coordinate, $z = \sqrt{1 - x^2 - y^2}$	Equation (32)
α_j	Coefficient in series for \mathcal{M}_n	Equation (70)
γ_j	Coefficient in series for \mathcal{N}_n	Equation (78)
Γ	Gamma function	
η	Parameter in quadratic limb darkening term	Equation (25)
θ	Polar angle on star with respect to observer	
Θ	Heaviside step function	Equation (11)
κ_0	Angular position of occulter/occulted intersection point	Equation (4)
κ_1	Angular position of occulter/occulted intersection point	Equation (4)
λ	Angular position of occulter/occulted intersection point	Equation (55)
	Term in cel identities	Equation (16)
Λ	Mandel & Agol (2002) function	Equation (11)
μ	Cosine of polar angle on star, $\mu = z$	Equation (1)
$\Pi(\cdot, \cdot)$	Complete elliptic integral of the third kind	Equation (13)
ϕ	Angular position of occulter/occulted intersection point	Equation (54)
φ	Dummy integration variable	
ξ	Transformed integration variable	Equation (60)