

The background of the slide features a series of four circular diagrams arranged from left to right, illustrating the stages of a particle collision. The first circle shows two incoming particles (red and blue spheres) with arrows indicating their path. The second circle shows the particles interacting, with yellow wavy lines (gluons) appearing between them. The third circle shows a more complex interaction with multiple gluons and colored spheres. The fourth circle, on the right, shows a dense, tangled mass of yellow wavy lines, representing a high-energy state or a quark-gluon plasma. A large white arrow at the bottom points from left to right, labeled 'ENERGY'.

Understanding QCD dynamics through p-Pb collisions at the LHC

Allencris J.R.R.

Supervisor: Prof. Ronan McNulty

Image Credit: "Brookhaven Lab to Lead New 'Saturated Glue' Theory
Collaboration" , Brookhaven National Laboratory, 2022.
<https://www.bnl.gov/newsroom/news.php?a=120965>




Fig1. Proton parton distribution functions plotted as a function of Bjorken x




Fig2. A map plotting the log of $\frac{1}{x}$ against Q^2 . Different Evolution equations govern different regimes

Introduction

Cross-section: Probability that a given process takes place

Vector Mesons: Bound states of $q\bar{q}$ with the quantum numbers $J^P = 1^-$

Central Exclusive Production: Class of reactions $A + B \rightarrow A + X + B$
A & B remain intact. X is fully measured

Rapidity (y): Measure of particle angle with respect to beam axis
 $y \approx \eta$ for UR

Pseudorapidity (η): Rapidity but easier to measure




Fig3. CEP of the J/ψ meson in an ion-ion collision. Phys. Rev. C97 (2018) 024901




Fig4. Pseudorapidity range

(Albrow et al, 2010)




Fig5. The particular process(es) we're interested in.

Overview of Project Goals

Weizsacker-Williams
Approximation

Equivalent photon
spectrum

$$\sigma(\mathbf{h}_1 + \mathbf{h}_2 \rightarrow \mathbf{h}_1 \otimes V \otimes \mathbf{h}_2) = \int d\omega \frac{n_{h1}(\omega)}{\omega} \sigma_{\gamma h_2 \rightarrow V \otimes h_2}(W_{\gamma h_2}^2) + \int d\omega \frac{n_{h2}(\omega)}{\omega} \sigma_{\gamma h_1 \rightarrow V \otimes h_1}(W_{\gamma h_1}^2)$$

Color-dipole
formalism

Photoproduction
cross-section

(Goncalves et al, 2017)

Equivalent Photon Approximation (EPA)

1. Compute E-field of A_1 in rest frame of A_2
2. Fourier transform $E(t) \rightarrow E(\omega)$
3. $P_1 \rightarrow E_2$ and $P_2 \rightarrow E_1$
4. Integrate $P = P_1 + P_2$ over impact parameter b to get $n_A(\omega)$



$$n_A(\omega) = \frac{2Z_A^2 \alpha_{EM}}{\pi} \left[\zeta K_0(\zeta) K_1(\zeta) - \frac{\zeta^2}{2} (K_1^2(\zeta) - K_0^2(\zeta)) \right]$$




Fig6. Diagram showing equivalence of the E-field and a flux of photons


(Pshenichnov, 2011)

(Fermi, 1924)

(Williams, 1935)

Photoproduction cross-section

- Photon fluctuates into $q\bar{q}$
- $q\bar{q}$ interacts with particle through 2 gluons (unusual because colorless)



- Then creates/ moves off as V

$$\left. \frac{d\sigma(\gamma A \rightarrow J/\Psi)}{dt} \right|_{t=0} = \frac{\pi^3 \Gamma_{ee} M_{J/\Psi}^3}{48\alpha} \frac{\alpha_S^2(\bar{Q}^2)}{\bar{Q}^8} \times [x G_A(x, \bar{Q}^2)]^2$$

$$\sigma(\gamma A \rightarrow J/\Psi) = \left. \frac{d\sigma(\gamma A \rightarrow J/\Psi)}{dt} \right|_{t=0} \int_{t_{min}}^{\infty} dt |F(t)|^2$$

References

- [1] A. Accardi et al. Electron Ion Collider: The Next QCD Frontier - Understanding the glue that binds us all. 2012. DOI: 10.48550/ARXIV.1212.1701. URL: <https://arxiv.org/abs/1212.1701>.
- [2] M.G. Albrow, T.D. Coughlin, and J.R. Forshaw. "Central exclusive particle production at high energy hadron colliders". In: Progress in Particle and Nuclear Physics 65.2 (2010), pp. 149–184. DOI: 10.1016/j.ppnp.2010.06.001. URL: <https://doi.org/10.1016%2Fj.ppnp.2010.06.001>.
- [3] Carlos A. Bertulani, Spencer R. Klein, and Joakim Nystrand. "PHYSICS OF ULTRA-PERIPHERAL NUCLEAR COLLISIONS". In: Annual Review of Nuclear and Particle Science 55.1 (2005), pp. 271–310. DOI: 10.1146/annurev.nucl.55.090704.151526. URL: <https://doi.org/10.1146%2Fannurev.nucl.55.090704.151526>.
- [4] Enrico Fermi. "On the Theory of Collisions between Atoms and Electrically Charged Particles". In: Electromagnetic Probes of Fundamental Physics. WORLD SCIENTIFIC, 2003. DOI: 10.1142/9789812704214_0026. URL: https://doi.org/10.1142%2F9789812704214_0026.
- [5] V. P. Gonçalves and C. A. Bertulani. "Peripheral heavy ion collisions as a probe of the nuclear gluon distribution". In: Physical Review C 65.5 (2002). DOI: 10.1103/physrevc.65.054905. URL: <https://doi.org/10.1103%2Fphysrevc.65.054905>.
- [6] V. P. Gonçalves and M. V. T. Machado. "Nuclear exclusive vector meson photoproduction". In: The European Physical Journal C 38.3 (2004), pp. 319–328. DOI: 10.1140/epjc/s2004-02044-7. URL: <https://doi.org/10.1140%2Fepjc%2Fs2004-02044-7>.
- [7] V. P. Gonçalves et al. "Color dipole predictions for the exclusive vector meson photoproduction in pp/pP b/P bP b collisions at run 2 LHC energies". In: Physical Review D 96.9 (2017). DOI: 10.1103/physrevd.96.094027. URL: <https://doi.org/10.1103%2Fphysrevd.96.094027>.
- [8] Spencer R. Klein et al. "STARlight: A Monte Carlo simulation program for ultra-peripheral collisions of relativistic ions". In: Computer Physics Communications 212 (2017), pp. 258–268. DOI: 10.1016/j.cpc.2016.10.016. URL: <https://doi.org/10.1016%2Fj.cpc.2016.10.016>.
- [9] I. A. Pshenichnov. "Electromagnetic excitation and fragmentation of ultrarelativistic nuclei". In: Phys. Part. Nucl. 42 (2011), pp. 215–250. DOI: 10.1134/S1063779611020067.
- [10] E. J. Williams. "Correlation of certain collision problems with radiation theory". In: Kong. Dan. Vid. Sel. Mat. Fys. Med. 13N4.4 (1935), pp. 1–50.